


1955

# KILGORE'S

## Bred-Rite

### SEEDS



**Fordhook**  
242 Bush  
Lima Bean  
(See Page 8)

**U.S. No. 4 or 191**  
(White Kentucky Wonder)  
Pole Bean  
(See Page 7)



A field of  
**Contender**  
Beans at  
Pompano Beach,  
Florida  
(See Page 6)

**Early**  
**Blood-Red**  
**Market**  
Beet  
(Asgrow Wonder)  
(See Page 9)

**Isana**  
**Sweet Corn**  
(See Page 18)

**Blight**  
**Resistant**  
**World Beater**  
(Thick wall strain)  
Pepper  
(See Page 26)

**THE KILGORE SEED COMPANY**  
*General Offices and Mail Order Department, Plant City, Florida*






**Wilt Resistant Grothen Globe Tomato**  
(See Page 33)




**Great Lakes Lettuce**  
(See Page 21)



**Purple Top White Globe Turnip**  
(See Page 35)



**Early Yellow Summer Crookneck (Baby Crookneck) Squash**  
(See Page 30)



**Sunburst Mixture Zinnias**  
(See Page 48)

*Keep Up the Home Front  
and  
Make Florida More Beautiful*

## *Do It With Flowers*

Kilgore's flower seeds have been especially developed and improved for Florida planting. We list only kinds and varieties which, as the result of over forty years' experience, we have found to be especially suited to the climatic and soil conditions prevailing in Florida. Different kinds of flowers can be grown in Florida every month of the year. (See page 48 for list of flower varieties.)

Flower culture and home beautification in Florida are fully discussed in "Kilgore's Flower Guide for Florida." Send for your free copy.



## PLANT CITY, FLORIDA

Exp. or Frt. Office.....Date.....19.....

|            |  |  |
|------------|--|--|
| Total..... |  |  |
|------------|--|--|

No legal action can be brought on any defect or variation resulting from the failure of any seed to be as represented unless notice is given within a reasonable time, and not later than 15 days from the time such defect or variation becomes apparent and is observed by the purchaser. Chapter 26814 Laws of Florida, Acts of 1951. THE KILGORE SEED COMPANY.

**SUBSTITUTES:**

☐ —Please do not substitute without first notifying me.

We would like to send our annual catalog to your friends. Space on the reverse side of this sheet is provided for this purpose.

If you are interested in flower culture and home beautification and wish a copy of Kilgore's Flower Guide for Florida be sure to mark X in this space ☐

[SEE OTHER SIDE]



Tear Off Along This Line

PLEASE WRITE BELOW

| NAMES | POST OFFICE | P.O. Box, Street, or R.F.D. | STATE |
|-------|-------------|-----------------------------|-------|
|       |             |                             |       |
|       |             |                             |       |
|       |             |                             |       |
|       |             |                             |       |
|       |             |                             |       |



# Important Directions to Customers

By carefully observing the following directions, mistakes and misunderstandings will be very largely avoided.

**How to order**—Please be careful to sign your name, box number, street or rural route, post office and state on every order. Be sure to tell us your nearest express or freight office should you desire the seed or merchandise forwarded by express, freight or truck line. **Please write plainly.**

**Cash with order or C. O. D.**—Ours is a cash business and our prices are made on that basis. Customers are requested to remit when ordering, and by so doing save the c.o.d. fee charged by express companies, banks, and postal authorities for collecting and returning money.

**Perishable items not sent C. O. D.**—Perishable items, such as strawberry plants, rose bushes, seed potatoes, onion sets, etc., are not sent c.o.d. under any circumstances. Remittances must be sent with order.

**How to send remittance**—Remittances should be made by postal money orders, drafts, checks or cash. **Please do not send stamps.**

**We pay postage to points in Florida**—The postpaid prices indicated in this catalog include delivery to any point in Florida.

**Prices subject to change**—This catalog is printed during the early fall months. The market on many items of garden and field seeds, as well as supplies, has not yet been fully established. Consequently, prices on these items indicate value at time these pages are printed. In practically all cases orders will be filled at prices indicated in this catalog, **but all prices are subject to change without notice.** Write for special prices on larger quantities.

**Export Shipments to Cuba and the West Indies**—For the convenience of our customers in the West Indies, we have established the following agencies: E. G. Del Campo, Nueva Gerona, Isle of Pines, Cuba; F. C. Albury, Nassau, Bahamas.

Each order of pepper and tomato seed sent to Cuba requires an export certificate costing \$3.30.

*For direct deliveries outside the continental United States, all prices quoted as "Postpaid" in this catalog are on an f.o.b. Plant City, Florida, basis, and the proper amount for postage must be added to the cost of the seeds.*

**Parcel Post rates to Cuba and the West Indies**—The parcel post rate to Cuba, Bahamas and Jamaica is 45 cents, for Haiti 61 cents, and for the Dominican Republic 58 cents for the first pound or fraction thereof. If the net weight of an order is 1 pound or more, 1 pound must always be added to cover weight after being packed for shipment. In each case add 22 cents per pound for each additional pound or fraction thereof to the above indicated pound rates. Packages of merchandise going to Cuba, Haiti or Dominican Republic, weighing 8 ounces or less are subject to the postage rate of 1½ cents for each 2 ounces or fraction of 2 ounces. To the Bahamas and Jamaica, packages weighing 8 ounces or less are subject to the postage rate of 3 cents for each 2 ounces or fraction thereof, with a minimum charge of 15 cents.

**WARRANTY AND LIMITATION OF LIABILITY**—"We warrant that the seeds, bulbs and plants we sell are, at the time of delivery, as described on the container within recognized tolerances; but we limit our liability on this warranty to the amount of the purchase price of such seeds, bulbs and plants. We give no other or further warranty express or implied." The Kilgore Seed Company.

No legal action can be brought on any defect or variation resulting from the failure of any seed to be as represented unless notice is given within a reasonable time, and not later than 15 days from the time such defect or variation becomes apparent and is observed by the purchaser. Chapter 26814 Laws of Florida, Acts of 1951. THE KILGORE SEED COMPANY.

Air express shipments can now be made from Miami or Tampa, Florida, to any point in the West Indies.

**Return of seeds for exchange**—To safeguard our customers and ourselves, seeds may not be returned if seed package has been broken. In case of error on our part, unbroken packages may be returned within 10 days after shipment and we will gladly make correction at our expense.

**Stock or lot number on all Kilgore's seed packages**—Every lot of seed distributed by the Kilgore Seed Company will indicate (on one end of the front of the envelope or on the back of the label on canned seed, or with large seed like beans, peas, and corn on a tag attached to the bag), a stock number or lot number. This stock number is a code reference to our records of the history, performance and production of that particular lot of seed, by means of which we can trace it back to the growers' field where it was produced and to our trial plantings. It is recommended that every commercial grower of vegetables retain the empty packages or tags with the stock numbers. Correspondence pertaining to the performance of Kilgore's Bred-Rite seeds should include the specific lot number of the seed under consideration to aid investigation.

**Postal, Express, and Sales Tax information**—The parcel post weight limit for first and second zones is 40 pounds; 20 pounds in the third to eighth zones. Any weight up to 40 pounds in the first and second zones and 20 pounds in the third zone will usually go just as cheaply by parcel post as by freight or express.

In sections of Florida that are not easily accessible by freight or express, parcel post service will be found not only more satisfactory, but more economical. If on a rural route, delivery will be made to your door by mail.

Most Florida points are in the 1st (within 50 miles from Plant City, Fla.) or 2nd (within 50 to 150 miles from Plant City, Fla.) Postal Zone. Jacksonville north, Live Oak north and west, and points on the east coast south of West Palm Beach are in the 3rd Zone. (Within 150 to 300 miles from Plant City, Fla.) Puerto Rico and Virgin Islands are in the 6th Zone for surface mail; 8th Zone for Air Parcel Post. Surface mail 28c for 1st lb., 12c for each additional lb. Air Parcel Post 80c lb.

| 1st and 2nd ZONES |     |         |     | 3rd ZONE |     |         |      |
|-------------------|-----|---------|-----|----------|-----|---------|------|
| 1 lb.             | .23 | 11 lbs. | .63 | 1 lb.    | .23 | 11 lbs. | .75  |
| 2 lbs.            | .27 | 12 "    | .67 | 2 lbs.   | .29 | 12 "    | .80  |
| 3 "               | .31 | 13 "    | .71 | 3 "      | .34 | 13 "    | .85  |
| 4 "               | .35 | 14 "    | .75 | 4 "      | .39 | 14 "    | .90  |
| 5 "               | .39 | 15 "    | .79 | 5 "      | .44 | 15 "    | .96  |
| 6 "               | .43 | 16 "    | .83 | 6 "      | .49 | 16 "    | 1.01 |
| 7 "               | .47 | 17 "    | .87 | 7 "      | .54 | 17 "    | 1.06 |
| 8 "               | .51 | 18 "    | .91 | 8 "      | .60 | 18 "    | 1.11 |
| 9 "               | .55 | 19 "    | .95 | 9 "      | .65 | 19 "    | 1.16 |
| 10 "              | .59 | 20 "    | .99 | 10 "     | .70 | 20 "    | 1.21 |

If in doubt about your zone rate, obtain it from your postmaster, and be sure to send enough money to cover necessary postage, if the items ordered are not indicated "postpaid" in this catalog. Do not have your order delayed or reduced by failing to add necessary money for postage. If you send too much money, it will be returned to you. It will be found plainly stated above each item whether or not it is postpaid. Also be sure to include with your remittance the State Sales Tax on non-exempt merchandise. (See Page 2).

When you order your seeds and supplies to come by express or freight we can send them collect, so you may pay the transportation charges on delivery. Give your express or freight station, if it is not the same as your post office.



# SALES TAX INFORMATION

The Florida Revenue Act of 1949 (Sales and Use Tax) imposes a 3% tax on the retail sales of many commodities. Numerous agricultural supply items are exempt from the tax, and for your guidance in determining tax exempt items shown in this catalog, we list below those items that are not subject to the tax.

**EXEMPT ITEMS**—Beekeeper's Supplies (except Hive Bodies and Frame).

|                        |                       |
|------------------------|-----------------------|
| Suppers                | Seeds—All Vegetable,  |
| Cans and Jars          | Field Crop, Pasture   |
| Crate Materials        | Grass Seeds, Onion    |
| Fencing—For farm use   | Sets, Seed Potatoes   |
| Feeds—For farm animals | Seedbed Cloth         |
| Grafting Wax and       | Seed Inoculants       |
| Rubber Strips          | Seed Treating         |
| Hotkaps                | Compounds             |
| Label Paste            | Starter Solutions and |
| Peat Moss              | Root Inducing         |
| Poultry and Stock      | Materials             |
| Remedies               | Terra-Lite            |
| Paper Pots—Plant Bands | Tractors              |
| and Flats              | Weed Killers          |

**EXEMPT ONLY IF USED FOR COMMERCIAL AGRICULTURAL PURPOSES**

|                     |                  |
|---------------------|------------------|
| Dusters             | Fumigants        |
| Farm Tools          | Garden Tools     |
| Fertilizers         | Raffia and Twine |
| Insecticides and    | Rubber Bands     |
| Fungicides          | Sprayers         |
| Soluble Plant Foods | Tarpaulins       |

All flower seeds and other merchandise listed in this catalog is subject to 3% State Sales Tax except sales to Dealers, Government Agencies, Schools, Churches and Charitable organizations provided Tax Exemption Certificate has been obtained from the State Comptroller. All orders from institutions or organizations claiming tax exemption must clearly indicate on order sent us, the certificate serial number and date of certificate.

**THE RATE OF TAX APPLICABLE TO NON-EXEMPT SALES IS AS FOLLOWS:**

- (1) On single sales of 10c or less no amount shall be added for taxes.
- (2) On single sales in amounts for 11c to 35c, both inclusive, 1c shall be added for taxes.
- (3) On sales in amounts from 35c to 65c, both inclusive, 2c shall be added for taxes.
- (4) On sales in amounts from 65c to \$1.00, both inclusive, 3c shall be added for taxes.
- (5) On sales in amounts of more than \$1.00, 3% shall be charged upon each dollar of price, plus the above bracket charges upon any fractional part of a dollar in excess of even dollars.

**SPECIAL NOTICE**—In order to avoid delay in filling your order, on all non-exempt items, be sure to include with your remittance the proper amount of sales tax as indicated above. We are not permitted by law to assume the Sales Tax.

## FLORIDA HOME VEGETABLE GARDENS IN 1955

Florida is one of the most important commercial vegetable producing states of the Union. Florida home vegetable gardens offer almost no competition to commercial vegetables which are grown mainly for shipment to northern markets. Many Florida families who do not get fresh vegetables from their own gardens, seldom get them, at least in sufficient quantity for good health, from the market. Florida people, city residents as well as rural folks, can help solve their food problems by producing vegetables in the home garden.

Florida climate and soils are favorable for producing the finest vegetables in the United States. In Flor-

ida fresh vegetables can be grown every month in the year, especially during the fall, winter and spring months when they cannot be grown in other sections of the United States. In addition to producing fresh vegetables in your Florida home garden, plans should also be made for preserving vegetables from the garden by canning or freezing for home use, which will mean not only better health for the family, but also a material saving in cost. For best results, order your seeds from the Kilgore Seed Company — Florida's Leading Seedsmen, Mail Order Department, Plant City, Florida.





## Location and plan of the vegetable garden

A level, well drained piece of ground, convenient to the house should be selected for the garden. It should be located where it can be irrigated if necessary and away from hedges, large trees or the shade of buildings. On the farm it should be well-fenced to protect it from rabbits, chickens and other animals.

After selection of a location, and determining the size of the garden, a list of vegetables should be made. In selecting varieties, primary consideration should be given to the likes of the family, also to the size of the plot available. Different vegetables give variety to the diet.

A map or plan of the garden should be made, showing the various vegetables to be grown, and their position in the garden, indicating length of rows and the

space between rows. In the case of a small, backyard garden, entirely worked by hand, the rows should be close together, i.e., the smaller figure in column 5 of our chart on page 4, so that large quantities of vegetables may be produced on a limited space. In large farm gardens cultivated by mule-drawn tools, or other equipment used in the fields for this purpose, the rows should run the long way of the plot, and all crops should be in wide rows as indicated by the larger figure in the chart on page 4.

The planting data given on page 4 of this catalog will help greatly in preparing a plan since seed requirements, planting depths, distances and time of maturity are given.

## Making a seedbed

Small seeds such as lettuce, cabbage, collard, pepper, eggplant, tomato, etc., should be sown in a seedbed and the young plants transplanted to the field or garden. A shallow box or flat may be used for this purpose, or a small plot in the garden may be set aside to use as a seedbed for starting plants for transplanting. The soil must be in perfectly fine condition, and in the seedbed it should be made into beds four to six inches high, to prevent overflow in case of heavy rains. In preparing seedbed soil it is desirable to mix in "Terra-Lite" (Vermiculite) in order to increase germination of seed and prevent damping off of seedlings. (See page 74). When the soil in the seedbed is in condition—thoroughly and finely pulverized, fertilized with 2 to 3 pounds per 100 sq. ft. of a commercial fertilizer such as Vigoro or Vertagreen (page 67), applied at least a week or ten days before sowing seed, moist, and made smooth and level—sow the seed in rows about six inches apart. Very fine seed like celery, should not be covered with soil, but may be pressed into the soil with a light roller or with a wide board. Larger seeds like tomato, pepper, and cabbage may be covered, not over  $\frac{1}{4}$  to  $\frac{1}{2}$  inch deep.

We recommend the use of burlap or white muslin as a cover over the seedbed, stretching it on a three- or four-foot wide tent-shaped or slanting frame built over the seedbed, with the sides about 10 to 12 inches from the ground to provide ample circulation of air under the cover. If plants are being started in late

summer for fall planting, the seedbed cover will serve as a shade to protect seedlings from the hot sun, also to break the force of heavy rains. If plants are started in winter for a spring crop, this seedbed cover will protect seedlings from frost or freezes. It may be also advisable to use ground covers of burlap or old fertilizer bags. When seeds begin to germinate these ground covers should be removed immediately to prevent leggy or spindly plants. After the seed is sown, it is desirable to sprinkle every evening or oftener if necessary with cool water by means of hand sprinkling pots in order to keep the soil cool and moist, which tends to increase germination of seeds during the hot summer months.

Success in getting transplanted plants to grow satisfactorily depends very much upon how the plants are grown in the seedbed. For a week or ten days before transplanting, the plants in seedbed should be gradually hardened. This means leaving the cover off the plants for a week or so before transplanting, and water should be withheld from the plants during this period. When plants in seedbed are about six inches high they are ready to transplant into the field or garden, preferably on a cloudy day or toward evening. Water the plants thoroughly several hours before removing from seedbed, and again immediately after setting. A starter solution is helpful (see page 67). To grow good vegetables, a large supply of moisture and fertilizer is essential until the crop is ready to harvest.

## Preparation of the soil for the vegetable garden

The soil in the garden should be carefully and well prepared. Grass and weeds should be removed with all roots. A little extra time spent in preparing the soil for planting will save many hours of hoeing later in the season. It is best to spade or plow, and apply commercial fertilizer to the soil ten days to two weeks before planting in order to avoid burning the seed or young plants. A sample of the soil should be tested, either by yourself with an inexpensive soil test kit (page 75) or by your county agent, and if found too acid or sour apply Hardwood Ashes or Hydrated Lime; if too alkaline or sweet apply Aluminum Sulphate and Manganese Sulphate (see page 68).

Well rotted animal manure may be used as a fertilizer, at the rate of approximately 25 to 30 lbs. of manure, and  $2\frac{1}{2}$  to 3 lbs. of Super Phosphate for each 100 square feet of garden. If manure is not available, leaf mold or black muck or hammock soil are ideal materials to work into the garden soil. If none of these materials is available, a two or three inch layer of horticultural grade peat moss (page 74) may be worked into the garden soil. This material is clean and easy to handle, and is wonderfully effective for breaking up hard soil, making it mellow, or for adding humus to sandy soil, making it more retentive of moisture and fertilizer materials. After this peat moss is worked

into the soil, an inch or two inch layer of Pulverized Sheep Manure, along with a liberal application of a good garden fertilizer such as Vigoro or Vertagreen (see page 67) should be used, applying one of these fertilizer materials broadcast at the rate of 4 to 6 lbs. per 100 square feet. If this material is raked into the surface soil a week or ten days before planting, and if well watered in case of no rain in order to speed the incorporation and decomposition of the fertilizer or manure, you will have the foundation for an excellent vegetable garden. If the fertilizers are applied just before planting, they are very apt to burn the young seedlings as they emerge from the seed.

After the vegetable plants are well started in the garden, fertilizer should be applied lightly on either side of the row of vegetables, but not in contact with the leaves or stems to avoid burning, every two or three weeks. This will make thrifty plants which produce an abundant supply of high quality vegetables for the home table.

*Note:—All materials indicated above, such as Pulverized Sheep Manure, Peat Moss, Terra-Lite, Super Phosphate, Vigoro, Vertagreen, Wood Ashes, Lime, Aluminum Sulphate, Manganese Sulphate, etc., are available at all Kilgore seed stores, and in our mail order department at Plant City, Florida.*

For further information on Florida home gardening write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida, for a free copy of Bulletin 131 entitled "The Florida Home Garden," and also for Circular 104 entitled "Vegetable Garden Production Guide."



# Reference Table for Florida Vegetable Gardens

Many mistakes may be avoided by following directions given in this table. The time requirements indicated for crops to reach edible size vary, of course, according to the season, varieties and soil conditions. Allowance also must be made for the relative estimate of the yields, due to climatic and soil conditions. However, the aver-

ages indicated in the table will be found to be fairly accurate for Florida growing conditions on the basis of many years of observation and experience. **For information on "when to sow vegetable seeds in Florida" see page 5.**

| Kind of vegetable<br>Months<br>to<br>Sow Seed<br>(Inclusive)** | Number<br>Running feet<br>of row for<br>a family of<br>3 or 4 people<br>(Includes for<br>canning) | Seeds or plants<br>to provide for<br>a family of<br>3 or 4 people | Depth to<br>cover<br>in<br>inches | Space<br>between<br>rows<br>in<br>feet | Set plants from<br>seedbed or thin<br>out to inches<br>in row | Approximate time<br>required to produce<br>crop, from seed<br>planted in garden or<br>from transplanting<br>plants to garden<br>(days) |
|--|---|---|-----------------------------------|--|---|--|
| Beans (Bush) 1-4, 9-10   | 100*  | 1/2 lb.†  | 1 1/2-2                           | 2-2 1/2                                | 3-4   | 48 to 56   |
| Beans (Pole) 3-5   | 50  | 1/4 lb.   | 1 1/2-2                           | 3-4                                    | 9 or 3 ft. hills  | 63 to 70   |
| Beans (Bush Lima) 3  | 100*  | 1/2 to 1 lb.†   | 1 1/2-2                           | 3                                      | 4-5   | 65 to 75   |
| Beans (Pole Lima) 3  | 50  | 1/2 lb.   | 1 1/2-2                           | 3-4                                    | 9 or 3 ft. hills  | 90   |
| Beet 1-3, 9-12   | 100*  | 1 oz.†  | 1 1/2-3/4                         | 1 1/2-2                                | 3-4   | 56 to 68   |
| Broccoli 8-12  | 40  | 1 pkt. or 25 plts.  | 1/4-1/2                           | 3                                      | 18-24   | 55 to 60 from plts.  |
| Brussels Sprouts 1-2, 8-12                                     | 25  | 15 plants   | 1/4-1/2                           | 2 1/2-3                                | 18-24   | 90 from plants   |
| Cabbage 1-3, 8-12  | 60  | 50 plants   | 1/4-1/2                           | 2 1/2-3                                | 12-16   | 62 to 98 from plts.  |
| Chinese Cabbage 1-2, 8-12                                      | 40*   | 1 pkt. or 50 plts.†   | 1/4-1/2                           | 2 1/2-3                                | 8-12  | 70 from seed   |
| Cantaloupe 1-4   | 50  | 1 pkt.†   | 1/2-3/4                           | 4-5                                    | 24 or 4 ft. hills   | 75 to 95   |
| Carrot 1-3, 9-12   | 100*  | 2 pkts.†  | 1/4-1/2                           | 1 1/2-2                                | 2-3   | 72 to 95   |
| Cauliflower 1-2, 8-12  | 50  | 30 plants   | 1/4-1/2                           | 3-4                                    | 18-24   | 55 to 62 from plts.  |
| Celery 7-12  | 50  | 150 plants  | 1/8                               | 3                                      | 4-5   | 105 to 130 from plts.  |
| Collard Every Month  | 50*   | 1 pkt. or 35 plts.†   | 1/4-1/2                           | 2 1/2-3                                | 14-18   | 50 to 52 from plts.  |
| Corn (Sweet) 1-3   | 200*  | 1/4 lb.†  | 1-2                               | 2 1/2-3                                | 12-15   | 65 to 80   |
| Corn (Roasting Ear) 1-3  | 200*  | 1/4 lb.†  | 1-2                               | 3-3 1/2                                | 15-18   | 68 to 87   |
| Cowpeas (Table) 3-8  | 150   | 1/2 to 3/4 lb.  | 1-1 1/2                           | 3-4                                    | 5-6   | 68 to 85   |
| Cucumber 1-3, 8-9  | 50  | 1 pkt.  | 3/4                               | 5-6                                    | 12-24   | 40 to 50   |
| Eggplant 1-3, 6-8  | 50  | 17 plants   | 1/2                               | 4-4 1/2                                | 36-18   | 80 to 85 from plts.  |
| Endive (Escarolle) 1-2, 8-12                                   | 40  | 1 pkt. or 50 plts.  | 1/4-1/2                           | 1 1/2-2                                | 9   | 90 to 95 from seed   |
| Garlic 8-11 (bulbs or sets)                                    | 30  | 1 lb.   | 1                                 | 1 1/2                                  | 4   | 90   |
| Kale 1-3, 8-12   | 25  | 1 pkt.  | 1/4-1/2                           | 2 1/2                                  | 20-24   | 70   |
| Kohl Rabi 1-3, 9-12  | 25*   | 1 pkt.†   | 1/4-1/2                           | 1 1/2-2                                | 6-8   | 50   |
| Leek 1-2, 9-12   | 50  | 1 pkt. or 120 plts.   | 1/2                               | 1 1/2                                  | 5-6   | 140 from seed  |
| Lettuce (Head) 1-2, 8-12                                       | 75*   | 1 pkt. or 90 plts.†   | 1/4-1/2                           | 1 1/2-2                                | 10-15   | 66 to 84 from seed   |
| Lettuce (Leaf) 1-3, 8-12                                       | 50*   | 1 pkt. or 60 plts.†   | 1/4-1/2                           | 1 1/2-2                                | 10-12   | 46 from seed   |
| Muskmelon 1-4  | 50  | 1 pkt.  | 1/2-3/4                           | 4-5                                    | 24 or 4 ft. hills   | 75 to 95   |
| Mustard 1-12   | 40*   | 1 pkt.†   | 1/4-1/2                           | 2                                      | 8-10  | 40 to 45   |
| Okra 3-9   | 75  | 1 oz.   | 1-1 1/2                           | 3-4                                    | 10-15   | 50 to 60   |
| Onion Seed 1-9, 9-12   | 80  | 1 pkt. or 400 plts.   | 1/2                               | 1 1/2                                  | 2-4   | 110 to 150 from seed   |
| Onion Sets 1-2, 8-12   | 40  | 1 pint.   | 1                                 | 1 1/4-1 1/2                            | 1-2   | 30 to 60   |
| Parsley 1-2, 9-12  | 30  | 1 pkt.  | 1/4-1/2                           | 1 1/2-2                                | 10-12   | 90   |
| Parsnip 1-2, 9-12  | 50  | 1 pkt.  | 1/2                               | 2 1/2-3                                | 4   | 120  |
| Peas (English) 1-2, 9-12                                       | 100*  | 1/2 lb.†  | 1 1/2-2                           | 3                                      | 1-2   | 50 to 53   |
| Pepper 1-2, 6-8, 11-12   | 40  | 24 plants   | 1/2                               | 2 1/2-3 1/2                            | 10-16   | 60 to 80 from plts.  |
| Potatoes (Irish) 1-3, 8-12                                     | 200   | 12 lbs.   | 1-4                               | 3-3 1/2                                | 9-12  | 75 to 90   |
| Pumpkin 3-5  | 50  | 1/2 oz.   | 1                                 | 8-10                                   | 5 ft. hills   | 100 to 118   |
| Radish 1-4, 8-12   | 25*   | 1 pkt.†   | 1/2                               | 1-1 1/2                                | 1-2   | 21 to 25   |
| Rhubarb 9-11   | 25  | pkt. or 15 plts.  | 1/2-3/4                           | 2 1/2-3                                | 18-24   | 90 from plants   |
| Roselle 3-4  | 25  | 1 pkt.  | 1/2-3/4                           | 5-6                                    | 24-36   | 175  |
| Rutabaga 1-2, 8-12   | 50  | 1 pkt.  | 1/4-1/2                           | 2                                      | 5-6   | 90   |
| Spinach-Savoy 1-2, 8-12  | 50*   | 1 pkt.†   | 1/2-3/4                           | 1 1/4-1 1/2                            | 6   | 40   |
| Spinach-New Zealand 3-7  | 25  | 1 pkt.  | 3/4-1                             | 3                                      | 20  | 60   |
| Squash (Bush) 1-3, 8-9   | 50  | 1 pkt.  | 1                                 | 3-4                                    | 2 ft. hills   | 42 to 52   |
| Squash (Running) 1-2   | 50  | 1 pkt.  | 1                                 | 6-8                                    | 3 or 4 ft. hills  | 60 to 115  |
| Swiss Chard 1-3, 9-12  | 25  | 1 pkt.  | 1/2-1                             | 2                                      | 12-15   | 58 to 60   |
| Tomato-Ground 1-3, 8-9, 12                                     | 150   | 50 plants   | 1/2                               | 3 1/2-5                                | 30-36   | 70 to 90 from plts.  |
| Tomato-Staked 1-3, 8-9, 12                                     | 75  | 50 plants   | 1/2                               | 4                                      | 16-20   | 70 to 90 from plts.  |
| Turnip 1-4, 7-12   | 50  | 1 pkt.†   | 1/4-1/2                           | 1 1/4-1 1/2                            | 4-6   | 35 to 50   |
| Watermelon 1-4   | 75  | 1 pkt.  | 1/2-1                             | 8-10                                   | 6 to 8 ft. hills  | 85 to 100  |

\* Make two or more plantings at different times during the season. Number of feet of row indicated is for each planting.

† Number of feet and amount of seed or number of plants indicated are for each planting.

\*\* Months indicated are for central and south Florida. In north Florida, some tender kinds will not stand the winter cold, and this fact should be taken into consideration with reference to time of sowing seed.

## THE KILGORE SEED COMPANY—A Florida Institution GENERAL OFFICES AND MAIL ORDER DEPARTMENT, PLANT CITY, FLORIDA

Fourteen Kilgore seed stores serving Florida, located at:

Belle Glade  
Fort Myers  
Gainesville  
Homestead

Miami —  
600 N.W. Fifth St.  
Ocala  
Pahokee

Palmetto  
Plant City  
Pompano Beach  
Sanford

Vero Beach  
Wauchula  
West Palm Beach —  
1310 Belvedere Road



# What Vegetables and When to Sow Seeds in Florida

**JANUARY AND FEBRUARY**—Bush beans, Beet, Brussels Sprouts, Cabbage, Chinese Cabbage, Cantaloupe, Carrot, Cauliflower, Collard, Corn—Sweet and Roasting Ear, Cucumber, Eggplant, Endive, Kale, Kohl Rabi, Leek, Lettuce, Mustard, Onion Seed and Sets, Parsley, Parsnip, Peas—English or Garden, Pepper, Potatoes, Radish, Romaine, Rutabaga, Spinach, Squash, Swiss Chard, Tomato, Turnip, Watermelon.

**MARCH**—Bush, Pole and Lima Beans, Beet, Cabbage, Cantaloupe, Carrot, Collard, Corn—Sweet and Roasting Ear, Cucumbers, Eggplant, Kale, Kohl Rabi, Lettuce, Mustard, Okra, Potatoes, Pumpkin, Radish, Roselle, Spinach—New Zealand, Squash, Swiss Chard, Tomato, Turnip, Watermelon.

**APRIL**—Bush and Pole Beans, Cantaloupe, Collard, Mustard, Okra, Pumpkin, Radish, Roselle, Spinach—New Zealand, Turnip, Watermelon.

**MAY**—Pole Beans, Collard, Mustard, Okra, Pumpkin, Spinach—New Zealand.

**JUNE**—Collard, Eggplant, Mustard, Okra, Pepper, Rutabaga, Spinach—New Zealand, Turnip.

**JULY**—Celery, Collard, Eggplant, Mustard, Okra, Pepper, Rutabaga, Spinach—New Zealand, Turnip.

**AUGUST**—Broccoli, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Celery, Collard, Cucumber, Eggplant, Endive,

Garlic, Kale, Lettuce, Mustard, Okra, Onion Sets, Pepper, Radish, Romaine, Rutabaga, Spinach, Squash, Tomato, Turnip.

**SEPTEMBER**—Bush Beans, Beet, Broccoli, Brussels Sprouts, Cabbage, Chinese Cabbage, Carrot, Cauliflower, Celery, Collard, Cucumber, Endive, Garlic, Kale, Kohl Rabi, Leek, Lettuce, Mustard, Okra, Onion Seed and Sets, Parsley, Parsnip, Peas—English or Garden, Radish, Rhubarb, Romaine, Rutabaga, Spinach, Squash, Swiss Chard, Tomato, Turnip.

**OCTOBER**—Bush Beans, Beet, Broccoli, Brussels Sprouts, Cabbage, Chinese Cabbage, Carrot, Cauliflower, Celery, Collard, Endive, Garlic, Kale, Kohl Rabi, Leek, Lettuce, Mustard, Onion Seed and Sets, Parsley, Parsnip, Peas—English or Garden, Potatoes, Radish, Rhubarb, Romaine, Rutabaga, Spinach, Swiss Chard, Turnip.

**NOVEMBER**—Beet, Broccoli, Brussels Sprouts, Cabbage, Chinese Cabbage, Carrot, Cauliflower, Celery, Collard, Eggplant, Endive, Garlic, Kale, Kohl Rabi, Leek, Lettuce, Mustard, Onion Seed and Sets, Parsley, Peas—English or Garden, Pepper, Potatoes, Radish, Rhubarb, Romaine, Rutabaga, Spinach, Swiss Chard, Turnip.

**DECEMBER**—Beet Broccoli, Brussels Sprouts, Cabbage, Chinese Cabbage, Carrot, Cauliflower, Celery, Collard, Eggplant, Endive, Kale, Kohl Rabi, Leek, Lettuce, Mustard, Onion Seed and Sets, Parsley, Parsnip, Peas—English or Garden, Pepper, Potatoes, Radish, Romaine, Rutabaga, Spinach, Swiss Chard, Tomato, Turnip.

## What grasses, clovers, and field crops, and when to sow seeds in Florida

**JANUARY AND FEBRUARY**—Alfalfa, Clover—Crimson, Grasses—Bahia, Oats, Rape, Rye.

**MARCH**—Beans—Velvet, Benne, Corn—Field, Roasting Ear, and Chicken, Crotalaria, Grasses—Bermuda, Carpet, Italian Rye, Buffel, Sudan and Bahia, Blanket Indigo, Hairy Indigo, Hegari, Millet, Peas—Southern or Cowpeas, Peanuts, Rape, Rice, Shallu (Egyptian Wheat), Sorghum, Sunflower.

**APRIL**—Beans—Velvet, Beggarweed, Benne, Chufas, Clover—Alyce, Corn—Field, Roasting Ear and Chicken, Crotalaria, Grasses—Bermuda, Carpet, Buffel, Sudan and Bahia, Blanket Indigo, Hairy Indigo, Hegari, Millet, Peas—Southern or Cowpeas, Peanuts, Rice, Shallu (Egyptian Wheat), Sorghum, Sunflower.

**MAY**—Beans—Velvet, Beggarweed, Benne, Chufas, Clover—Alyce, Corn—Field and Chicken, Crotalaria, Grasses—Bermuda, Carpet, Buffel, Sudan and Bahia, Blanket Indigo, Hairy Indigo, Hegari, Millet, Peas—Southern or Cowpeas, Peanuts, Sesbania, Shallu (Egyptian Wheat), Sorghum, Sunflower.

**JUNE**—Beans—Velvet, Beggarweed, Benne, Chufas, Clover —Alyce, Corn—Chicken, Cowpeas, Crotalaria, Grasses—Bermuda,

Carpet, Buffel, Sudan and Bahia, Blanket Indigo, Hairy Indigo, Hegari, Millet, Peanuts, Rice, Sesbania, Shallu (Egyptian Wheat), Sorghum, Sunflower.

**JULY**—Beans—Velvet, Beggarweed, Chufas, Corn—Chicken, Cowpeas, Grasses—Bermuda, Buffel, Sudan and Bahia, Hegari, Millet.

**AUGUST**—Cowpeas, Grasses—Bermuda, Buffel, Sudan and Bahia, Millet, Sunflower.

**SEPTEMBER**—Clover—Sweet, Kenland Red, Grasses—Bermuda, Bahia, Lupines, Rape, Rye.

**OCTOBER**—Clover—Sweet, Kenland Red, Grasses—Bermuda, Italian Rye, Carpet, Bahia, Lupines, Oats, Rape, Rye.

**NOVEMBER**—Clover—Crimson, Sweet, White Dutch, Ladino, Kenland Red, Grasses—Bermuda, Italian Rye, Carpet, Bahia, Lupines, Oats, Rape, Rye.

**DECEMBER**—Clover—Crimson, Sweet, White Dutch, Ladino, Kenland Red, Grasses—Italian Rye, Carpet, Bahia, Lupines, Oats, Rape, Rye.

See pages 38 to 47 for complete cultural directions on field corn, grasses, clovers and field crops in Florida.

## What flowers and when to sow seeds in Florida

| Kind of Flower—When to Sow Seeds*                  | Kind of Flower—When to Sow Seeds*                        | Kind of Flower—When to Sow Seeds*                |
|--|--|--|
| AGERATUM (Floss Flower)—Sept. through March        | CHINESE FORGET-ME-NOT OR CYNOGLOSSUM—Sept. through March | NASTURTIUM—Sept. through March                   |
| ALYSSUM—Sept. through Jan.                         | COSMOS—March through May                                 | PANSY—Sept. through Dec.                         |
| ASTER—Sept. through April                          | DAHLIA—March through April                               | PETUNIA—Sept. through June                       |
| BABY'S BREATH OR GYPSOPHILA—Sept. through Jan.     | DELPHINIUM (PERENNIAL LARKSPUR)—Oct. through Nov.        | PHLOX—Sept. through Feb.                         |
| BACHELOR'S BUTTON OR CENTAUREA—Sept. through Jan.  | FLOSS FLOWER OR AGERATUM—Sept. through March             | PINKS OR DIANTHUS—Sept. through March            |
| BALSAM—Sept. through March                         | GLOBE AMARANTH (GOMPHRENA)—Feb. through July             | QUEEN ANNE'S LACE—Sept. through Jan.             |
| BEAN—SCARLET RUNNER—Feb. through May               | GOULD (ORNAMENTAL)—Feb. through May                      | SALVIA—Sept. through June                        |
| GAILLARDIA OR BLANKET FLOWER—Any month of the year | HOLLYHOCK, ANNUAL INDIAN SPRING—Aug. through Jan.        | SHASTA DAISY—Sept. through Jan.                  |
| AFRICAN DAISY OR ARCTOTIS—Sept. through Jan.       | LARKSPUR—Nov. through Jan.                               | SNAPDRAGON OR ANTIRRHINUM—Sept. through Dec.     |
| BLUE LACE FLOWER OR DIDISCUS—Sept. through Jan.    | LINARIA—Sept. through Dec.                               | STATICE—Sept. through Jan.                       |
| CALENDULA—Sept. through Jan.                       | LUPINE—Sept. through Dec.                                | STRAWFLOWER OR HELICHRYSUM—Sept. through March   |
| CALIFORNIA POPPY—Sept. through Jan.                | MARIGOLD—Feb. through May and again in Sept.             | SUNFLOWER OR HELIANTHUS—Feb. through Aug.        |
| CALLIOPSIS OR COREOPSIS—Sept. through Feb.         | MORNING GLORY OR IMPROVED IPOMEA—March through June      | SUN PLANT OR PORTULACA—Feb. through Aug.         |
| CANDYTUFT—Sept. through Feb.                       | MOURNING BRIDE OR SCABIOSA—Sept. through Dec.            | SWEET PEA—Sept. through Dec.                     |
| CARNATION—Sept. through Dec.                       |  | TITHONIA OR FLORIDA SUNFLOWER—Feb. through June. |
| CELOSIA (COCKSCOMB)—Jan. through June              |  | VERBENA—Any month of the year                    |
|  |  | VINCA OR PERIWINKLE—Any month                    |
|  |  | ZINNIA—Feb. through Aug.                         |

See page 48 for complete list of flower varieties for Florida.

\* "Time to sow seed" indicated is for central and south Florida. In north Florida, many kinds will not withstand the winter cold, and this fact should be taken into consideration with reference to time of planting.

If interested in flowers write for "Kilgore's Flower Guide for Florida"



# BEANS

Our bean seed is high germinating, guaranteed western-grown stock, from the best western growing areas, all carefully milled, graded and fitted for the critical grower.

See page 34 for details of the Kilgore method of handling seeds

**Culture of Green and Wax Podded Bush Beans.** Beans are grown on all types of soil in Florida, and are not very particular with reference to soil requirements. Beans are a very quick maturing crop, and are usually preceded or followed by other crops.

Spring planting in the southern and central part of the State begins January 1 and continues until April. Fall planting starts in August. For a fall crop, plant from the middle of August until the middle of October; for a main crop on the Florida east coast plant from October to December. Sow seed in drills, 2 to 2½ feet apart, dropping a bean every 2 to 2½ inches, and cover 1½ to 2 inches deep. On muck soils apply 300 to 600 pounds of a 3-8-8 fertilizer mixture per acre. On sandy loam apply 1200 to 2000 pounds of a 4-7-5 fertilizer per acre, making one application. The fertilizer put out from a week to ten days before planting. There is always a risk of the fertilizer burning and killing the

vitality of the seed when both are put in at the same time.

For the control of insects and diseases of beans see page 53. In order to control soil diseases and get a better stand, it is always desirable to treat bean seed with **Spergon Seed Protectant** at the rate of 2 ounces per bushel of seed before planting. (See pages 52 and 62). This often means the difference between success and failure in getting a stand. For control of watery soft rot or white mold use Granular Aero Cyanamid.

Inoculate bean seed with Nitragin or A.B.C. Inoculant. This hastens maturity, increases yields and enriches the soil. (See page 69).

The number of days to picking after each variety represents average results secured over a period of years under Florida growing conditions, and will vary somewhat in different sections, seasons, and on different soils. The number of days indicated will serve to show the relative maturity of different varieties.

## Green Podded Bush Beans

Plant ½ lb. to 100 feet of row at each planting

Plant 3 pecks (45 lbs.) to 1 bushel (60 lbs.) per acre. In the garden make four plantings at different times.

Write for special prices on larger quantities



Contender (Mosaic Resistant) Bush Beans

**CONTENDER (Mosaic Resistant).** (45 days.) A consistently early, high yielding, attractive appearing, stringless bean, adapted for shipment to fresh markets as well as processing. Plant is very vigorous and produces pods similar to Black Valentine, although the Contender pods will average three-fourths of an inch longer, are slightly heavier and thicker, are round to oval, and have a tendency to curve slightly. Consistently an enormous yielder.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.10; 60 lbs. (bu.) \$15.30

**ASGROW BLACK VALENTINE.** (48 days.) The plant is large, vigorous, erect and very prolific. The oval pods are nearly straight, medium green, entirely stringless, and of fine quality and delicate flavor. They retain their color and texture in shipping long distances. Very attractive in appearance, of high edible quality and a good shipper.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.10; 60 lbs. (bu.) \$15.30

**TENDERLONG 15 (Improved Tendergreen Type).** (Mosaic Resistant). (50 days.) Developed and introduced for disease resistance; longer, straighter pods than Tendergreen. The plant is an erect, compact bush, holding the pods well off the ground. It is highly productive and resistant to common bean Mosaic and several other virus diseases, one form of powdery mildew; with considerable tolerance to Halo blight. The pods are uniformly round and thick, similar to Tendergreen, but

more attractive, being longer, averaging six inches, straighter, meaty, stringless and fiberless, of excellent flavor and quality for use fresh, frozen or canned, of very attractive medium dark green color.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.70; 60 lbs. (bu.) \$17.70

**TENDERGREEN.** (50 days.) This variety is popular due to its excellent shipping quality. Plants are large, erect, and thick-stemmed. The pods average about five inches in length, are attractive in appearance, being nearly straight, round, fleshy, dark green, entirely stringless, and of excellent quality.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.50; 60 lbs. (bu.) \$17.10

**WADE (B1515) (Mosaic Resistant)** (48 days.) The vigorous plants are tall and hold the pods well off the ground. Because of the heavy foliage, the pods do not wind-scar as readily as do other varieties of this type. It is a fleshy, round podded bean of the Tendergreen type, but being powdery mildew and mosaic resistant, it is a consistently higher yielder and produces longer, straighter pods than Tendergreen. The pods are of a dark green color and hold their green color exceptionally well in transit and on the market. Entirely stringless and little fiber. Good for market as well as for processing. Because of its long bearing season, high quality and disease resistance, it is highly recommended for home gardens and is a good shipper.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid

Not prepaid, 15 lbs. (pk.) \$4.85; 60 lbs. (bu.) \$18.30



Tenderlong 15

Green Podded Bush Beans (Continued on next page)



# BEANS

## Green Podded Bush Beans (Cont'd)

**PLENTIFUL.** (50 days.) Plants more spreading in growth than Bountiful, and should be given a little wider spacing in the row. More resistant to rust than Bountiful. The pods are longer, slightly thicker, and narrower than Bountiful, flat, straight, and of medium green color.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$4.10; 60 lbs. (bu.) \$15.30


## WAX PODDED BUSH BEANS

Plant 3 pecks (45 lbs.) to 1 bushel (60 lbs.) per acre.

In the garden make four plantings at different times.

Plant ½ lb. to 100 feet at each planting for a family of 3 or 4 people.

*Write for special prices on larger quantities.*



*Cherokee*

**CHEROKEE.** (50 days.) Plant large, vigorous, erect and prolific. Early and an enormous yielder of beautiful, attractive appearing, golden-yellow, long, thick oval pods, nearly straight, entirely stringless and of excellent quality. Considered the best wax bean for Florida and the South.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$4.25; 60 lbs. (bu.) \$15.90

## GREEN PODDED POLE BEANS

Plant 2 pecks (30 lbs.) per acre.

In garden ¼ pound to 50 feet

**Culture of Green Podded Pole Beans.** Pole beans are later in maturing than bush beans, but bear for a longer season and are extremely productive, yielding more than bush beans, and also produce bigger pods. They are somewhat more tender than bush beans if harvested young. For best results, pole beans should be supported by wires or by good strong poles to keep them off the ground. Poles should be 8 to 9 feet above ground, and set firmly in the ground, in rows 3 to 4 feet apart, and three feet apart in the row. Plant three to four seeds around each pole. When plants are well established, thin to two or three plants to a pole. A better arrangement would be to place three or four poles tent-fashion at 3 ft. spacings in the row, and then thin to one plant at a pole. In order to keep pole beans bearing over a long season, it is important to keep them closely picked. Inoculate bean seed with Nitragin or A.B.C. Inoculant

for best results. (See page 69.) Pole beans are more susceptible to mildew, rust and other foliage diseases than are bush beans, and therefore should be sprayed with Sulfocide or dusted with sulphur quite often. (See pages 53 and 62.)

*Write for special prices on larger quantities.*

**U. S. NO. 4 (No. 191) (WHITE KENTUCKY WONDER).** (63 days.) A high yielding type of white-seeded Kentucky Wonder, resistant to certain forms of rust. Plant is vigorous, with dark green leaves. The pods are dark green, oval, straight, long, averaging from 8 to 9 inches, and attractive in appearance. The pods are stringless in the marketable stage. Productive and very desirable for market or home use. Grown most extensively on the west coast of Florida for shipment to southern and northern markets.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$5.25; 60 lbs. (bu.) \$20.10

**KENTUCKY WONDER (OLD HOMESTEAD).** (65 days.) This is a popular, vigorous, hardy, climbing or pole bean, early and very productive over a long season. Pods are very long, ranging from eight to ten inches, curved, nearly round, with undulating surface. Though thick, they appear slender on account of their great length. Pods are meaty, very brittle, and of good quality if harvested young. The pods hang in large clusters from the top to the bottom of the vine. Our strain has been selected for productivity, earliness and freedom from flat pods.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$4.25; 60 lbs. (bu.) \$15.90

**McCASLAN.** (65 days.) A vigorous grower and heavy producer, unlike other varieties, producing from the ground up over a long season. The pods are medium green, eight inches long, large, flat, slightly curved, thick and meaty, stringless, brittle, fine grained, and of excellent quality if harvested young. Because of the high yield of large, attractive pods, this variety has become the leading pole bean on the lower east coast of Florida for shipment to northern markets.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$4.50; 60 lbs. (bu.) \$17.10



*McCaslan pole beans as grown from Kilgore's high vitality seed in the Homestead section.*



# LIMA BEANS

## BUSH LIMA BEANS

Fordhook—Plant 1 bushel (60 lbs.) per acre.

Henderson's—Plant 1 to 2 pecks (15 to 30 lbs.) per acre.


In the garden make two plantings at different times. With Fordhook plant 1 pound to 100 feet or with Henderson's plant ½ pound to 100 feet at each planting for a family 3 or 4 people.

**Culture of Bush Lima Beans.** In general, cultural requirements of Bush Lima Beans are similar to those given for green and wax podded bush beans, but they need more room (see page 7). Lima beans are not so hardy as string beans, and the ground should be warm when planting the seed. Fertilizer should be applied at least ten days before planting. In order to control soil diseases and get a better stand, it is always desirable to treat lima bean seed before planting (see pages 52 and 62). Plant in rows three feet apart, dropping a bean every two or two and one-half inches in the row, covering an inch deep. Thin to stand 4 to 5 inches in the row.

*Write for special prices on larger quantities.*

**FORDHOOK 242.** (72 days.) It is an excellent continuous bearer, producing a heavy early yield, and is especially suitable to Florida because of its ability to make a heavy set and yield under high temperature conditions. Numerous tests all over the south have shown that Fordhook 242 outyields regular Fordhook by an average of from 35 to 50 per cent in different seasons. The plants are as tall as regular Fordhook, grow more erect and have a more open appearance. Pods average 3 to 4 inches long by 1 and 1½ inches wide and about ½ inch thick, and contain 3 to 4 thick beans. Green shell beans are very slightly smaller than regular Fordhook, have light green seed coats, and are of delicious, finely flavored edible quality. If you have not been successful in growing regular Fordhook in Florida by all means try Fordhook 242.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$4.85; 60 lbs. (bu.) \$18.30



*Jackson Wonder (Old Florida Bush Speckled Butter Bean)*

**FORDHOOK-CONCENTRATED** (75 days.) This highly bred strain has been developed for concentration of pods on the plant, making it easy to harvest, and for concentration in time of reaching maturity, practically all the beans maturing at one time, making successive harvests unnecessary. The plants are stocky, without tendrils and very prolific. Pods are dark green, very large, with thick fibrous walls giving them body for long distance shipments. Pods contain 3 to 4 large, very thick, plump beans of excellent quality.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$4.85; 60 lbs. (bu.) \$18.30

**HENDERSON'S BUSH LIMA** (Baby Lima). (65 days.) An early variety, hardy and a sure cropper. The plants are small and very productive. The pods are small in size, very short and narrow, thin, flat, straight to slightly curved. The seeds are smaller and not as plump as Fordhook. This variety matures about ten days earlier than Fordhook. The plants are more prolific than Fordhook, but the seeds and pods are smaller.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$3.75; 60 lbs. (bu.) \$14.10

## Butter or Green Shell Beans

Used principally as Green Shell Beans, like Lima Beans

Old Florida Pole—Plant 1 peck (15 lbs.) per acre.

Jackson Wonder Bush—Plant 1 to 2 pecks per acre.

French's Horticultural—Plant 1 to 2 pecks per acre.

In the garden plant ¾ pound Bush to 100 feet,

½ pound Pole to 100 feet for a family of 3 or 4 people.

*Write for special prices on larger quantities.*

**FLORIDA POLE SPECKLED BUTTER BEAN** (Old Florida Pole). (75 days.) Culture similar to Green Podded Pole Beans (See page 7). Plant is tall, vigorous, and very prolific over a long bearing season. Pods, borne in clusters, are of medium size, flat, containing an average of 3 medium-sized, flat seeds. Seeds are of buff color spotted with reddish brown. A dependable, prolific variety for Florida.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$4.35; 60 lbs. (bu.) \$16.30

**JACKSON WONDER** (Old Florida Bush Speckled Butter Bean). (65 days.) Culture similar to Bush Lima Beans. This variety is similar to Old Florida Pole, but produces a bush or dwarf plant. Plant is small to medium in size, spreading, with semi-runners resembling somewhat Henderson's Bush Lima. Pods are flat, containing 3 or 4 beans. The seeds are small, flat and of light buff color. It is earlier than Old Florida Pole, very prolific, drouth-resistant, a sure cropper and of good quality.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$3.75; 60 lbs. (bu.) \$14.10

**FRENCH'S HORTICULTURAL** (68 days.) This is the most attractive bean of the speckled Cranberry type making many long pods (6 to 8 inches), straight, very heavily splashed with bright red or deep carmine, which gives them a very striking and attractive appearance. They grow in profusion on strong, erect, sturdy bush plants, which produce short runners. The shelled beans are pinkish buff color, mottled and streaked. Can be used as a dry shell or green shell bean. Of fine flavor.

¼ lb. 20c; 1 lb. 65c; 2 lbs. \$1.10, Postpaid  
Not prepaid, 15 lbs. (pk.) \$5.60; 60 lbs. (bu.) \$21.30



# BEET

Plant 6 pounds of seed per acre.

In the garden make three plantings at different times.

Plant 1 oz. to 100 feet of row at each planting for a family of 3 or 4 people.

**Culture.** Successful culture demands good soil, to which has been applied 600 to 1,200 pounds per acre of good standard vegetable fertilizer. Plant seeds in rows 1½ to 2 feet apart and when 3 inches high thin to 3 or 4 inches apart in the row. In Florida, sow from latter part of September to March.

Prices quoted are postpaid.

Write for special prices on larger quantities.

**ASGROW WONDER (Early Blood Red Market).** (56 days.) One of the earliest and most uniform beets ever introduced. Roots very attractive in appearance, slightly flattened globe, smooth, with small tap root, of dark red color outside. Flesh of deep dark red, tender, and excellent quality.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50  
5 to 25 lbs. \$1.40 per lb.

**CROSBY'S EGYPTIAN (Improved).** (60 days.) This standard variety produces beets of uniform flattened globe shape, very much flattened at the top, of deep red color outside, with a deep purple color with little zoning inside. Sweet and tender.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50  
5 to 25 lbs. \$1.40 per lb.

**DETROIT DARK RED.** (68 days.) A very desirable, globe-shaped beet, of dark red color, outside as well as inside, without white zoning; sweet and of excellent quality.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50  
5 to 25 lbs. \$1.40 per lb.

## SWISS CHARD

Plant 4 pounds per acre.

In the garden plant one packet to 25 feet of row

**Culture.** Plant like beets in the fall, winter, or spring, in rows 2 feet apart, and thin to 12 or 15 inches in the row.

**FORDHOOK GIANT.** (68 days.) The thick, fleshy, dark green leaves are much crinkled or savoyed, and the pearly white stems are very broad, thick, and fleshy. A very vigorous, hardy grower.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50  
5 to 25 lbs. \$1.40 per lb.

**RHUBARB CHARD.** (60 days.) Stalks of rhubarb ruby-red color, which reddish color extends out through the veins into the rich dark green, heavily crumpled leaves.

Pkt. 10c; oz. 30c; ¼ lb. 85c; 1 lb. \$3.00  
5 to 25 lbs. \$2.90 per lb.

## SUGAR AND STOCK BEET

**Culture.** Plant 4 to 6 pounds per acre in rows 2½ to 3 feet apart, and thin to 6 inches apart in the row.

**SUGAR BEET, KLEIN WANZLEBEN.** (85 days.) This variety is the most valuable for sugar making and stock feeding.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

**STOCK BEET, MAMMOTH LONG RED.** (80 days.) An enormous yielder and one of the very best for dairy-men.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

# BRUSSELS SPROUTS

Sow ¼ pound in seedbed to plant an acre.

In the garden sow 1 packet in seedbed.

Sow seed and cultivate exactly the same as cabbage.

**LONG ISLAND IMPROVED.** (90 days from setting out plants.) Plants grow to a height of two feet. Stems are thickly set with small, firm, cabbage-like heads, 1½ inches in diameter, maturing small heads successively. Cooked and served like cabbage.

Pkt. 10c; ½ oz. 25c; oz. 45c; ¼ lb. \$1.40; 1 lb. \$5.00  
5 to 25 lbs. \$4.90 per lb.



# BROCCOLI

Sow ¼ to ½ pound in seedbed to plant an acre.

In the garden 1 packet to 40 feet of row, or 1 packet in seedbed for 25 plants to set 40 feet of row.

**Culture.** In Florida, seed may be sown in seedbeds from September to January. Plants should be set in rows 3 feet apart, and 1½ to 2 feet apart in the row. When the main head is cut, thick stems form at the base of leaves on the stalk below. At the end of each branch a medium-sized green head develops. These small heads should be cut with 4 to 5 inches of stem, and tied in bunches for market.

**EARLY GREEN SPROUTING OR CALABRESE.** (55 days from plants.) This is the old standard early variety of Broccoli which produces tall, erect, bushy plants, with an attractive dark, bluish-green color to the head. Forms a large, compact central head. After the central head is cut the side shoots send out small heads over a long season. An exceptionally high yielder.

Pkt. 10c; ½ oz. 20c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50  
5 to 25 lbs. \$3.40 per lb.

**TEXAS 107.** (50 days from plants.) Early maturing variety. In numerous Florida trials in different sections of the State it consistently ranked top in yield. Produces a medium size main head with a great number of good size side shoots. Individual bud formation in head and sprouts, of medium size and very tight and compact, slightly lighter green in color than Early Green Sprouting.

Pkt. 10c; ½ oz. 20c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50  
5 to 25 lbs. \$3.40 per lb.

**WALTHAM 29.** (60 days.) Low growing, heavy, short compact plant, which produces a large, central head. The plant is heavily branched with dark green, tight buds that are of medium size, evenly distributed. Both the plant and head have a most attractive dark blue-green color. It has ability to come through periods of hot weather without the heads opening up. Because of its uniformity, a large portion of the crop matures at one time. Its excellent dark green color, fine head and bud texture and high yield make it a valuable addition for Florida. Of excellent quality and appearance for fresh market and for freezing.

Pkt. 10c; ½ oz. 30c; 1 oz. 50c; ¼ lb. \$1.50; 1 lb. \$5.50  
5 to 25 lbs. \$5.40 per lb.



# CABBAGE

Sow 6 ounces in seedbed to plant an acre or 12 ounces per acre in field.  
For the garden, sow 1 packet in seedbed for 50 plants to set 60 feet of row.



A typical plant of Marion Market (Yellows Resistant) Cabbage

**Culture.** Cabbage is a heavy feeder and should be grown rapidly. For maximum yields on sandy loam soils 1500 to 2000 lbs. per acre of a 4-7-5 fertilizer mixture should be applied in the row a few days prior to setting plants. In addition, two or three side dressings of Nitrate of Potash or Nitrate of Soda should be made at three to four week intervals. You cannot grow a big, heavy crop of cabbage on a starvation ration. Where planted in seedbeds, which is the desirable and common practice, plant seed thin in drills 6 inches apart, cover  $\frac{1}{4}$  to  $\frac{1}{2}$  inch deep, using 6 ounces of seed for an acre of plants, and do not make the soil in seedbeds as rich as soil where plants are to be set. When seed is planted in the field, use 12 ounces of seed per acre. Make rows  $2\frac{1}{2}$  to 3 feet apart and set plants 12 to 18 inches apart in row, depending on variety and size of head you wish to produce. If you desire small heads, crowd the plants; for large heads, place further apart. For a winter crop of cabbage in Florida seed is sown in September and early October. For an early spring crop plant in November or early December. For the control of cabbage insects and diseases see page 53.

To control downy mildew disease in cabbage seedbeds use Wettable Spergon Spray or Spergon Dust (See page 53). For the prevention of downy mildew and other cabbage diseases in the field use Dithane D-14 spray or Dithane dust. (See page 53).

Our cabbage seed is grown in disease-free territory on the Pacific coast where Black Rot has never been found by plant pathologists who inspect our seed growing fields each year.

The number of days in parentheses after each variety indicates the relative number of days to secure marketable heads from setting out plants in the field or garden. From the time seed is sown until the plants are ready for the field or garden thirty to thirty-five days are usually required.

Prices quoted are postpaid.

Write for prices on larger quantities.

**JERSEY WAKEFIELD.** (62 days.) Our stock has been bred and selected especially for earliness. Very resistant to cold weather. The small, medium-sized, conical-shaped, pointed heads are very firm, of finest quality, and especially desirable for home use.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**CHARLESTON WAKEFIELD.** (70 days.) The heads are larger and broader at the base, not quite so pointed and later in maturing than Jersey Wakefield. Of uniform size, very solid and of very desirable market quality. Very resistant to cold weather and premature seeding.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**MARION MARKET (Yellows Resistant).** (75 days.) This is a late maturing, disease-resistant strain of Copenhagen Market, highly resistant to Yellows. Larger and coarser in growth than Copenhagen Market but makes somewhat heavier yields, with large, firm, solid heavy heads. Will produce a normal crop and heavy yield on Yellows infested soil where non-resistant varieties will fail. We have an improved, true to type strain of this popular variety.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**MIDSEASON MARKET.** (75 days.) Our strain is uniform as to type and time of maturity, cutting over 50% perfect heads at the first harvest when seeded in the field. The heads are of medium size, hard, globe-shaped on short stems. Leaves are dark green, thick and leathery, making this variety hardy and resistant to cold damage, and also making it a good shipper in bags, the heads not being bruised so readily as other varieties. Of crisp texture and good quality.


Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**RESISTANT COPENHAGEN MARKET (Yellows Resistant)** (New) (70 days.) With the increased interest in Cabbage Yellows Resistance in Florida, which Cabbage Yellows disease has now been found in certain north Florida soils, the only way to combat it being by planting Yellows Resistant strains or varieties, we are offering a resistant strain of Copenhagen Market which can be successfully grown in Yellows infested soils where non-resistant varieties or strains would be a failure. Our strain of Yellows Resistant Copenhagen Market is slightly later than our regular Copenhagen Market (Early Strain No. 86). In other general characters it is similar, producing uniform, medium size, round, hard firm heads, bright to dark green in color, maturing practically all at one time. A most desirable strain of Copenhagen Market to grow as insurance against the dreaded "Cabbage Yellows Disease."

Pkt. 10c; oz. 45c;  $\frac{1}{4}$  lb. \$1.40; 1 lb. \$5.00  
5 to 25 lbs. \$4.90 per lb.


**COPENHAGEN MARKET (Early Strain No. 86).** (68 days.) Our early strain of Copenhagen Market has become very popular in all the large cabbage producing sections of Florida. It is short-stemmed, compact with very solid, round, medium-sized heads, having few outer leaves, thus facilitating close planting, very uniform in shape, and matures practically all at the same time. This uniform maturity is desirable from a marketing standpoint.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.



Copenhagen Market. (Early Strain No. 86.)





Golden Acre



Early Glory



Savoy

**GOLDEN ACRE.** (64 days.) An extremely early, small to medium-sized round headed cabbage of the Copenhagen Market type, but considerably earlier than most stocks of Copenhagen Market. Our strain of this variety heads up very uniformly and is exceptionally well bred.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**EARLY GLORY.** (72 days.) A medium-early, round headed type. Somewhat similar to Copenhagen Market, but slightly later, larger, with a darker green color, and a heavier yield. The heads are protected by large, thick, overlapping wrapper leaves, thus resisting cold damage. Heads are solid, of medium size, and of excellent shipping and market quality. An improved early maturing strain of the old Glory of Enkhuizen.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**EARLY ROUND DUTCH.** (71 days.) A hardy, and dependable variety of round headed cabbage. Heads are firm and of high quality. Plants vigorous, small, and compact, with erect leaves and short stems. Suitable for close planting, but must be fertilized more heavily if planted close in order to produce size. Very slow to bolt or produce "seeders" and withstands cold weather. Especially well suited for muck land production.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**BADGER MARKET (Yellows Resistant) (new).** (72 days.) This new yellows resistant cabbage developed by the U.S.D.A. and the Wisconsin Agricultural Experiment Station, is showing much promise on Florida muck soils. On sandy soils it is inclined to be too small unless heavily fertilized. In season it is only a few days later than Early Copenhagen Market No. 86. It is extremely uniform as to type and time of maturity. It makes a uniform, small, compact, round head, of attractive dark blue-green color, very firm and hard, even on loose muck, with a short to medium core. It is very resistant to splitting, and can be left in the field for ten days to two weeks after it is mature and still show no signs of splitting. It possesses excellent edible quality. This new variety is strongly recommended for Florida muck and the better sandy loam soils where it makes heads two and one-half to three pounds in weight, just the right size for northern markets. Seed supply extremely limited.

Pkt. 10c; oz. 75c;  $\frac{1}{4}$  lb. \$2.25; 1 lb. \$8.00  
5 to 25 lbs. \$7.90 per lb.

**SAVOY-CHIEFTAIN.** (90 days.) Savoy cabbage is commonly called "curly leaf" or "curly" cabbage. It has a distinct flavor which is favored by many people. The plant is of large growth. The large, rich, dark green leaves are uniformly wrinkled or savoyed and curled. The heads are medium in size, hard, and stand a long time without bursting. The best variety of the savoy type for Florida.

Pkt. 10c; oz. 30c;  $\frac{1}{4}$  lb. 85c; 1 lb. \$3.00  
5 to 25 lbs. \$2.90 per lb.

**RED ACRE.** (72 days.) This variety fills a long felt need in being the only red cabbage that matures very early for a red, only a few days later than Early Copenhagen Market. The plant is small, compact and makes good-sized round, hard, deep red colored heads. Very uniform as to type. Especially desirable for muck land production in Florida, but also does exceedingly well on fertile sandy loams.

Pkt. 10c; oz. 35c;  $\frac{1}{4}$  lb. \$1.00; 1 lb. \$3.50  
5 to 25 lbs. \$3.40 per lb.

**ROUND RED DUTCH.** (80 days.) A medium early variety, similar to Red Acre. Is a little more vigorous but with larger head at maturity. It is possible to cut Round Red Dutch just as early as Red Acre and get as much or more yield per acre. The heads are round, solid, with a deep red color.

Pkt. 10c; oz. 35c;  $\frac{1}{4}$  lb. \$1.00; 1 lb. \$3.50  
5 to 25 lbs. \$3.40 per lb.

## CHINESE CABBAGE


Also known as Celery Cabbage, although it is not a true cabbage, and grows more like a Cos or Romaine lettuce. Plant 1 pound per acre.

In the garden make two plantings at different times. 1 packet to 40 feet of row at each planting.

**Culture.** This is a cool-weather plant and should not be grown for a spring or summer crop, because it bolts to seed in hot weather. Seeds should be planted in  $2\frac{1}{2}$  to 3 foot rows where the crop is to mature, thinning out as the plants develop to stand 8 to 12 inches in the row. Chinese Cabbage makes excellent cole slaw.

**MICHILHI.** (70 days.) From seed planting in field or garden to edible stage. Sure headed, maturing uniformly into long, slender, very firm, well blanching, crisp, tender, and sweet heads. It is slightly more vigorous and darker green than other strains, the heads are more uniform, and slower to bolt to seed. The heads become 19 to 20 inches tall,  $3\frac{1}{2}$  to 4 inches thick, tapered near the tip, solid, sweet and tender.

Pkt. 10c; oz. 35c;  $\frac{1}{4}$  lb. \$1.00; 1 lb. \$3.50  
5 to 25 lbs. \$3.40 per lb.



Michihli (Celery Cabbage)



# CANTALOUPE

Plant 1 to 2 pounds per acre.

In the garden plant 1 packet to 50 feet of row.



Smith's Perfect (Florida Golden Dew) Cantaloupes

**Culture.** General cultural requirements, diseases and insects and their control, similar to those of cucumber. (See page 16.) Plant in hills 4 to 5 feet apart, 5 to 6 seeds to a hill. Thin out to two plants per hill when 4 inches high. Plant in 5 foot rows, thinning to 2 feet in the row. Apply 600 lbs. fertilizer 10 days before planting, side dress with 400 to 600 lbs. at vining. For earlier melon crop ahead of the melon worms, use **Hotkaps**. (See page 75). Use Lindane Dust or Spray to control melon and pickle worms. (See page 54).

Prices quoted are postpaid.

Write for special prices on larger quantities.

**SMITH'S PERFECT (Florida Golden Dew or Rock Melon).** (90 days.) A good, strong, vigorous vine growth, with thick, hard, tough upturned leaves, withstanding wind and cold damage. Very resistant to downy mildew disease commonly called blight or rust. Also resists aphids. A very good commercial type, making medium size fruits (average 4 lbs.), very heavy for their size, round in shape, slightly flattened at stem and blossom ends, covered with a fine netting, free of ribbing, a tough rind, makes a good shipper and market melon. The fruits have a greenish rind, which upon ripening turns to a beautiful greenish-yellow. As the fruits ripen they produce a delicious melon odor. For market, it is best to harvest them a little on the green side just as they begin to turn yellow. The flesh is firm, very thick, with a small seed cavity, of attractive deep orange or salmon color, and has a most delicious flavor.

Pkt. 10c; oz. 30c; ¼ lb. 85c; 1 lb. \$3.00;  
5 to 25 lbs. \$2.90 per lb.

**HALE'S BEST, JUMBO STRAIN.** (78 days.) A large fruited strain of Hale's Best, grown very extensively in Georgia and other southern states. Fruits are oval in shape, showing some trace of ribbing, with heavy coarse netting. Average about 4 to 5 lbs. Flesh thick, relatively small seed cavity, deep orange color.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00;  
5 to 25 lbs. \$1.90 per lb.

**HALE'S BEST, No. 36.** (75 days.) An improved strain of Hale's Best, developed for uniformity in shape and complete netting. Fruits are round in shape, with very heavy, coarse netting, and average about three pounds. Flesh thick, small seed cavity, deep salmon color, very sweet, fine quality. Excellent shipper.

Pkt. 10c; oz. 20c; ¼ lb. 60c; 1 lb. \$2.00;  
5 to 25 lbs. \$1.90 per lb.

**GEORGIA 47 (Mildew Resistant) (New).** (72 days.) This new downy mildew and powdery mildew resistant variety is very early, exceedingly prolific, with firm,

thick flesh of golden orange color, very sweet. Fruits are generally too small for a shipping cantaloupe, and have a tendency to produce an unattractive rough swelling at blossom end. For home garden or local markets this variety is highly recommended.

Pkt. 10c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50;  
5 to 25 lbs. \$3.40 per lb.

**NO. 45, POWDERY MILDEW RESISTANT (Hale's Best type).** (80 days.) Attracting much favorable attention in Florida because of its high yielding, good shipping and holding quality. Vines moderately large and produce a very heavy crop. Highly resistant to powdery mildew, but not to downy mildew. Medium sized fruits (about 3½ lbs.) are short-oval, indistinct ribbing and well netted. Flesh thick, firm, sweet and of good quality, light orange color. Can be picked almost mature for local markets or distant shipping.

Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25  
5 to 25 lbs. \$2.15 per lb.

# CAULIFLOWER

Sow 4 ounces in seedbed to plant an acre.

In the garden sow one packet in protected seedbed for thirty plants.

**Culture.** Starting plants in well prepared seedbeds where conditions can be provided for continuous rapid growth. Plant in seedbeds from Sept. to Dec. and cultivate the same as cabbage, spacing 3 to 4 feet between rows and 18 to 24 inches in row. Cauliflower needs rich soil and plenty of moisture. When head develops to the size of an apple, draw outer leaves over head and tie with soft cord to produce attractive, pure white, solid heads. Our cauliflower seed is grown in black rot free territory. The number of days after each variety indicates time to produce edible heads from setting out plants.

Prices quoted are postpaid.


Write for special prices on larger quantities.

**SUPER SNOWBALL.** (55 days.) An early maturing strain. Snowball type, in which curled formation of leaves over head protects it from sun and from freezing. Dwarf, compact plants are strong, vigorous, with many short, large, round, broad, heavy, spreading light green leaves, and short stout stem. Heads are large, smooth, solid, pure white and very uniform; proper growing conditions produce over 90% grade-one heads. Recommended as an early winter crop, to mature in cold weather. Susceptible to "whiptail" and should be grown in alkaline soils with high pH to avoid trouble.

Pkt. 25c; ½ oz. \$1.00; oz. \$1.75; ¼ lb. \$5.25;  
1 lb. \$20.00

**SNOWDRIFT.** (62 days.) One of the surest headers known, especially well adapted to withstand the hot, humid climatic conditions prevalent in Florida. Vigorous plants grow larger and taller than Snowball, are more compact and upright in growth with long, rather narrow, erect growing, dark green leaves which protect head or curd in its early growth. Large, snow-white attractive heads are very deep. Heavy and firm, even under relatively hot, humid growing conditions. An ideal variety for early spring production in Florida. Very free of leafy and ricey heads. Very resistant to "whiptail" disease.

Pkt. 25c; ½ oz. \$1.00; oz. \$1.75; ¼ lb. \$5.25;  
1 lb. \$20.00



Snowdrift Cauliflower



# CARROT

Plant 4 pounds of seed per acre. In the garden make three plantings at different times.  
Plant 2 packets to 100 feet of row at each planting for a family of 3 or 4 people.

**Culture.** Sow in good, deeply worked, fine soil, enriched with 800 to 1200 pounds per acre of a good vegetable fertilizer, in drills  $1\frac{1}{2}$  to 2 feet apart, and cover seed  $\frac{1}{4}$  to  $\frac{1}{2}$  inch deep. When 2 inches high, thin out to 2 to 3 inches apart. Carrots can be planted in Florida from August to March.

Prices quoted are postpaid.  
Write for special prices on larger quantities.




Comparison of root characteristics of carrot varieties. From left to right: Imperator, Nantes (Coreless), Danvers Half Long, Red Core Chantenay.

**DANVERS HALF LONG.** (75 days.) Our stock of this variety is half stump or semi-pointed, from 6 to 7 inches in length from the crown to the tip. The shoulder at the crown is from  $1\frac{1}{2}$  to  $1\frac{3}{4}$  inches thick, and tapers uniformly to a thickness of about  $\frac{3}{4}$ -inch just where the root begins to taper to the tip. The tops are a little larger than those of Chantenay, and the crown is flat without a depression. The roots are smooth and attractive in appearance.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

**LONG IMPERATOR.** (77 days.) The roots average about 10 inches in length, with a shoulder diameter of  $1\frac{1}{4}$  to  $1\frac{3}{4}$  inches, smooth, uniformly tapered to a semi-blunt end, deep rich orange color, both exterior and interior with comparatively inconspicuous core. The flesh is of rich orange color, extending to the center of the root, with an indistinct core; fine grained, tender and of excellent quality. The tops are of medium size, but strong enough for good bunching. For shipping quality, attractiveness and popularity in the market, Kilgore's Bred-Rite strain of Long Imperator is superior to other long-rooted varieties.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.



Long Imperator Carrot

Developed to meet the need for a better market type.

**RED CORE CHANTENAY.** (72 days.) The roots are tapered, less stump rooted, and slightly longer than the old Chantenay, averaging about 6 inches. The interior is solid orange-colored without the usual lemon color in the core. The flesh is very tender and sweet, which with the excellent outside color and shape combines to make this variety valuable for the market.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

**NANTES.** (70 days.) Cylindrical, roots about  $5\frac{1}{2}$  to  $6\frac{1}{2}$  inches long and 1 to  $1\frac{1}{2}$  inches thick, very stump rooted with a small rat tail. The color is clear orange-red and the core is rather undefined, being of the same color as the flesh so that it is called "Coreless." The tops are small and short, but heavier than ordinary stocks of Nantes, and the root is very fine-grained, tender, delicate, sweet and of highest edible quality. An excellent high quality variety for the home garden and local market.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

**LONG ORANGE.** (86 days.) Our stock is the true thick-shouldered strain of Long Orange,  $1\frac{3}{4}$  to 2 inches thick at the shoulders. 10 to 12 inches long, and tapering in a straight line to the tip. The tops are rather large. Grown in sections of Florida where a smooth, reddish-colored, long-rooted carrot is desired.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

**YELLOW BELGIAN (Stock Carrot).** (95 days.) The giant among carrots. Grown exclusively for stock, very large, 12 inches long from crown to tip, and  $2\frac{1}{2}$  inches thick at shoulder. Pale yellow skin and flesh. Hardy, heavy yielder and very nutritious for stock.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.



# CELERY

Sow ½ pound in seedbed to plant an acre.

For the garden sow 1 packet in a well protected seedbed for 150 plants to set 50 feet of row  
Prices quoted on celery seed are postpaid. Write for special prices on larger quantities.



*Typical plants of Green Florida Pascal Celery ready for packing in Howard crates.*

**Culture of Celery.** First sowings in Florida are made the latter part of June and continue to December. The main sowings for a winter crop are usually made in July or early August, and for the spring crop, grown principally on muck, they are usually made in October and early November. Celery is always started in protected seedbeds. The soil must be put in perfectly fine condition and made into beds 4 to 6 or more inches high, to prevent overflow in case of heavy rains. Sow the seed broadcast or in rows 6 inches apart, using 1 pound of seed to 125-150 yards of bed 3 feet wide. The seed should be barely covered with soil.

We recommend use of burlap or white muslin as a cover over the seedbed stretching it on a tent-shaped or slanting frame built with the sides about 12 inches from the ground to provide ample circulation of air. This cover will serve as shade and break the force of heavy rains. After the seed is sown, it is desirable to sprinkle with a very fine spray in order to keep the soil cool and moist, which tends to increase the germination of celery seed during the hot summer months. With surface irrigation, water should be circulated continuously around the beds.

When plants in seedbed are 6 inches high, transplant to the field in double rows 3 to 3½ feet apart, with 7 inches between the double rows, and set the plants 4 to 5 inches apart in the row. The plants may be set in single rows 3 feet apart, allowing 4 to 5 inches between plants in the row. The Pascal varieties should always be planted at least 6 inches in the row. To grow fine celery, a large supply of moisture and fertilizer is essential from the time the seed is sown in the seedbeds until the crop is ready for harvest.

The number of days in parentheses after each variety indicates the relative number of days to marketable stage or harvest from setting out plants in the field or garden. It usually requires about seven or eight weeks to grow plants large enough to be set in the field.

In May and June we test samples of all our celery seed in our laboratory germinator, and also in outdoor seedbeds at Sanford and Plant City, and the germination is printed on every package of celery seed so our customers will know how much to plant in order to secure a satisfactory stand in the seedbed.

For the control of insects and diseases of celery see page 53. For the control of pink rot of celery use Granular Aero Cyanamid.

Good control of black heart in celery has been obtained by spraying weekly with 10 lbs. Calcium Nitrate dissolved in 100 gal. water per acre.

**GREEN FLORIDA PASCAL.** (125 days.) A very vigorous, compact-growing variety, with large, broad, rounded, heavy, dark green leaves, and with broad, semi-round, medium-long, exceedingly thick, very meaty, solid, smooth, glossy, attractive appearing ribs. The ribs or petioles will average from 7½ to 8 inches to the first joint and plants will range from 20 to 23 inches overall. It develops a much better heart than other varieties of the Green Pascal type. The ribs or stalks are of ideal edible quality, being tender, brittle, stringless and crisp. Even the outermost stalks possess an unsurpassed flavor and are surprisingly rich and nutty. Our Florida Pascal strain has been developed for smooth glossy ribs, a taller, more compact growth, and better heart formation than other strains of Green Pascal.

This stock is not subject to premature seeding, and can be grown at any time during the fall, winter or spring in Florida. It is important not to cut this variety too soon, as it requires 2 to 3 weeks more time in the field than varieties of the Golden type, but will out-yield the Golden type. The yield and quality are materially cut down if harvested too soon. Set plants of this variety at least 6 to 7 inches apart in the row for large sizes and heavy yields. The seeds of Green Pascal are larger than Golden type celery, and therefore 25 to 50% more seed by weight is required to get the same stand.

Pkt. 15c; ½ oz. \$1.00; 1 oz. \$1.50  
¼ lb. \$400; 1 lb. \$15.00

**SUMMER PASCAL** (Waltham Strain). (115 days.) This is the earliest maturing of all the Green Pascal varieties. Developed by the Waltham, Mass., Agricultural Experiment Station for a rapidly growing Green Pascal type of celery with long, upright growing, compact ribs or petioles, making a very attractive appearing celery for the markets. The ribs are thick and semi-rounded, but not quite as smooth as our strain of Green Florida Pascal, although the Waltham strain will mature about a week earlier, and makes a high yield of large sizes. This strain must be cut promptly when ready or it will tend to go down, and won't hold up in the field after maturity as well as Green Florida Pascal.

Pkt. 15c; ½ oz. 85c; oz. \$1.35; ¼ lb. \$3.40;  
1 lb. \$12.50



## CELERY (Cont'd)

**FLORIDA GOLDEN NO. 15.** (115 days.) One of the best varieties for a mid-winter crop in Florida, because it withstands cold weather better than most varieties of the Special or Golden Plume type. Selected for uniformly long, thick, round ribs, large full hearts of rich golden yellow color, and for heavy yields of good sizes. The ribs average eight to ten inches in the first joint, are not thin and flat, but are rounded and exceptionally thick. The plants range from 21 to 24 inches over all, making a most attractive pack. We introduced this improved variety. It is intermediate between the Old Golden and Special or Golden Plume types, possessing the desirable characteristics of both types. Recommended especially for a Florida mid-winter crop.

Pkt. 15c; ½ oz. \$1.00; 1 oz. \$1.50; ¼ lb. \$4.00;  
1 lb. \$15.00



Part of typical row of Florida Golden No. 15 Celery

## COLLARD

Sow ½ pound in seedbed to plant an acre, or plant 2 pounds per acre in field.

For the garden make two sowings at different times. Sow one packet in seedbed for 35 plants for each planting to set 50 feet of row, or one packet in 50 feet of row.

**Culture.** Collard is an old standby for winter greens all over the South. May be started in seedbeds or direct in the field, in rows 2½ to 3 feet apart, with plants 14 to 18 inches apart in the row. Use about 1200 pounds of fertilizer per acre before setting plants. Collard can be grown every month of the year in Florida, but it grows best and produces the highest edible quality during cool weather. This vegetable should be included in every Florida family garden, as it is very hardy, easy to grow and will furnish an abundance of greens, high in health-producing vitamins.

The number of days in parentheses after each variety indicates the time from setting out plants to harvest. It requires about a month in the seedbed to get plants from seed sowing to transplanting stage.

Prices quoted are postpaid.

Write for special prices on larger quantities.


**GEORGIA OR SOUTHERN.** (50 days.) A non-heading type of the cabbage family, very extensively grown for greens in home and commercial gardens in Florida.



Georgia or Southern

Plant grows from two to three feet tall, is erect and spreading, with many large, undulated leaves on a white stem. Withstands cold as well as heat, and adverse soil conditions. A high yielding hardy type, of excellent quality when cooked. This is the old standard variety for Florida and the South, and is used more extensively than any other. Our strain has been selected for uniformity and resistance to early seeding.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.



Vates

**VATES.** (55 days.) A very large, vigorous, rather low growing plant, not subject to wind damage, with very large, broad, heavy, thick leaf blades, the leaf blade developing from the base of the short stem or petiole. The leaves are smooth with an attractive deep, dark green color, the stems and leaves never turning yellow or purplish even in the coldest weather or on poor soils. It retains this deep green color in shipment and on the markets. A heavy yielder slow to bolt to seed in cold or hot weather. Uniform as to plant type and free of mixture, our stock having been grown direct from the originator's most recent selection. In quality it is claimed to be superior to any of the old standard varieties.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.



# CUCUMBER

In the garden plant 1 packet to 50 feet of row.  
Plant 2 pounds per acre.

**Culture.** Rich, moist, sandy loam soil is the best for growing cucumbers. The soil should be well fertilized with a good vegetable fertilizer applied at the rate of 1200 lbs. per acre a week or ten days before planting; 600 to 1000 lbs. is generally used as a side-dresser just before the plants start to run. Fall plantings are made in late August and in September. Spring planting in Florida begins early in January and continues through February. Plant in rows 5 to 6 feet apart, 6 inches apart in the row, and thin to 1 or 2 feet between every two plants in the row. Sometimes it is difficult to get a stand early if the ground is wet and cold, as the seed will rot before it germinates. Therefore seed should be treated with Spergon.

If young plants are very badly damaged with frost or cold, it is best to plant over. When young plants are stunted they never make a normal crop and usually not much earlier than if they were planted over. A row of oats planted between every two rows of cucumbers makes a good windbreak.

In isolated locations it is always desirable to provide a hive of bees in cucumber fields to ensure perfect pollination and a good set of fruits. The bees should not be placed in the field until the cucumbers begin to blossom, in order to provide food for the bees so they will remain in the cucumber fields. Without proper pollination the cucumbers are deformed, or at least a large percentage of bottlenecks and nubbins are produced, and a small set of fruits develops.

For the control of insects and diseases of cucumbers see page 54. To control angular leaf spot treat seed with a mercuric chloride. (See page 52.) The best control for downy mildew, sometimes called "rust," is Dithane D-14 spray or Dithane dust. (See page 54.) The most effective and safest control for aphids and some other cucumber insects is Isotox spray or Kilgram dust. (See page 54).

For an earlier cucumber crop and a protection against beetles, use **Hotkaps**. (See page 75).

For dill to make cucumber pickles see Page 20.

The number of days after each variety represents the time from seed planting to first picking under average spring or fall conditions in Florida.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**MARKETER (Early Green Market).** (50 days.) The leading variety for yield and quality. Holds its dark green color much better than most varieties, when grown in late spring or early fall. It averages about 8 inches in length, and is especially desired by the eastern and southern markets where a medium size cucumber is preferred. It is particularly admired for the neat appearance and uniformity of its many fruits, which are of an exceptionally rich, uniform, dark green color down to the blossom end. It seems to stand heat of late spring better than other varieties. Exceedingly prolific, throwing an exceptionally large number of female blossoms per plant. The fruits are exceptionally uniform in size, shape and color, with practically no jumbos or oversize even when not harvested for several days. It will average over 90 per cent fancies. Buyers pay a premium for this cucumber, when other varieties go begging at lower prices.

**Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25**  
**5 to 25 lbs. \$2.15 per lb.**

**SANTEE (Downy Mildew Resistant).** (New dark green strain) (45 days.) Developed by Dr. W. C. Barnes of the South Carolina Experiment Station. Similar to Palmetto, but earlier and more prolific, and darker green color. Attractive cylindrical shape with blunt ends. Similar to but darker green than Cubit. Quite resistant to downy mildew.

**Pkt. 10c; oz. 30c; ¼ lb. 85c; 1 lb. \$2.75**  
**5 to 25 lbs. \$2.65 per lb.**



*A field and typical fruits of Marketer (Early Green Market) Cucumbers at Fort Myers, Florida*






*Packing Marketer (Early Green Market) Cucumbers on a farm in the Plant City section*

**CUBIT.** (42 days.) The plants are very prolific; the smooth fruits are dark green color, and average about 9 inches in length, are cylindrical in shape with blunt ends, without any taper, and with a very small seed cavity making them ideally adapted for slicing. They hold their color, crisp texture and fresh quality well in shipping and on the market. This variety makes the most uniformly beautiful shaped cucumber of any variety. Attractive appearing, high quality, early maturing, and good shipping and keeping variety.

**Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25**  
5 to 25 lbs. \$2.15 per lb.

*Kilgore's Bred-Rite Seeds are selected, tested and proven in Florida for Florida.*

*(See page 34.)*



**CUBIT**

*Uniform, large attractive fruits.*

**PUERTO RICO NO. 39 (Mildew Resistant) (Special Long Strain).** (42 days.) This variety was developed by the Puerto Rico Agricultural Experiment station for resistance to downy mildew or blight. In cool, rainy weather, especially in fall, when other varieties go down with mildew disease, this variety continues to grow and produce an abundance of fruits, making it especially valuable as a home garden variety. Although this variety is highly resistant to downy mildew it is not immune, and should be sprayed or dusted regularly with a good fungicide like Dithane. The vines are rather compact in growth, hardy and very productive. As originally introduced the fruits of this variety were rather short, but we have selected for long fruits until with our special long strain the fruits will average about 7½ inches in length with dark green color.

**Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25**  
5 to 25 lbs. \$2.15 per lb.

**STRAIGHT-8.** (47 days.) This plant is vigorous in growth, very prolific and produces symmetrical, almost cylindrical fruits. As indicated by the name, the fruits are straight, average about eight inches long, uniform in size, shape and color, making it an ideal packing and shipping type. Recommended for late fall planting in

Florida, not for a late spring crop because the fruits have a tendency to be too light in color late in the season.

**Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25**  
5 to 25 lbs. \$2.15 per lb.

**WAUCHULA STAYS GREEN (Special Long Strain).** (40 days.) Plants are prolific, producing a heavy crop of fancy fruits, and are very early. Fruits average 8 inches in length and are smooth, cylindrical in shape, medium green in color, and very uniform in size, shape and color. This strain is especially well suited for fall and early spring planting in Florida, but is not desirable for late spring.


**Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25**  
5 to 25 lbs. \$2.15 per lb.

**STAYS GREEN (Pickling Type).** (40 days.) This variety should be grown only for pickles, not for a slicing or as a fresh market cucumber. It is very early and exceedingly productive of short, dark green, warty cucumbers most desirable for dill pickles.

**Pkt. 10c; oz. 25c; ¼ lb. 65c; 1 lb. \$2.25**  
5 to 25 lbs. \$2.15 per lb.

**PALMETTO (Mildew Resistant).** (50 days.) Developed by Dr. W. C. Barnes of the Clemson College Truck Experiment Station, Charleston, S. C., from a cross of Puerto Rico 40 and Cubit in order to develop a good market type possessing the downy mildew resistance of the Puerto Rico 40. This variety produces vigorous vine growth, and is highly resistant to downy mildew, the most serious cucumber disease of the south. Palmetto will survive and produce a good crop, when other varieties may go down with mildew under cool, wet weather conditions.

**Pkt. 10c; oz. 30c; ¼ lb. 85c; 1 lb. \$2.75**  
5 to 25 lbs. \$2.65 per lb.



*Palmetto (Mildew Resistant) Cucumbers*



# SWEET CORN

10 pounds seed required per acre. In the garden make two plantings. Plant  $\frac{1}{4}$  lb. to 200 feet of row at each planting.

Our Hybrid Sweet Corn seed is now graded and sized for better, more accurate stands and more uniform maturity.

**Culture.** Rows should be  $2\frac{1}{2}$  to 3 feet apart with the plants 9 to 15 inches depending on type and season. Sweet corn requires much more fertilizer than field corn. Use about 1000 lbs. per acre of a good commercial fertilizer mixture high in potash. It is claimed that potash helps the tips of the ears to fill better and improves the edible quality of sweet corn. When corn is about knee high apply 200 lbs. per acre of sodium nitrate or its equivalent. This fertilizer practice will greatly increase the yield. Early plantings are made in Florida from January 1st until March. Ship in special corn crates, five to six dozen ears per crate. For control of corn ear worms and leaf blight see page 54.

**GOLDEN CROSS BANTAM.** (73 days.) Golden Cross Bantam is a hybrid sweet corn that has done well as a high quality, yellow-kerneled, early variety in Florida. The small, short stalk ( $5\frac{1}{2}$  to 6 feet) is very sturdy and vigorous. Suckers quite badly. Ears 6 to 7 inches long, 10 to 14 rows. Kernels are attractive, yellow in color. Uniform, higher yielding. Recommended for home use, local markets and for shipping to select markets.

$\frac{1}{4}$  lb. 20c; 1 lb. 75c; 2 lbs. \$1.25, postpaid  
Not prepaid, 10 lbs. \$3.80; 25 lbs. \$9.25; 100 lbs. \$36.00


**CALUMET.** (76 days.) In comparative trial plantings at experiment stations and in commercial plantings all over Florida, this new hybrid variety has received the high ratings for yield, uniformity and attractive appearance, and appears to have some resistance to ear worms. The plant is  $6\frac{1}{2}$  to 7 feet, slightly taller than Ioana, is vigorous, relatively resistant to drouth, and almost suckerless, with large, dark green leaves. The ears are long, averaging about 8 inches, cylindrical, well filled to the tips and with tight husks of good length. Kernels are light yellow, medium wide, very deep, usually in 12 to 14 rows. Very uniform in maturing and a high yielder. Makes a very attractive pack of from 5 to 6 dozen per crate usually 5 to  $5\frac{1}{2}$  dozen. One of the finest of the many hybrids recently introduced.

$\frac{1}{4}$  lb. 20c; 1 lb. 75c; 2 lbs. \$1.25, postpaid  
Not prepaid, 10 lbs. \$4.70; 25 lbs. \$11.50; 100 lbs. \$45.00



Calumet Sweet Corn. Note kernel covering over tip of ears.

(For Field and Roasting Ear Corn, See Pages 38 and 39)



Ioana

**IOANA.** (74 days.) A very productive, yellow-kerneled, hybrid. Vigorous plants are highly resistant to drouth. Medium tall, 6 feet, sturdy, with large, broad and numerous dark green leaves. Ears  $7\frac{1}{2}$  to 8 inches long, 12 to 14 rowed, well-filled with medium wide, attractive light yellow kernels of fine quality and flavor, making it highly resistant to worms. Husks retain green color for a long time, making for fresh, attractive market appearance. Uniform in plant and ear characters, and very productive.

$\frac{1}{4}$  lb. 20c; 1 lb. 75c; 2 lbs. \$1.25, postpaid  
Not prepaid, 10 lbs. \$3.80; 25 lbs. \$9.25; 100 lbs. \$36.00

**GOLD RUSH HYBRID.** (70 days.) This early maturing hybrid makes small, short plants,  $5\frac{1}{2}$  to 6 feet tall, sturdy, vigorous and heavy yielding, mostly two ears per plant, relatively few suckers, fairly resistant to Helminthosporium leaf blight. Medium to large size slightly tapering ears, rather short, averaging  $7\frac{1}{2}$  to 8 inches, fairly thick, with tight, good heavy husk, 12 to 14 rows.

$\frac{1}{4}$  lb. 20c; 1 lb. 75c; 2 lbs. \$1.25, postpaid  
Not prepaid, 10 lbs. \$4.70; 25 lbs. \$11.50; 100 lbs. \$45.00

**ERIE** (75 days.) This hybrid, which won an All-America award several years ago, has been thoroughly tested a number of years in different sections of Florida, and without exception it has always ranked among the top in yield, appearance, quality and general rating. Is slightly later than Golden Cross Bantam and Ioana. Vigorous and relatively resistant to drouth. It is widely adapted and always high yielding. Makes very attractive appearing, long cylindrical ears, mainly 12-14 rowed. Especially well suited to Florida growing conditions. It should be given a thorough trial by every Florida sweet corn grower.

$\frac{1}{4}$  lb. 20c; 1 lb. 80c; 2 lbs. \$1.40, postpaid  
Not prepaid 10 lbs. \$3.90; 25 lbs. \$12.00; 100 lbs. \$47.00

**HURON.** (74 days.) This hybrid in numerous trial plantings in Florida shows much promise because of its vigorous growth. Heavy, dark green foliage, resistant to heat and drouth and comparatively resistant to ear worms. Especially desirable for late spring crop in Florida. Seems to be adapted to sandy loam soils. Medium height, 7 feet, and very sturdy, with few suckers. Ears are long, averaging 8 to  $8\frac{1}{2}$  inches, slim with a small cob, with 14 rows of deep, narrow, light yellow kernels. Makes high yields even under poor growing conditions.

$\frac{1}{4}$  lb. 20c; 1 lb. 75c; 2 lbs. \$1.25, postpaid  
Not prepaid, 10 lbs. \$4.70; 25 lbs. \$11.50; 100 lbs. \$45.00

**ILLINOIS GOLDEN No. 10.** (74 days.) A heavy yielding main crop corn. Seems well adapted to muckland production in Florida. Plants are vigorous,  $6\frac{1}{2}$  feet tall, with some suckers, drouth tolerant. Somewhat resistant to ear worms. Medium long, large, thick, heavy ears, covered with large, fairly tight husks, with 12 to 14 straight rows of large medium-yellow kernels.

$\frac{1}{4}$  lb. 20c; 1 lb. 75c; 2 lbs. \$1.25, postpaid  
Not prepaid, 10 lbs. \$4.50; 25 lbs. \$11.00; 100 lbs. \$43.00



# EGGPLANT

Sow  $\frac{1}{4}$  pound in seedbed to plant an acre.

For the garden sow one packet in seedbed for 17 plants to set 50 feet of row.

**Culture.** For a spring crop, plant seed in November, December, and January; for fall crop, plant in June, July, and August, in muslin or burlap-covered seedbeds. When set in the open, the plants may need some protection from the sun. Some growers use palmetto fans, stuck slanting on the south side of the hill, which will keep the sun from striking the newly set plants with full force. Have 4 to 4½ feet apart and 2½ to 3 feet between the plants. Eggplant being a long season crop requires several fertilizer applications. Best results are obtained by applying half or more of a complete fertilizer at planting time and the remainder when the crop is one-third to one-half grown. The initial application should be made in two bands each located 2 or 3 inches below and 3 inches to the side of the plant row. Subsequent application should be drilled close to the plant row.

Top-dressing applications of nitrogen or a combination of nitrogen and potash vary in amount and frequency according to seasonal conditions. Two or three applications at rates equivalent to 100 pounds nitrate of soda and 25 pounds muriate of potash per acre generally meet the needs during a given growing period. For the control of insects and diseases of eggplant see page 54.

The number of days in parentheses after each variety indicates the time to first marketable fruits, from setting out plants in the field or garden. It usually requires six to eight weeks to produce plants for field setting.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*

If interested particularly in Eggplant production write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida for a free copy of circular 109 entitled "Eggplant Production Guide." Similarly, if particularly interested in the production of any other vegetable crop, write the Experiment Station for a free copy of their production guide on that crop, being sure to indicate the crop on which you desire guidance.

**FORT MYERS MARKET.** (83 days.) We introduced this variety, which is today one of the leading varieties grown in the South. Our breeding department has made special selections to improve the variety for uniformity in type of plant, size, shape and color of fruit. In numerous tests we have found our strain superior to others in that it produces immense yields of uniform, large-sized fancy fruits over a long period. The plants are of the high-bush type, with very tall, vigorous growth. The fruits are slightly elongated, cylindrical in shape, and they possess the intense deep black color so much in demand among buyers and in the markets.

**Pkt. 10c; oz. 65c;  $\frac{1}{4}$  lb. \$2.15; 1 lb. \$7.50**  
**5 to 25 lbs. \$7.40 per lb.**

**FLORIDA MARKET (Cook's Strain) (phomopsis resistant).** (75 days.) This is the disease resistant variety of eggplant which Florida growers have been waiting for in order to make eggplant production on diseased soils a profitable venture. Tip over, leaf blight and fruit rot disease all caused by *Phomopsis* fungus has become so severe in the state as to make eggplant production unprofitable in some sections. This new variety was developed by Dr. Phares Decker, Plant Pathologist at the Florida Agricultural Experiment Station, Gainesville, Fla. It makes a very vigorous growing, sturdy bush, holding the fruits well off the ground. It is extremely early and a heavy yielder, bearing over a long season. This makes it a very desirable variety for south Florida where it can be planted in early fall for production throughout the winter and spring months. The attractive fruits are cylindrical in shape and develop to a large size. They are of an attractive, dark color. After several years of selection on the original Florida Market variety, we have in this **Cook's strain** a great improvement in color, shape and uniformity. Under good growing conditions it makes uniform dark colored fruits.


**Pkt. 10c; oz. 75c;  $\frac{1}{4}$  lb. \$2.50; 1 lb. \$9.00**  
**5 to 25 lbs. \$8.90 per lb.**



*A field of Fort Myers Market Eggplant at harvest time.*



# ENDIVE



*Improved Florida Deep Heart Escarolle*

Sow  $\frac{1}{2}$  to 1 pound in seedbed to plant an acre, or 1 to  $1\frac{1}{2}$  pounds per acre in the field.

For the garden sow one packet in seedbed for 50 plants to set 40 feet of row, or plant one packet to 40 feet of row.

**Culture.** Plant from August to February in seedbeds or in the fields or garden the same as lettuce. Set in rows  $1\frac{1}{2}$  to 2 feet apart and 9 inches apart in the row. Soil and fertilizer requirements are the same as for lettuce. (See page 21).

The number of days indicated after each variety is the number of days from seed planting in the field or garden to edible or harvest stage. For insect control see page 54.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**ESCAROLLE—IMPROVED FLORIDA DEEP HEART.** (90 days.) One of the best and most prolific types to grow commercially. In great demand on northern markets, and well adapted to Florida soil and climatic conditions. The leaves are large, broad, thick, slightly crumpled, of medium green color, with thick, nearly white midribs. Our improved stock of this variety is of the Full-Heart type, developing an upright growth, producing a very heavy, thick matted plant with large, deep, well balanced creamy white hearts.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

**RUFFEC GREEN CURLED.** (Green Ribbed). (95 days.) Sometimes called "Chicory." A hardy vigorous sort, with deep green, curled, finely cut leaves, giving a mossy appearance. The midrib is thick, broad, green, tender and fleshy. A dense mass of deeply divided leaves forms the head which blanches readily to a beautiful creamy white.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

# DILL

A packet will sow 10 ft. of row; an ounce 50 ft. of row.

**DILL—LONG ISLAND MAMMOTH.** (65 days.) Partially matured seed clusters and flowerheads are useful for flavoring dill pickles. A tall growing plant, 2 to  $2\frac{1}{2}$  feet high, vigorous, with very finely divided leaves. Seed clusters are borne in umbels or clusters, have a strong aromatic odor, and mildly pungent flavor.

Pkt. 10c; oz. 35c;  $\frac{1}{4}$  lb. \$1.00; 1 lb. \$3.50

# GARLIC

(90 days)

Plant 1 pound to 30 feet of row.

Garlic sets come in the form of bulbs which are divided into so-called sets or cloves, which are planted the same as onion sets, from August to November, in rows 15 to 20 inches apart, setting the cloves 3 to 4 inches apart in the row, being sure not to cover the tips of the cloves. Garlic is a southern crop and does well in Florida. After thorough curing, the bulbs should be stored in a dry, airy place until used.

$\frac{1}{2}$  lb. 45c; 1 lb. 85c; 5 to 25 lbs. 75c per lb.; postpaid

# KALE

Plant 1 ounce to 100 feet of row, or 1 packet to 25 feet of row.

**Culture.** Kale is grown for greens. Plant in Florida from August to April, in rows  $2\frac{1}{2}$  feet apart, and for large plants thin from 20 to 40 inches apart in the row. The plants are extremely hardy, withstanding the coldest weather in any part of Florida. The plants also stand heat, and can be grown almost the year around in Florida, except in midsummer.

**DWARF BLUE CURLED.** (70 days from seed sowing to edible stage.) This is the most desirable hardy variety for Florida and for the markets. It forms a handsome, dwarf, spreading plant with many long, heavy, finely curled, attractive, blue-green, plumelike leaves. Our stock is exceptionally well-bred, uniform and very attractive. This vegetable ranks high in vitamin content.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

# KOHL RABI

For the garden make short row plantings at different times.

Plant 1 packet to 25 feet of row.

**Culture.** Grow in row  $1\frac{1}{2}$  to 2 feet apart, thinning from 6 to 8 inches apart in the rows. Planting every ten days will give a succession of tender, fleshy bulbs until hot weather. Start fall planting September, and successive plantings can be made until early spring.

**WHITE VIENNA.** (50 days from seed sowing to edible stage.) The fleshy stems, or so-called bulbs, are formed above ground and are very light green, tender and delicate, and of best quality if used when about 2 to 3 inches in diameter, before fully grown. Prepare by boiling the same as turnips. They make a delicious and nutritious dish.

Pkt. 10c;  $\frac{1}{2}$  oz. 25c; oz. 40c;  $\frac{1}{4}$  lb. \$1.15; 1 lb. \$4.00  
5 to 25 lbs. \$3.90 per lb.

# LEEK

Sow  $1\frac{1}{2}$  lbs. in seedbed to plant an acre, or plant 3 lbs. per acre in the field.

For the garden sow 1 pkt. in seedbed for 120 plants to set 50 feet of row, or plant 1 pkt. to 50 feet of row.

**Culture.** Leek seed may be planted any time during the fall or winter directly in the field, or preferably in seedbeds, in good rich soil. When plants in seedbeds are six to eight inches high, transplant to rows 18 inches apart, and 5 to 6 inches apart in the row, or thin out to this spacing in the field. The long, thick stems when boiled and served with a butter or cream dressing are delicious. They are also used for seasoning and flavoring soups and stews, and in salads, and may be eaten like green onions. The stems are eight to ten inches long and  $1\frac{1}{2}$  inches thick. When marketed they are tied in bunches with twelve Leeks to a bunch.

**LARGE AMERICAN FLAG.** (140 days from seed planting in field or garden to full edible stage.) The heaviest yielding and the best variety for Florida growing conditions. It makes larger and thicker, more fleshy stems than other varieties.

Pkt. 10c;  $\frac{1}{2}$  oz. 25c; oz. 45c;  $\frac{1}{4}$  lb. \$1.40; 1 lb. \$5.00  
5 to 25 lbs. \$4.90 per lb.



# LETTUCE

Sow  $\frac{1}{2}$  to 1 pound in seedbed to plant an acre, or plant 2 to 3 pounds per acre in field. For the garden make two plantings at different times. Sow 1 packet in seedbed for 60 to 80 plants to set 50 to 75 ft. of row at each planting.

**Culture.** First sowings are made in Florida the beginning of September and continue until March. Low, damp soil is best suited for this crop. Lettuce must be grown rapidly in order to give it size, tenderness, crispness and high edible quality. Therefore, the land must be highly fertilized and moist. In Florida about one ton per acre of 4-7-5 fertilizer is commonly used for the Boston, Romaine and leaf types, where 800 to 1200 pounds per acre of a 4-8-8 fertilizer is recommended for the Iceberg type. Broadcast the seed on good moist seedbeds that have been thoroughly prepared, or sow in rows six inches apart and use a light roller to press seed into soil. A pound of seed sown broadcast plants a bed three feet by a hundred yards. Place burlap or well-washed old fertilizer sacks over the beds until seeds sprout. Seedbeds should be kept thoroughly moist at all times after the seed is planted. In hot weather germination may be hastened and improved by placing lettuce seed in moist cloth in the ice box for several days before sowing. In level culture, lettuce plants are usually set 15 by 15 inches in the field, or in rows  $1\frac{1}{2}$  to 2 feet apart, and 10 to 15 inches in the row, or in double rows 16 inches apart, with  $3\frac{1}{2}$  to 4 feet from center to center of double rows, and 14 inches in the rows. Plants should be watered as set in field. For insect control see page 54.

Rusting of head lettuce and Romaine is usually due to boron deficiency. To prevent or stop this trouble, it is recommended to spray with 1 lb. borax to 50 gallons of water applied at the rate of not more than a total of 4 pounds borax per acre in two applications. Too much borax is toxic.

The number of days in parentheses after each variety represents the time required to develop marketable heads from the planting of seed in seedbed or in the field or garden under average Florida conditions.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**KILGORE'S MAMMOTH ROMAINE.** (73 days.) Our Mammoth Romaine is taller, more vigorous and makes larger heads with correspondingly heavier yields than other strains of Romaines or Cos lettuce. It can be left in the fields for 10 days to two weeks after it is ready to harvest without danger of bolting to seed. It forms firm, solid heads with the leaves well capped over the top. We have selected and re-selected this strain until it is very uniform in all respects. See second paragraph under "Culture" for control of rusting, to which this strain is somewhat susceptible.

**Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00**  
5 to 25 lbs. \$1.90 per lb.

**PARRIS ISLAND COS (Mosaic Resistant) (new).** (68 days.) This is a new Cos or Romaine type resistant or tolerant to lettuce mosaic, and very resistant to premature seeding. Similar to Dark Green Cos, but a shade lighter green, with the leaves of somewhat heavier texture. Of value wherever mosaic is a limiting factor in the growing of Cos type lettuce. Very uniform and attractive.

**Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50**  
5 to 25 lbs. \$2.40 per lb.

**SIMPSON, BLACK-SEEDED.** (46 days.) The best large, very early, nonheading or leaf variety. The leaves are very broad, much blistered and crumpled, and excessively frilled on the borders, light yellowish-green in color, very tender, of fine delicate texture, and of excellent quality.

**Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00**  
5 to 25 lbs. \$1.90 per lb.

**WHITE BOSTON (Ocoee Strain).** (75 days.) This strain of White Boston was originated and developed in the Winter Garden section of Florida, and has been selected under Florida conditions for the development of a large framed lettuce similar to Big Boston. Produces large heads and heavy yields.

**Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00**  
5 to 25 lbs. \$1.90 per lb.

**BIG BOSTON.** (75 days.) The heads are of cabbage or butterhead type, large sized, broad, slightly spreading, globular, with a broad shoulder at the base. The outer leaves are broad, almost smooth, but wavy at the edge, medium light green with a tinge of brown on outer leaf margins.

**Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00**  
5 to 25 lbs. \$1.90 per lb.

**MIGNONETTE.** (67 days.) An early maturing, exceedingly high quality, small crisp head lettuce, large, tinged with white hearts, not grown for commercial shipment, but useful for the Florida home gardener and for local markets because of its exceedingly high edible quality, being sweet-flavored, crisp and tender.


**Pkt. 10c; oz. 35c;  $\frac{1}{4}$  lb. \$1.00; 1 lb. \$3.50**  
5 to 25 lbs. \$3.40 per lb.

**BIBB.** (60 days.) A very small, early maturing, loose butterhead type of lettuce, with thick, tender leaves of deep glossy green color, rich in vitamins, rich, creamy yellow interior head possessing superior crispness and flavor. Recommended for home gardens and local markets where high edible quality is desired. Set plants only 6 inches apart because of their small size.

**Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00**  
5 to 25 lbs. \$1.90 per lb.

**GREAT LAKES (No. 118).** (83 days.) This strain 118 has been developed for uniformity of size and type, and is the very best strain for Florida. The plants of Great Lakes are large, vigorous and have a distinctive, deep grassy-green, attractive appearance. The heads are relatively large, compact, firm and solid, with abundant wrapper leaves which cover the heads, protecting them from cold and heat injury. Experience indicates that Great Lakes will stand high temperatures, is much less subject to bottom rot, tip burn and slime rot, and does not bolt so readily as other varieties of this type. It offers a possibility of prolonging the Florida lettuce growing season in late spring, and in early fall.

**Pkt. 10c; oz. 40c;  $\frac{1}{4}$  lb. \$1.15; 1 lb. \$4.00**  
5 to 25 lbs. \$3.90 per lb.



*Great Lakes Lettuce (No. 118)*



# M U S T A R D

Plant 4 pounds of seed per acre.  
In the garden make two plantings at different times.  
1 packet to 40 feet of row at each planting.



*Chinese Broad Leaf (Elephant's Ear) Mustard*



*Florida Broad Leaf*

**Culture.** General cultural directions similar to those for cabbage (see page 10), but is always planted direct in the field or garden. Mustard is one of the quickest growing greens. Sow seed August to March in good rich garden soil in rows 24 inches apart and thin from 8 to 10 inches in the row. Heavy fertilization with a fertilizer mixture high in nitrogen or ammonia, and moist soil are conducive for quick growth, high yield and good edible quality. Does best when grown in cool weather of late fall, winter and early spring. Makes a nutritious dish, high in food value, when boiled for greens.

The number of days after each variety indicates the time from seed to harvest.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*



*Fordhook Fancy Mustard*

**FLORIDA BROAD LEAF.** (40 days.) Tall and upright, large, thick, broad, smooth, round or oval, bright medium green leaves (8 to 10 inches long and 8 inches wide). The leaves have a large, distinct, flattened, whitish stem and midrib. It produces more leaf growth than other varieties. Plants stand a long time without bolting to seed, and do well in hot weather. Of excellent edible quality, the flavor being very mild. Fine for greens and salads.

**Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00**  
**5 to 25 lbs. 90c per lb.**

**CHINESE BROAD LEAF (ELEPHANT'S EAR).** (40 days). A hardy and vigorous grower, producing large, broad, elongated or oblong leaves (9 to 10 inches long and 6 inches wide) which are slightly crumpled and of medium dark green color. The leaves have a small, rounded midrib and differ in this respect as well as in shape of leaf from the Florida Broad Leaf. Tender and of excellent quality, with mildly pungent flavor, when boiled for greens.

**Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00**  
**5 to 25 lbs. 90c per lb.**

**FORDHOOK FANCY.** (42 days.) A vigorous, upright grower, producing large, bright green, very deeply fringed leaves, attractive in appearance. A very heavy yielder. The plants stand relatively long without bolting to seed. Of good edible quality, crisp and tender in texture, and mild in flavor. Our stock of this variety is well bred and uniformly true to type.

**Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00**  
**5 to 25 lbs. 90c per lb.**

**TENDERGREEN (MUSTARD SPINACH).** (45 days.) Although not a true mustard, it is like a quick growing mild mustard, with a spinach flavor, and remains tender even in dry, hot weather. An all-season plant, it withstands heat and may be grown throughout the spring and summer. The smooth, oblong leaves are dark green in color, with small white midribs. Relatively slow to bolt to seed.

**Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00**  
**5 to 25 lbs. 90c per lb.**



# OKRA

Plant 8 pounds per acre.

In the garden plant 1 ounce to 75 feet of row for a family of 3 or 4 people.

Okra does wonderfully well all over Florida where the young seed pods are used in different ways and are highly esteemed. In addition to being served alone, pods are used in soups and stews to which they give body. It is a healthful vegetable, and should be planted in every Florida garden. We have done a great deal of work on our seed stocks of this valuable Florida vegetable.

Each year, we make special selections for stock seed from only the best, earliest maturing plants. Our seed growing fields are carefully rogued so as to eliminate any slightly off-type plants. We are confident we have the best strains of okra seed obtainable anywhere.

**Culture.** Okra can be grown in Florida any time from February until September, but it is a warm-weather plant and will not do much until the ground becomes warm. For early planting when the ground is cold always treat okra with Spergon or Arasan. (See pages 61 & 62.) It can be grown in almost any kind of soil except very sandy or poorly drained soils. Plant in rows 3 to 4 feet apart and sow thinly in the drill. Okra should be cut every second day. If this is not done some of the pods become too old for market and the young plants will stop bearing pods. Plants should bear for several months after harvesting starts. Okra does best on sandy loam soil with a fair amount of fertility. On sandy loam soils, 500 to 700 lbs. per acre of fertilizer at time of seeding, and two or three side dressings of nitrate of potash during the bearing period are desirable. Okra is very susceptible to nematodes causing root knot disease, and therefore it is best to plant on land not previously used for growing crops that are subject to attack by nematodes. For the control of root knot nematodes use "D-D" or Soilfume (see page 64). For insect control see page 55.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**CLEMSON SPINELESS.** (55 days.) Plants are semi-dwarf, growing to a height of 4½ feet on fairly good soil and produce uniform, straight, large, thick, ridged pods of the Perkins' Long Green type, of good length, averaging 3 to 3½ inches when ready for market, green in color, and entirely free from spines, thus making it easy to pick without discomfort. Pods are thicker than Perkins' and fill the crates quickly. Plants are close jointed, and produce an enormous yield. Our strain of this variety is highly bred, uniform, and one of the best, the fields showing practically no variation in plant type of pods. Very well adapted for home gardens and local markets, but does not stand long distance shipments as well as Perkins'. In greatest demand on southern markets.

**Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00**  
5 to 25 lbs. 90c per lb.



*Clemson Spineless Okra*



*A field and a hamper of typical pods Perkins' Spineless*

**PERKINS' SPINELESS.** (53 days.) Developed from a cross of Clemson Spineless and Long Green, selected for darker green, longer, more slender pods of spineless type for shipment to northern markets. Pods similar in size, shape and appearance to Perkins' Long Green, pods and plants are practically spineless. Early, productive, makes a semi-dwarf plant 4½ to 5 feet tall. Bred and selected for uniformity in higher yield of dark green, long, slender pods desired in the markets. This Perkins Spineless should not be confused with Clemson Spineless, being an entirely different and distinct variety, the pods having the long, slender shape and dark green appearance of Perkins' Long Green. Perkins' Spineless can be sold in any market for Perkins' Long Green.

**Pkt. 10c; oz. 15c; ¼ lb. 40c; 1 lb. \$1.25**  
5 to 25 lbs. \$1.15 per lb.

**DWARF STALK LONG GREEN POD—CERTIFIED** (Perkins' Long Green type). (50 days.) This improved strain of the Perkins' Long Green type is a big improvement in uniformity and yield over ordinary strains of this variety. Developed by several years of careful painstaking selection. Seed crops, grown specially for us, are inspected several times each session, and certified for uniformity and trueness to type by Georgia Crop Improvement Association. Plants average 3½ to 4 ft. tall, close jointed and branching, and loaded with pods from ground up. Pods are long, usually about 3½ to 4 inches when ready for market, straight, slender, tapered, ridged, deep dark green and meaty. Retain their tenderness and green color for a long time in transit and on the markets. This improved strain earlier, more productive, makes shorter, closer jointed plants than regular old strains of Perkins' Long Green, and has been selected for uniformity in plant and pod type. This Certified seed sold only in 5-lb. and 10-lb. sealed bags with official Georgia Crop Improvement Association certification tag attached to each sealed bag.

**5-lb. sealed bag, \$1.50 per pound;**  
**10-lb. sealed bag, \$1.40 per pound**

For many years, the Kilgore Seed Company has specialized in vegetable seeds for Florida, until today we pride ourselves on the reputation we have gained of handling the very best seeds that can be procured anywhere for Florida growers.


Kilgore's research means better seeds for you because they are bred in Florida for Florida. (See page 34).



# ONION SETS and SEED

For dry bulbs sow 1½ lbs. in seedbed to plant an acre, or plant 3 lbs. per acre in the field. For the garden sow 1 packet in seedbed for 350 to 450 plants to set 80 feet of row or plant 1 packet to 80 feet of row for a family of 3 or 4 people.

## ONION SETS



For green onions or early maturing bulbs.

Plant 8 to 12 bushels of sets per acre. In the garden 1 pint of sets to 40 feet of row.

A bushel weighs about 32 lbs.

Set out in rows 1½ feet apart and 1 to 2 inches apart in rows for green onions. Plant from September to March.

Prices quoted are f.o.b. Plant City except pints and quarts which are postpaid. 1 bushel weighs about 32 lbs.

|                   | Pint | Quart | Peck | Bushel |
|-------------------|------|-------|------|--------|
| White Sets .....  | .40  | .65   | 1.40 | 5.00   |
| Yellow Sets ..... | .40  | .65   | 1.40 | 5.00   |

Available from September to March.

## ONIONS FROM SEED

**Culture.** For large mature bulbs in Florida, first plantings are made the end of August and continued until November. September and October are the best months to plant. The seed is generally planted in the open field, where the onions are to grow, in rows 18 inches apart, and when 4 to 5 inches high they are thinned out from 3 to 4 inches apart, depending on variety and size of bulbs desired. Those taken out can be used to transplant any possible skips. Onion seed can also be planted in seedbeds and transplanted to the field later. This is a safe and sure method, and is becoming more popular in Florida. Transplant when plants are small. The larger the plants when transplanting, the more splits will develop. It is also important to avoid checking plants in growth when transplanting to avoid splits. Do not work soil to onions after they start bulbing. Onions do best in sandy loam or muck soil, and require at least a ton of fertilizer mixture per acre for sandy soils or 1,000 to 1,200 on muck.

The number of days after each variety indicates the time from planting of seed in the field or garden to large mature bulbs under average Florida conditions.

Prices quoted are postpaid.

Write for special prices on larger quantities.

**TEXAS EARLY GRANO (Babosa).** (130 days.) This strain of early Grano or Babosa was developed by the Texas Agricultural Experiment Station for greater uniformity of maturity and earliness. It is fully ten days earlier, more uniform in shape, and slightly smaller in size than regular Grano. Bulbs have light straw-colored skin, with pure white flesh, are globe- or top-shaped, slightly tapering to the root. Mild in flavor with a sweet taste.

Pkt. 10c; ½ oz. 25c; 1 oz. 40c; ¼ lb. \$1.25; 1 lb. \$4.50  
5 to 25 lbs. \$4.40 per lb.

**EXCEL (YELLOW BERMUDA 986).** (100 days.) Produces more onions of high marketable quality ten to fourteen days earlier than regular Yellow Bermuda. It is practically free of bolters, doubles and splits, and color defects, very uniform in maturity, and in size, color and shape of bulbs, the shape being a deep or thick-flat, the color a light yellow. Crisp and mild in flavor, and keeps better in storage than other Yellow Bermudas.

Pkt. 10c; ½ oz. 25c; oz. 40c; ¼ lb. \$1.25; 1 lb. \$4.50  
5 to 25 lbs. \$4.40 per lb.

**GRANEX HYBRID (new).** (120 days.) Developed and released by the U.S.D.A. The first short-day F<sub>1</sub> hybrid onion adapted to the South and most nearly fits the requirements for an onion suitable for growing in Florida. Very early, extremely uniform for type, color and maturity, and does not have the large, thick neck but dries down to a small neck. Practically no splits or bolting to seed. In Texas trials, it has out-yielded both its Grano type and Excel type parents. Yields of 600 to 900—50-lb. bags of U. S. No. 1 onions per acre have been reported. The bulb of Granex is short top-shaped or slightly flat, mild in flavor, yet keeps unusually well. On the basis of earliness, total yield, percentage of marketable bulbs, attractive appearance and general all-around good quality, Granex is strongly recommended.

Pkt. 10c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.50; 1 lb. \$9.00  
5 to 25 lbs. \$8.90 per lb.

**CRYSTAL WAX BERMUDA.** (110 days.) Early maturing, flat in shape, of waxy white color, attractive in appearance, and of mild flavor.

Pkt. 10c; ½ oz. 20c; oz. 35c; ¼ lb. \$1.00; 1 lb. \$3.50  
5 to 25 lbs. \$3.40 per lb.

**WHITE GRANO (White Babosa) (new).** (140 days.) A new variety similar to Early Grano except that the shape is somewhat more spherical and the skin color is a clear attractive white. This new variety is well suited to Florida growing conditions, resistant to bolting or premature seeding and splitting, of excellent edible quality.


Pkt. 10c; ½ oz. 30c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00  
5 to 25 lbs. \$5.90 per lb.

**RED CREOLE.** (150 days.) Very productive, of medium to large size, semi-globe shaped, quite resistant to thrips. Strong in flavor, but one of the very best keepers. Grows slowly and is late in maturing.

Pkt. 10c; ½ oz. 30c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00  
5 to 25 lbs. \$5.90 per lb.


**NEBUKA (Long White Bunching).** (60 days to green onions.) Makes a long stem, much longer than ordinary bunching onions, pure white in color, and without a bulb. Resistant to pinkroot, thrips and smut.

Pkt. 10c; ½ oz. 25c; oz. 45c; ¼ lb. \$1.40; 1 lb. \$5.00  
5 to 25 lbs. \$4.90 per lb.



From left to right—Texas Early Grano, Granex Hybrid, Excel.





*Paramount (Improved Moss Curled)*

## PARSLEY

Plant 1 packet to 150 feet of row, or 2 pounds per acre.  
In the garden plant 1 packet to 30 feet of row.

**Culture.** Parsley can be planted in the fall or spring, in drills 18 inches apart, and thinned to 6 inches in the row. Seed is slow to germinate and it usually takes from two to three weeks to produce a satisfactory stand.

## PEAS

### ENGLISH OR GARDEN

Our pea seed is high germinating, guaranteed western-grown stock, all carefully fitted for the critical grower.

Plant 1 bushel (56 lbs.) per acre in single rows; 1½ bushels (84 lbs.) per acre in double rows.

In the garden make three plantings at different times.

Plant ½ pound to 100 feet of row at each planting.

Although there are hundreds of distinct varieties of English or Garden Peas, we list only two which we have found particularly well adapted to Florida growing conditions, and on which two varieties we have concentrated our efforts in an endeavor to maintain stocks second to none for Florida.

**Culture.** English peas are very hardy and are not commonly attacked by insects. Peas can be grown on almost any kind of soil and with about the same cultivation as beans. Planting in double rows, six to eight inches apart, is a good plan, because this enables the plants to support each other. Single rows should be spaced two and a half to three feet apart, and double rows three to three and a half feet apart from center of double rows. Plant seed 1 to 2 inches apart in the row, and 1½ to 2 inches deep. By working the soil up against the stems of the plants they will bear longer. Use a high grade fertilizer, at the rate of 1,000 to 1,200 pounds per acre. To control soil diseases, and to get better stands, more vigorous growth and increased yields, treat seed with **Sperguson Seed Protectant** at the rate of two ounces per bushel of seed (see pages 52 and 62). Being a legume, the seed should be inoculated with Nitragin. (See page 69.)

The number of days after each variety indicates the relative time from planting to first picking. This time will vary in different sections and in different seasons.

*Write for special prices on larger quantities.*

**HUNDREDFOLD (Kilgore's Winner).** (53 days.) This variety was introduced by us and it has become very popular. It is one of the finest large early peas for Florida sandy loam soils, and in cool seasons is also adapted to Florida muck soils. It is a surer cropper and

**PARAMOUNT (Improved Moss Curled).** (90 days from seed planting to first harvest.) Plant very vigorous in growth, usually 12 inches tall and spreading 20 inches when given sufficient room, with very long, stout stems for bunching. This variety has been bred and selected for uniform, attractive, leaves of very dark green color.

Pkt. 10c; oz. 20c; ¼ lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

## PARSNIP

Plant ½ ounce to 150 feet of row.

In the garden plant 1 packet to 50 feet of row.

**Culture.** Parsnips do best on deep, rich, sandy soil. Sow seed in Florida from September to December, in rows 1½ to 2 feet apart; cover with one-half inch of soil. When plants are well up, thin to 4 inches apart in row. Parsnip seed germinates very slowly. The Parsnip is a winter vegetable with more starch and sugar than other root crops. To prepare, cook roots one hour in water, scrape off the skin. Slice or chop coarsely or mash, and season with milk, pepper and salt, or slices may be fried after seasoning. A delicious and nutritious dish.

**HOLLOW CROWN.** (120 days from seeding to harvest.) Roots 10 to 12 inches long 2½ to 3 inches in diameter at shoulders.

Pkt. 10c; oz. 20c; ¼ lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

a better yielder than other early large podded varieties. The plants are vigorous in growth, eighteen to twenty inches in height, with heavy, very dark green vine and foliage, producing a great abundance of large (4 to 4½ inches long) broad, deep green, pointed pods containing usually eight large dark green peas of fine quality. A very desirable sort for home gardens and for shipping to markets desiring a large podded pea of high quality.

¼ lb. 20c; 1 lb. 55c; 2 lbs. \$1.00, Postpaid

Not prepaid, 14 lbs. (pk.) \$3.50; 56 lbs. (bu.) \$13.00

**LITTLE MARVEL.** (52 days.) The most popular pea for Florida, because of its high quality, earliness and high yield of well filled pods. The most desirable early variety, especially well adapted to Florida muck soils where it is a sure cropper with well filled pods, even in warm seasons. The vines are vigorous, stocky, dark green in color, growing about 1½ feet tall, and are loaded with straight, round, blunt, plump pods about 3 inches long, well filled with 7 to 8 tender peas. A heavy yielder and sure-cropper, valuable for home garden and for early shipping.

¼ lb. 20c; 1 lb. 55c; 2 lbs. \$1.00, Postpaid

Not prepaid, 14 lbs. (pk.) \$3.50; 56 lbs. (bu.) \$13.00



*Little Marvel, a heavy yielding, early maturing, sure-cropper, especially well adapted to Florida.*



# PEPPER

Sow  $\frac{1}{2}$  to 1 pound of seed in seedbed to plant an acre.  
For the garden sow 1 packet in a protected seedbed for  
24 plants to set 40 feet of row.

**Culture.** Peppers, like eggplants, are usually started in cloth-covered seedbeds. For a fall crop in central Florida sow the seed in June, July and August; for a long season winter crop on the east coast plant in August, September and October. For an early spring crop in Florida west coast sections sow seed in November and in north Florida in Decemebr or January.

High temperature is needed to germinate either pepper or eggplant seed satisfactorily, and it is difficult to get a good stand when the ground is wet and cold. Hammock land, or rather moist soil, is best suited to this crop, although any medium good sandy loam soil, properly fertilized, will grow good peppers. Make rows three feet apart and set plants 10 to 16 inches in the row. As this crop is in bearing for a long time, it is most profitable to make several applications of fertilizer, using 400 pounds per acre about every three weeks until at least a ton has been used.

For the control of insects and diseases see page 55.


The number of days in parentheses after each variety indicates the relative time from setting out plants in the field or garden to picking of marketable green peppers. It requires six to eight weeks to produce plants for the field or garden.

**WONDER GIANT.** (74 days.) This variety of the thick meat Florida Giant type is becoming more popular, especially on the west coast of Florida where it has been grown the past few seasons. The vigorous growing plant is more spreading and does not grow tall as Florida Giant. Developed and selected for uniformity in plant and fruit type and productivity. The blocky fruits borne upright on the plant are mostly 4-lobed, with sweet, thick flesh, of deep glossy green color, making an attractive pack.

Pkt. 10c; oz. 60c;  $\frac{1}{4}$  lb. \$1.90; 1 lb. \$7.00  
5 to 25 lbs. \$6.90 per lb.

**YOLO WONDER (Mosaic Resistant Pendent California Wonder Type) (New).** (70 days.) This new variety of thick meat pepper possesses a combination of two desirable qualities not found in any other California Wonder type pepper, namely high resistance to tobacco mosaic disease, and in addition it produces almost 100% pendent fruits. With the fruit hanging down they are protected to a great extent by the large, deep green leaves, and hence do not sunburn so badly as other strains of thick meat pepper which produce upright fruits exposed to the sun. A rather short bush, which makes it well adapted to muck soil. The fruit is rather short and blocky, with a tendency to be indented at the blossom end, resembles California Wonder. Under mosaic conditions it will produce large yields of high quality, dark green, fancy fruits when other varieties produce low yields of malformed and poor color fruits. This promising new disease resistant variety deserves a trial this season.

Pkt. 10c; oz. 70c;  $\frac{1}{4}$  lb. \$2.25; 1 lb. \$8.50  
5 to 25 lbs. \$8.40 per lb.



Yolo Wonder



Harvesting Kilgore's Florida Giant Pepper at Plant City

Prices quoted are postpaid.

Write for special prices on larger quantities.

**FLORIDA GIANT.** (75 days.) This variety, introduced by us, produces an abundant yield of large, heavy, thick walled, fleshy, dark green fruits, over a long period, but requires heavier and more frequent fertilization, and a richer, more moist soil than thin fleshed varieties of the World Beater type. The tall plants are compact but very vigorous and upright in growth. Slightly later than World Beater, they are heavy producers, often bearing six to eight large fruits at one time. The fruits are longer than California Wonder and other thick-meated peppers, about 4 to 4½ inches long, and 3 to 3½ inches in diameter, usually with four lobes, are smooth and well filled at the blossom end. Makes a very attractive pack and carries well to distant markets.

Pkt. 10c; oz. 60c;  $\frac{1}{4}$  lb. \$1.90; 1 lb. \$7.00

5 to 25 lbs. \$6.90 per lb.

**CALIFORNIA WONDER.** (75 days.) An outstanding medium-late variety. Plant is vigorous, upright, prolific. The thick meated, smooth, large, heavy fruits average about four inches in length and about the same in diameter. This is an excellent shipping type, holding up well in transit and on the market, and is in great demand by truckers because of its good holding-up qualities. Solid thick walls and very sweet and mild. Blocky in shape. Color an attractive, glossy dark green. On poor soil, devoid of plant food and moisture, fruits after the first harvest may tend to "button" or be of small size. For best results, and to produce large, fancy fruits over a long season, it requires heavier fertilization and continuous soil moisture.

Pkt. 10c; oz. 60c;  $\frac{1}{4}$  lb. \$1.90; 1 lb. \$7.00


5 to 25 lbs. \$6.90 per lb.



**EARLY CALWONDER.** (65 days.) An early maturing strain of California Wonder, maturing 7 to 10 days ahead of other strains, and producing fruits over a long season. The plant is not so tall and erect in growth as regular California Wonder, but is very vigorous and prolific. The fruits usually are four-lobed and attractive in appearance, being smooth, dark green, uniform and thick meated. This early strain is rapidly growing in popularity.

Pkt. 10c; oz. 60c; ¼ lb. \$1.90; 1 lb. \$7.00

5 to 25 lbs. \$6.90 per lb.



*Early Calwonder*

**RUBY GIANT.** (70 days.) Our improved strain of Ruby Giant makes strong, vigorous, upright growing plants, with some disease resistance and withstanding poor growing conditions, yet producing a good crop. Dark green fruits are somewhat similar to World Beater in shape, being large, slightly tapering, and attractive in appearance. The thick flesh is sweet, mild, and of excellent quality. Productive, and bears over a long period, producing very large fancy fruits. No better strain of this popular type can be found anywhere for Florida growing conditions.

Pkt. 10c; oz. 50c; ¼ lb. \$1.50; 1 lb. \$5.50

5 to 25 lbs. \$5.40 per lb.

**NEAPOLITAN (Italian).** (50 days.) This is an early, medium large, prolific variety, growing in popularity on some northern markets where it is in demand among Italian people. This plant is wonderfully productive, carrying as many as forty fruits on a single plant. The fruits are upright in growth, averaging 4 inches long and 2 inches in diameter, three-lobed, yellowish green, becoming bright red when ripe. The flesh is mild and sweet.

Pkt. 10c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00

5 to 25 lbs. \$5.90 per lb.

**BLIGHT RESISTANT WORLD BEATER (Thick Wall Strain).** (70 days.) This strain of World Beater Pepper was developed in our breeding station at Bradenton, Florida for a more uniform, thicker walled, heavy yielding strain. In numerous trial plantings in different sections of Florida we have found this to be a heavy yielder of uniform, true to type, smoother, thicker walled fruits, which are larger in size and carry better to distant markets than regular World Beater which usually are very thin-walled. In some sections it has brought a premium on the market. The plants are erect in growth and hold the large heavy fruits well off the ground.

Pkt. 10c; oz. 65c; ¼ lb. \$2.15; 1 lb. \$7.50

5 lbs. to 25 lbs. \$7.40 per lb.

**ITALIAN "EL" (Green Neapolitan).** (65 days.) This green-fruited strain of Neapolitan is in increasing demand on northern Italian markets where it is often preferred to the regular yellow-fruited Neapolitan because of its peculiar sharp taste. Fruits are slightly larger and thicker fleshed than the regular yellow Neapolitan. The vigorous growing plants produce a heavy yield of light green fruits over a long season.

Pkt. 10c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00

5 to 25 lbs. \$5.90 per lb.

**ANAHEIM CHILI (Hot).** (80 days.) A popular, rather long, mildly pungent variety, not as hot as the small fruited Long Red Cayenne. The fruits are meaty, 6 to 7 inches in length and 1 to 1½ inches in diameter at stem end, green, ripening to a bright red. Often dried for winter use.

Pkt. 10c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00

5 to 25 lbs. \$5.90 per lb.

**HUNGARIAN YELLOW WAX (Hot).** (60 days.) A very early, hot sort, valuable for canners. Plants are rather dwarf in growth, and exceedingly prolific. Fruits pendent, medium slender, long, tapering, smooth and very hot or pungent. Color waxy yellow, changing to bright crimson at maturity.

Pkt. 10c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00

5 to 25 lbs. \$5.90 per lb.

**LONG RED CAYENNE (FINGER) (Hot).** (70 days.) This favorite hot variety produces large and productive plants. Fruit 4 to 5 inches long, ½ inch thick, slender, tapering, pendent, very hot or pungent, deep green in color, ripening to a brilliant red. Used largely for canning and in pickles.

Pkt. 10c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00

5 to 25 lbs. \$5.90 per lb.



*Blight Resistant World Beater*



# SEED POTATOES


Plant 600 to 900 pounds (10 to 15 bu.) per acre in 3-ft. rows.  
In the garden plant 12 pounds to 200 feet of row.

**Culture.** Plant in rows 36 to 42 inches apart, dropping a seed piece every nine to ten inches in the row. The seed piece should be at least one to one and one-half ounces in size. It is best to cut seed within one day of planting. Fertilizer should be put in seven or eight days before planting. Use from 1,500 to 2,000 pounds per acre, depending on type of soil. A 4-7-5 fertilizer mixture is a good standard for potatoes. On rich Everglades muck from 200 to 500 pounds per acre of 0-10-12 or 0-8-16 mixture applied in one application at planting time is a common practice. To control blight and increase the yield, spray with Dithane D-14, using 2 quarts Dithane D-14,  $\frac{3}{4}$  lb. zinc sulphate, to 100 gallons water. For control of insects and diseases see page 55.

**RED PONTIAC (Dakota Chief) (State Certified).** (80 days.) Packed in 100-lb. bags. This variety is the result of a cross, Red Bliss and Katahdin, made by the Michigan Experiment Station. In tests at the Subtropical Experiment Station in Homestead and elsewhere in Florida, it has greatly outyielded Red Bliss. The tubers of Red Pontiac are slightly longer, and have shallower eyes, but the color is a bright red like Red

Bliss. Pontiac tubers are tougher, stand shipment better and are better keepers than Bliss.

Write Plant City or your nearest Kilgore store for price on seed potatoes stating variety and quantity wanted.



Red Pontiac

## PUMPKIN

Plant two pounds seed per acre.

In the garden plant  $\frac{1}{2}$  ounce to 50 feet.

**Culture.** Plant 4 to 5 seeds in hills 8 to 10 feet apart one way and 5 feet apart the other way, when the weather gets warm. Can be planted in corn. Thin to two or three plants after well started. Give shallow and frequent hoeings until vines begin to run. Cut pumpkins when fully ripe.

Prices quoted are postpaid.

Write for special prices on larger quantities.

**KING OF THE MAMMOTHS.** (100 days.) Commonly known as a pumpkin, but is really the very large fruited Mammoth Chili squash, known as the giant among squashes. Fruits are globular, slightly ribbed, light yellow color mottled with orange. Flesh is solid, light yellow-orange, coarse, of fair quality. Grown mainly for stock feed or exhibition purposes. Fruits measure 20 to 22 inches across and weigh 60 to 80 pounds.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.



Small  
Sugar

**SMALL SUGAR.** (110 days.) The best variety for general use, and particularly desirable for pies. Fruits are round, flattened at ends. Skin is hard, smooth, somewhat ribbed, of deep orange color. Flesh is thick, sweet and dry, of bright orange color, and very high quality. Makes delicious pumpkin pies. Fruits measure eight to ten inches and weigh six to eight pounds.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

**CORNFIELD.** (118 days.) Prolific and vigorous, usually planted in cornfields. Extensively grown for making pies, for canning and for stock feed. Fruits are large, round to slightly flattened, with a hard, smooth surface, slightly ribbed, deep yellow or orange color. Fruits measure 12 to 14 inches and weigh 15 to 20 pounds.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

## RHUBARB

Sow 3 to 4 pounds in seedbed for an acre of plants, or plant 5 to 6 pounds per acre in field.

For the garden sow a packet in seedbed for 15 plants to set 25 feet of row.

**Culture.** Plant in the fall in rows  $2\frac{1}{2}$  to 3 feet apart and thin to about 18 to 24 inches in the row. It is best to plant seed in seedbed and transplant to field or garden. Rhubarb does best in well drained, fertile, moist soil, rich in humus. In Florida it is necessary to plant Rhubarb each season.

**VICTORIA.** (90 days from plants set in field or garden to edible stalks.) This vigorous growing variety produces thick, fleshy stalks of the best quality and flavor, and with attractive red color. It is the best variety for Florida.

Pkt. 10c; oz. 30c;  $\frac{1}{4}$  lb. 85c; 1 lb. \$3.00  
5 to 25 lbs. \$2.90 per lb.

## ROSELLE

Jamaica Sorrell or Florida Cranberry

(175 days from seed planting to ripe fruits.)

Plant 3 pounds per acre. In the garden plant one packet to 25 feet of row.

The red fruit is used when young for making jelly and jam. It is grown commercially for local markets, fruits being marketed in quart berry baskets.

**Culture.** Plant seed in field in March and April in Florida, when the ground is warm, spacing row 5 to 6 feet apart and thinning to about  $2\frac{1}{2}$  feet apart in the row, leaving one strong plant at a place. Roselle is an all-season crop requiring from five to six months from seeding to ripe fruits.

Pkt. 10c;  $\frac{1}{2}$  oz. 35c; oz. 60c;  $\frac{1}{4}$  lb. \$1.75; 1 lb. \$6.50  
5 to 25 lbs. \$6.40 per lb.



# RADISH

Plant 6 to 8 pounds of seed per acre.

In the garden make four plantings at different times. 1 packet to 25 feet of row at each planting

Our Radish Seed is now graded and sized for more accurate planting, better stands and more uniform maturity. Plant only Kilgore's graded radish seed for best results.

**Culture.** Plant in rows 12 to 18 inches apart and thin to 1 or 2 inches in the row. Successive plantings should be made every few weeks for a continuous supply. For best results, a rich, loose, moist sandy loam or muck soil is required. Radishes must be grown rapidly to be crisp, tender, and of best quality. Our radish seed stocks are produced each year from carefully selected transplanted roots. For insect and disease control see page 55.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**EARLY SCARLET GLOBE.** (21 days.) We have a perfected stock of this early variety that was grown from specially selected, transplanted roots. It is the earliest of all varieties; roots are globe-shaped, with a medium top. They are crisp and tender, of uniformly attractive bright, scarlet color. In great demand on all markets.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

**SCARLET TURNIP WHITE TIPPED (Sparkler).** (22 days.) A superior strain of this popular early variety. The roots are nearly round, being slightly flattened on the underside, deep scarlet, shading to a white tip on the bottom. They are very solid, crisp, tender, and sweet. Attractive in appearance.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

**BUCCANEER (Scarlet Globe Type) (New).** (21 days.) A new development of the Scarlet Globe type, developed for uniformity and attractive market appearance. The root is globe shaped, with a lustrous scarlet exterior, and with short top growth. Crisp-textured and a most uniformly attractive market radish.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.



*Early Scarlet Globe Radish.*

**LONG WHITE ICICLE.** (23 days.) The leading long radish. The straight roots are from 5 to 6 inches long, cylindrical, and very smooth and waxy white throughout. The flesh is mild, crisp, tender and of fine quality.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

# SPINACH

**Virginia Savoy**—Plant 12 to 15 pounds per acre in rows, or 20 to 25 pounds per acre broadcast.

In the garden make two plantings at different times.

Plant one packet to 50 feet of row at each planting.

*All prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**Culture of Virginia Savoy Spinach.** Successful for fall and winter planting. Does not grow well in hot weather. Requires a moist, rich soil. Plant seed in rows 16 to 20 inches apart, and thin to 6 inches apart in the row.

**VIRGINIA SAVOY (Blight Resistant).** (40 days.) Better for market than the smooth leaved varieties, because it does not pack as closely and thus is not as subject to heating as are the smooth leaved varieties, nor does it wilt or turn yellow as quickly before reaching the consumer. Developed for resistance to blight or mosaic disease. The plant is upright and compact in growth, with very thick, curly or savoyed dark green leaves.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

**New Zealand**—Plant 3 to 4 pounds of seed per acre  
In the garden plant one packet to 25 feet of row.

**Culture of New Zealand Spinach.** As seeds are very hard and germinate with difficulty, soak them in warm water for 24 hours to hasten germination. Plant in rows three feet apart, and drop three or four seeds per hill about 20 inches apart. Thin to one plant per hill.

**NEW ZEALAND.** (60 days.) Entirely distinct from the true Spinach. The plant is spreading, with numerous side shoots. It grows to a height of 1 foot and spreads over a wide radius. Leaves are thick, light green, rather small, broad and pointed. It grows well in hot weather, and is sometimes called "Summer" or "Hot-weather Spinach." The leafy tips of the branches are removed and new growth follows, making possible many cuttings. It does best on moist, rich land.

Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50  
5 to 25 lbs. \$1.40 per lb.

# SAGE

A popular garden herb used in flavoring or seasoning. Leaves should be cut as plant blooms, and should be dried quickly in the shade. Grows 18 inches tall.

**Culture.** Sage does well on a sandy soil. Plant seed in rows 15 to 18 inches apart. Thin to six inches in the row. The plant is a hardy perennial, and of easy culture in Florida.

Pkt. 10c; ¼ oz. 35c; 1 oz. \$1.00



# SQUASH

Squashes exhibit greater variation in plant and fruit character and hybridize or cross and become mixed more readily in seed production than any other vegetable. Consequently squashes are given special care and attention in our breeding grounds and production fields in the north and west. Pedigreed lines are established by covering buds on most desirable plants with bags to control pollination and prevent crossing. Market seed is grown from these pedigreed lines in fields well isolated from any other squash variety. In this way we are able to supply our customers with uniformly true-to-type strains of squash seed.

**Culture.** This is one of the quickest and easiest crops that can be grown, and usually proves very profitable. The earliest varieties begin bearing in six or seven weeks from seed. Squashes can be grown on almost any kind of soil. Use from 800 to 1,000 pounds of fertilizer per acre.

For bush varieties, have rows at least 3 to 4 feet apart and hills 2 feet apart in the row. Plant 3 to 5 seeds in each hill, and thin out to two plants when 3 inches high. Bush squash in Florida is a good paying crop for fall or spring. Plant for spring, during January, February and March; for fall, from August until October. Mosaic disease has become a most serious disease of squash in many sections of Florida. See Sanitation on page 50. Insect and Disease Control on page 54.

The number of days after each variety represents the average time required from seed planting to edible or marketable fruits.

*All prices quoted are postpaid  
Write for special prices on larger quantities.*

## BUSH VARIETIES

Plant 3 to 4 pounds of seed per acre.  
In the garden one packet to 50 feet.


**COCOZELLE (Italian Vegetable Marrow).** (50 days.) This is a true Italian Vegetable Marrow. When ready for market, the fruit is 10 to 12 inches long, slender, 1 to 1½ inches in diameter, cylindrical in shape, straight, smooth, dark green with lighter green stripes.

Our stock of this variety produces a bushy, open type of plant, with deeply cleft leaves, and a heavy yield of uniformly true-to-type fruits. We have devoted many years to the development of this superior strain, and we can say with full confidence that our Cocozelle is the earliest, most prolific, and uniformly true-to-type strain you can secure anywhere.

**Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50**  
**5 to 25 lbs. \$1.40 per lb.**



*Cocozelle Squash, or Long Slender Italian Vegetable Marrow*



*Early Yellow Summer Crookneck or Baby Crookneck*

**EARLY YELLOW SUMMER CROOKNECK (Baby Crookneck).** (45 days.) A very early and exceedingly prolific, light lemon-yellow color strain with a more open type of plant growth. The most popular and most extensively grown of any variety of squash in Florida. Fruits are small with thin, curved neck and uniform. Its rich, bright, lemon-yellow color gives it a very good market appearance, and it is in demand on all southern markets. Our strain of this variety is very early and exceedingly prolific, and has been selected for uniformly shaped fruits with a pronounced crook. No better strain of this most popular variety can be secured anywhere for Florida planting.

**Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50**  
**5 to 25 lbs. \$1.40 per lb.**

**BLACK ZUCCHINI (Kilgore's Improved Selection).** (45 days.) The plant of this improved selection, instead of developing a large, vigorous growing bush like the regular Black Zucchini, forms a short, single semi-runner, and produces a heavy yield of fruit over a long period. The plant is an open type of growth, with deeply cleft leaves, allowing the air and sunlight to circulate and prevent fruit rot. The fruits are dark green in color, six to eight inches long at market stage. A very early, prolific, improved strain of Black Zucchini.

**Pkt. 10c; oz. 20c; ¼ lb. 50c; 1 lb. \$1.75**  
**5 to 25 lbs. \$1.65 per lb.**

**EARLY PROLIFIC STRAIGHTNECK.** (42 days.) This development is ten days earlier and more productive than other strains of Straightneck Squash. The plants are vigorous, and the fruits are of brighter lemon-yellow color. Valuable for both home and market use. Because of its uniformity, earliness, high yield and attractive appearance this variety has grown greatly in popularity especially in northern markets. Makes a very attractive pack.

**Pkt. 10c; oz. 15c; ¼ lb. 45c; 1 lb. \$1.50**  
**5 to 25 lbs. \$1.40 per lb.**



## SQUASH—BUSH VARIETIES (Continued)

**WOOD'S EARLY PROLIFIC (Patty Pan Type).** (42 days.) The earliest, meatiest, and most prolific of the patty pan type. It will average seven days earlier than Early White Bush, matures its fruits quickly. The plants are prolific, producing round, smooth, deep, thick fruits, slightly scalloped, and of creamy-white color, usually with a light greenish cast, being bulged at both stem and blossom ends. A very good variety to grow where early squashes of the bush scallop type are desired for home use or for shipment.

Pkt. 10c; oz. 15c;  $\frac{1}{4}$  lb. 40c; 1 lb. \$1.25  
5 to 25 lbs. \$1.15 per lb.

**EARLY WHITE BUSH (Patty Pan Type).** (48 days.) This is a popular type of squash for shipment to northern markets. The smooth fruits are of pale green color when very young, becoming white as they grow, are round and flat but deep, with ridged or scalloped edges, measuring up to five or six inches across by three inches thick when ready to market.

Pkt. 10c; oz. 15c;  $\frac{1}{4}$  lb. 40c; 1 lb. \$1.25  
5 to 25 lbs. \$1.15 per lb.

**MAMMOTH WHITE BUSH (Patty Pan Type).** (50 days.) A large strain of the Early White Bush, fruits being about one-third larger, measuring to 8 inches in diameter. It is slightly later in maturing and not quite so prolific as Early White Bush. The fruits are thick, tender, and of good quality.

Pkt. 10c; oz. 15c;  $\frac{1}{4}$  lb. 40c; 1 lb. \$1.25  
5 to 25 lbs. \$1.15 per lb.

## SQUASH - RUNNING VARIETIES

Plant two pounds per acre of running varieties


In the garden one packet to 50 feet.

**Culture.** Cultural requirements are similar to bush squash (see page 30), but running varieties require much more room, planting in rows 6 to 8 feet apart and hills at least 3 to 4 feet apart. Plant 3 to 5 seeds in each hill and thin to two plants when three inches high. Generally grown as a spring crop in Florida.

In these types of squash the vines run 10 to 12 feet. The fruits are left on the vines until fully grown with a hard rind, and are then harvested by cutting the stem. Running squash may be stored in a cool, dry place until desired for use, but must be handled carefully to avoid bruising. Unlike bush squash, fruits of the running varieties may be stored and used boiled, baked or for making squash pies.



Royal Acorn (Mammoth Table Queen)



Butternut

**BUTTERNUT.** (80 days.) Vines produce pear-shaped yellowish brown or deep buff-colored fruits. Fruits weigh  $2\frac{1}{2}$  to 4 lbs., 8 to 10 inches long, and 3 to 4 inches in diameter at the largest diameter. The outer shell is smooth and hard, making this squash a wonderful shipper and keeper. The dry flesh is yellow, fine-textured, sweet, and of excellent flavor. Also excellent for making pies.

Pkt. 10c; oz. 30c;  $\frac{1}{4}$  lb. 85c; 1 lb. \$3.00  
5 to 25 lbs. \$2.90 per lb.

**BOSTON MARROW.** (95 days.) A very productive early maturing variety, making large fruit and maturing about two weeks earlier than the old Blue Hubbard which it resembles in shape. Our strain of this variety has an attractive deep-orange skin which is somewhat rough and quite hard. Thick, fine grained yellow flesh of excellent quality. Useful for canning and for fresh market. Can be held for a long time if fully mature when harvested and kept in a dry, cool, airy place.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00  
5 to 25 lbs. \$1.90 per lb.

**TABLE QUEEN (ACORN).** (60 days.) Extremely early for a running squash. Makes an enormous yield over a long season. Fruits dark green, weighing about 2 lbs., with sides deeply ribbed, and with a point at the blossom end. Yellow-colored flesh is very fine-grained and sweet when baked like Hubbard squash. It is a very good keeper and desirable for home and market use, also for shipping.

Pkt. 10c; oz. 15c;  $\frac{1}{4}$  lb. 45c; 1 lb. \$1.50  
5 to 25 lbs. \$1.40 per lb.

**ROYAL ACORN (MAMMOTH TABLE QUEEN).** (65 days.) To meet an increasing demand in Florida for a larger fruited strain of the Table Queen or Acorn Squash, we are offering this large-fruited strain. It is slightly later than regular Table Queen, not quite so prolific but the fruits are consistently larger and of a darker green color. In other respects this strain is similar to the regular Table Queen.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00  
5 to 25 lbs. \$1.90 per lb.

**ALAGOLD (Improved African Squash).** (100 days.) Resistant to most insects and diseases which commonly attack squash in the South. Average about 5 pounds in weight. The rind is of a dark green color, maturing to a deep yellow or buff color. A very good keeper.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00  
5 to 25 lbs. \$1.90 per lb.



# TOMATO

Sow  $\frac{1}{4}$  pound in seedbed to plant an acre, or  $\frac{1}{2}$  pound per acre in field.  
For the garden sow one packet in a seedbed for 50 plants to set 150 feet of row.  
If plants are staked, sow one packet in seedbed for 50 plants to set 75 feet of row.



*Jefferson (Wilt Resistant)*

This year again, like last year, tomato fields planted to Kilgore's Bred-Rite tomato seeds won first and second place in the judging of tomato fields at the annual Florida Tomato Festival at Ruskin, Fla. This year, the new Kokomo variety grown on Elsbury Farms at Ruskin won the coveted tomato sweepstakes award as the championship tomato field. Last year a field of our Improved Rutgers grown by W. D. Miller at Ruskin won this honor. Both years second place went to a field of Jefferson tomato, last year on the Powell & Braman Farms, and this year on the Beaudette Farms at Ruskin.

**Culture.** Tomatoes are grown on all kinds of soil, from a light sand to a muck. The land should be put in good condition. Lay off rows  $3\frac{1}{2}$  to 5 feet apart, open a furrow, and drill in 1200 pounds per acre of high grade fertilizer such as 4-7-5, and mix thoroughly with the soil. It is good policy to let this stand for seven to ten days and then plant the seed or set the plants. Seed may be planted direct in the field or preferably in a cloth-covered seedbed. Plants started in a seedbed should be 6 to 8 inches high when ready to transplant to the field in 4 to 7 foot rows and  $2\frac{1}{2}$  to 3 feet apart in a row, or if staked 16 to 20 inches in the row, rows 4 feet apart.

On the east coast of Florida, first plantings are made during August and continue until January. North and central Florida begin planting for a spring crop the middle to the latter part of December and continue to the middle of February. For a fall crop in central Florida, plantings are made in July and August.

For control of insects and diseases of Tomatoes see page 56. Spraying or dusting tomato plants with Dithane before the disease appears will help prevent the destructive late blight disease from developing.

The number of days indicated after each variety represents the time required from setting of plants in the field or garden to produce marketable fruits. It usually takes four or five weeks to produce plants for field setting.

*All prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**PONDEROSA (Brimmer).** (88 days.) A large, pink-fruited variety. Fruits semi-flat, solid with few seeds, and of splendid edible quality. Recommended only for home gardens.

**Pkt. 10c;  $\frac{1}{2}$  oz. 45c; oz. 85c;  $\frac{1}{4}$  lb. \$2.75; 1 lb. \$10.00**  
5 to 25 lbs. \$9.90 per lb.

**OXHEART.** (90 days.) A smooth, thick, tender and almost seedless pink-fruited tomato of delicious flavor. An ideal variety for the home garden and for the local market.

**Pkt. 15c;  $\frac{1}{2}$  oz. 65c; oz. \$1.25;  $\frac{1}{4}$  lb. \$4.00; 1 lb. \$15.00**  
5 to 25 lbs. \$14.90 per lb.

**PRITCHARD.** (75 days.) Vines medium dense, compact, erect, self-topping. Fruit matures very early, large, smooth, meaty, globular in shape. This variety is a heavy cropper and therefore requires rich or liberally fertilized soil. Does especially well on Everglades muck.

**Pkt. 10c;  $\frac{1}{2}$  oz. 30c; oz. 55c;  $\frac{1}{4}$  lb. \$1.65; 1 lb. \$6.00**  
5 to 25 lbs. \$5.90 per lb.

**JEFFERSON (Fusarium Wilt Resistant).** (76 days.) The plant is very vigorous and makes densely-leaved tall vines. It is more open and less subject to bunching than Rutgers. Produces well under favorable growing conditions, and well suited as a spring tomato and for staking in wilt infested soils. Under good growing conditions the fruits are globe-shaped, ripening to a uniform, attractive dark red color, with a good flavor, thick walled with a small core, and relatively free from cracking and catfacing. Fruits of this variety are slow-ripening so care must be used not to harvest them too immature. This variety holds up under adverse growing conditions better than any other variety. During each of the last two years our strain of this variety won second place in the judging of tomato fields at the annual Florida Tomato Festival at Ruskin, Florida.

**Pkt. 10c;  $\frac{1}{2}$  oz. 40c; oz. 75c;  $\frac{1}{4}$  lb. \$2.50; 1 lb. \$9.00**  
5 to 25 lbs. \$8.90 per lb.


**RUTGERS.** (82 days.) Our improved strain of this important variety is uniform in plant growth and produces uniform, smooth, thick wall, deep globe-shaped fruits. The plant is vigorous and a heavy producer of firm green tomatoes which ripen to a bright red, producing heavy, large yields over a long season. Very firm and free of puffs. Fruits of this strain size up well clear out to the end of the vine, and are smooth and free of cracks at the stem end. This variety is especially recommended for a spring crop in Florida and is well suited for light sandy soils and also for shallow, dry pine land on the lower east coast of Florida. Not suitable for deep, wet glade lands where it makes too much vegetative growth. Stands more rain as well as more drouth than other varieties. Many growers claim our Improved Rutgers makes the smoothest as well as the fanciest fruits of any strain of this variety on the market today.

**Pkt. 10c;  $\frac{1}{2}$  oz. 30c; oz. 55c;  $\frac{1}{4}$  lb. \$1.65; 1 lb. \$6.00**  
5 to 25 lbs. \$5.90 per lb.



*Rutgers Tomato*





*Kokomo (Wilt resistant Rutgers type)*

**KOKOMO (Wilt resistant Rutgers type) (new).** (80 days.) As a result of numerous trials in different sections of Florida during the past few years we are fully convinced that this new, Fusarium Wilt resistant variety of the Rutgers type is exceedingly well adapted to Florida growing conditions. It will do well wherever Rutgers does well. It is of the general Rutgers type, not as vigorous or bunchy in growth as Rutgers. Vines open up more, making it easier to spray or dust and to harvest. Kokomo sets a heavy crop of good size, smooth, deep globe-shape, meaty, thick-walled fruit, is a good shipper. It makes a very high percentage of smooth, fancy fruits and it is claimed to set well in hot weather. Is particularly valuable where Fusarium Wilt is a serious crop hazard, under which conditions Rutgers cannot be successfully grown. Kokomo grown on Elsbury Farms at Ruskin, Fla., won the coveted tomato sweepstakes award as the champion tomato field in the judging of tomato fields at the annual Florida Tomato Festival during the week of April 19, 1954.

**Pkt. 10c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.35; 1 lb. \$8.50**  
5 to 25 lbs. \$8.40 per lb.

**MARGLOBE.** (77 days.) This is an excellent strain of the old standard type Marglobe. It produces a vigorous plant resistant to nailhead rust and Fusarium Wilt. Fruit deep globe-shaped, smooth and thick walls, free of navel and blossom-end scars. Ripens to a beautiful, deep scarlet-red color. Well adapted to and recommended for north Florida.

**Pkt. 10c; ½ oz. 30c; oz. 55c; ¼ lb. \$1.65; 1 lb. \$6.00**  
5 to 25 lbs. \$5.90 per lb.

**GROTHEN'S GLOBE.** (70 days.) An early maturing, high yielding variety especially well adapted for fall planting in Florida. The plants are of open growth with rather small foliage requiring rich, deep, moist land for best results, maturing fruits about 12 days earlier than Rutgers, and ripening to a beautiful red color. A heavy producer of smooth, deep globe-shaped, large fruits, sizing up well clear to the top of the plants, producing a high percentage of fancies. The fruits are firm and solid, and the interior ripens slowly, making this variety an exceptionally good shipper. Earliness and high yields of large fruits, of excellent shipping quality, are its main factors. It does not do so well on light sandy soils, but is especially recommended for a fall, winter and early spring crop in Florida and is well suited to deep, wet glades, hammock, and rich sandy loam soils.

**Pkt. 10c; ½ oz. 35c; oz. 60c; ¼ lb. \$1.75; 1 lb. \$6.50**  
5 to 25 lbs. \$6.40 per lb.

**HOMESTEAD 24 (Wilt Resistant) (new).** (70 days.) A selected strain of Homestead tomato. Our aim in developing this strain has been elimination of off types in the original Homestead leading to uniformity of plants and fruits and a better crop. Compared with Rutgers, the plants of Homestead 24 are similar but semi-determinate, earlier in maturity and resistant to Fusarium Wilt. Fruits are similar, very free from puffiness and cracking, meaty and firm, excellent for green wrap. Homestead 24 is a heavy yielder and will make good crops even on wilt-infested soils. It has done well in the Homestead area, both on heavy marl soil and on the rockland. Seed supply limited.

**Pkt. 10c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.50; 1 lb. \$9.00**  
5 to 25 lbs. \$8.90 per lb.

**MANALUCIE (Wilt Resistant) (new).** (82 days.) This new, promising, vigorous growing, disease resistant variety was developed and recently released by the Gulf Coast Experiment Station at Bradenton, Fla. It possesses multiple resistance, being very resistant to four serious Florida tomato diseases, namely, Fusarium Wilt, gray leaf spot, early blight, and leaf mold. It is also mildly resistant to black spot stem canker caused by the Phoma fungus. Makes a very strong, vigorous growing plant with large leaves and large, strong stems. Fruits are late, but are large size throughout the harvest period. Smooth and relatively resistant to cracking. Firm and fleshy, and good edible quality. Slow ripening and must be allowed to fully mature before harvesting. Ripens to a deep red color. High-yielding, and a good shipper. Shows much promise, especially as a spring tomato on old tomato land.

**Pkt. 15c; ½ oz. 60c; oz. \$1.00; ¼ lb. \$3.15; 1 lb. \$11.50**  
5 to 25 lbs. \$11.40 per lb.

**KILGORE'S WILT RESISTANT GROTHEN'S GLOBE.** (70 days.) This variety has been developed from a cross of Pan American and regular Grothen's Globe which was made several years ago by the Kilgore Seed Company. Wilt Resistant Grothen's Globe possesses all the good qualities of the regular Grothen's Globe, and in addition it is very resistant to Fusarium Wilt, although slightly earlier and more vigorous than regular Grothen. In numerous comparative trials between Wilt Resistant and regular Grothen's Globe, even where wilt was not a factor, the Wilt Resistant strain yielded just as high as the regular strain. This Fusarium Wilt Resistant strain is especially recommended for old, wilt infested soils where the regular Grothen's Globe and other varieties cannot be grown successfully. Is not resistant to all wilts, but is highly resistant to the most serious wilt disease of tomatoes caused by the Fusarium soil fungus.

**Pkt. 10c; ½ oz. 40c; oz. 75c; ¼ lb. \$2.25; 1 lb. \$8.00**  
5 to 25 lbs. \$7.90 per lb.



*Grothen's Globe*



# What Will Your Harvest Be ?

The answer to that question depends principally on the kind of seeds you plant. Kilgore's Bred-Rite Seeds Are Bred and Developed in Florida for Florida. This is why The Kilgore Seed Co. is referred to as "Florida's Leading Seedsmen."



*Partial view of Experimental Farm and Breeding Station at Bradenton, Florida.*

Samples of different stocks of Kilgore seeds are planted from time to time on an Experimental Farm, at Bradenton, Florida. In addition, there also are included in these trials, samples from other seed dealers and seed growers to compare with Kilgore stocks. This shows just how Kilgore's strains and varieties compare in earliness, yield, uniformity, trueness to type and adaptability with others growing side by side under identically the same conditions in Florida. It also enables the company to develop new or improved varieties of distinct merit for Florida conditions, which we offer to our customers as soon as a supply of seeds can be produced. Modern, scientific breeding methods are employed both in maintaining and improving seed stocks. New products are carefully tested under Florida conditions, and superior stocks are then sent to our seed production fields in the North and West. Careful notes are kept on all experimental farm results which enable

us to check up with the constant improvements which are being made in standard varieties of vegetables. Our principal objective is to breed, develop or select strains and varieties of vegetable seeds adapted to Florida growing conditions, which differ materially from those found elsewhere in the United States. A well trained and experienced plant breeder supervises this work.

In addition, the Kilgore Seed Company personnel is in constant contact with the scientific research workers of the Florida Agricultural Experiment Station and field laboratories throughout the state where plant pathologists, horticulturists and plant breeders are doing a splendid work in developing new and improved vegetable varieties especially adapted to the climate and growing conditions of Florida, which we introduce to Florida growers as soon as available.

## HIGH VITALITY SEED MUST BE PLANTED TO SECURE A GOOD CROP

*Kilgore's vegetable seeds are air-conditioned and tested.*

*This means higher vitality, better germination and stronger plants.*

The Kilgore Seed Company, has in its central warehouse in Plant City, Florida, the most up-to-date dehumidifying machinery in order to maintain high vitality in all Kilgore seed. Surplus seed supplies in our stores are also kept in specially constructed dry rooms where the humidity is maintained at a uniformly low degree.

We are now putting up certain kinds of vegetable seeds which naturally lose their germination rather rapidly, such as celery, eggplant, pepper and hybrid onion, in special sealed air-tight and moisture-proof, one-pound containers after the seed has been treated with a seed protectant and the moisture content of the seed has been artificially reduced by a special process. The sealed cans should be **opened only as needed for planting** in order to retain the vitality of the seed until planted. If kept in unopened cans the seed will retain

its vigor and germination even under Florida conditions of high humidity and temperature.

In addition to keeping our seeds in artificially dried storerooms, we have a seed testing laboratory where samples of all our seed stocks are tested at regular intervals to further insure our customers against planting low vitality seeds. In our seed testing laboratory, tests are made every six weeks of all seed stocks in our warehouse, mail order department and fourteen Florida stores. We know at all times the correct germination of all seed we have on hand. If any seed stocks are found to be dropping in germination they are immediately removed from our warehouse and stores and are destroyed.

We are taking all these precautions and giving our customers this extra service without any additional cost, in order to provide in Kilgore's Bred-Rite Seeds the best that money can buy. Play safe and get your seed from the Kilgore Seed Company, Florida's leading seedsmen.



# TURNIP

Plant 2 to 3 pounds of seed per acre.  
In the garden make three plantings at different times.  
1 packet of turnip seed to 50 feet of row.

**Culture.** Plant in rows 15 to 20 inches apart. Thin early to one inch, because any over crowding will result in misshapen, slow growing, poor quality roots. Final thinning should be 4 to 6 inches apart. Thinnings may be used for turnip greens. For best quality, turnips should be grown rapidly in rich, moist soil. When turnips are grown for commercial canning and freezing, where only the tops are used and harvested with machinery, plant in double rows 4½ to 5 feet apart, 14 inches apart in the double row. 6 to 8 lbs. of seed is required, no thinning. Turnips can be planted every month in the year.

The number of days for each variety is the time required to produce edible roots from seed plantings.

*Prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**TOKYO MARKET.** (35 days.) Extremely early in maturity. Attractive, white, globe-shaped roots. Small foliage grows upright, and is somewhat aphid resistant. Our special stock of this variety is exceedingly uniform, which makes it a good variety to grow for market. Exceptionally fine quality.

Pkt. 10c; oz. 15c; ¼ lb. 40c; 1 lb. \$1.25  
5 to 25 lbs. \$1.15 per lb.

**PURPLE TOP MILAN (EARLY FLORIDA).** (35 days.) The roots are of medium size, smooth, flat, and white, with a bright purple top, and extremely early. Recommended as an extra early home garden and local market variety.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

**WHITE GLOBE.** (44 days.) A handsome, white, very large, perfectly round turnip of fine quality. It has small tops, grows rapidly, and gets very large, producing a heavy yield per acre. Resembles Purple Top White Globe except for color.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

**SHOGGIN (FOLIAGE) TURNIP.** (50 days.) An aphid-resistant foliage turnip which resists the hot, dry weather. It produces an immense yield of upright growing foliage, always in good demand during late spring and summer in Florida. The roots are white, semi-globe shaped, but more largely planted for the tops which are large, thick, succulent, tender, and of finest mild flavor.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.



*A typical root of Purple Top White Globe*

**PURPLE TOP WHITE GLOBE.** (50 days.) Ours is a very uniform and well colored strain of this popular variety. Roots are almost globe-shaped, and the flesh and skin are white, with a beautiful purple or red at the stem end above ground. Fine grained and sweet, of superior quality. Grown extensively in Florida for shipment to northern markets.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

**WHITE EGG.** (45 days.) An early, oval or egg-shaped variety with clear white roots of medium size. The flesh is white, firm, fine grained and sweet.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.

## RUTABAGA


Plant 1 pound per acre.

In the garden plant one packet to 100 feet of row.

**Culture.** Rutabaga is best grown as a late fall and winter crop. Planted in rows 2 feet apart, and the plants thinned to about 5 to 6 inches in the row.

**AMERICAN IMPROVED.** (90 days.) The smooth roots attain a much larger size than turnips; are almost globular, being slightly oblong, pale yellow with purple top, and with a very small neck. Flesh is yellow, solid, tender, not woody and stringy.

Pkt. 10c; oz. 15c; ¼ lb. 35c; 1 lb. \$1.00  
5 to 25 lbs. 90c per lb.



*Shogoin (Foliage) Turnip*



# WATERMELON

**Culture.** Watermelons may be grown on nearly all kinds of Florida soils that have good drainage, but rolling, sandy pine land is preferred. In general, new land is best for this crop in order to avoid diseases. The field should be checked off, planting in hills 8 feet apart each way or 7 by 9 feet. Seed should be planted  $\frac{1}{2}$  to 1 inch deep, and plants thinned to 2 or 3 plants per hill when they are still small and when all danger of frost is past, later thinning to 1 plant per hill for best results. The most successful growers apply 600 to 800 lbs. of fertilizer mixture about a week before planting the seed. A second application of 400 to 600 lbs. of fertilizer or top dressing may be made if necessary when the vines begin to run. The first planting of watermelons for an early crop in Florida should be made about January 1st and successive plantings may be made up to about the middle of February in central and south Florida. Plantings in north Florida are made 2 or 3 weeks later. For earlier melons use **Hotkaps**. (See page 75).

The leaf, stem and fruit disease of watermelons, anthracnose as well as gummy stem blight, the spores of which diseases are frequently present on melon seed, are killed by treating the seed with Spergon Seed Protectant. (See pages 52 and 62.) If for disease and insect control see page 56.

For pollination of watermelons, see second paragraph under cucumber culture—page 16.

All our melon seed is now treated with Spergon to control seed-borne diseases and to give better stands, especially if planted early in cold ground.

The number of days after each variety indicates the time from seed planting to ripe fruit.

*All prices quoted are postpaid.*

*Write for special prices on larger quantities.*

**FAIRFAX (New).** (92 days.) This new variety was developed by the U.S.D.A. Vegetable Breeding Laboratory at Charleston, S. C. It has the general appearance, shape and color of Garrison but is distinguishable by a darker green stripe on the light green base. Also it has a hard rind in contrast to the soft rind of Garrison, thus making it a good shipping melon, but the flesh texture and edible quality (sugar content) do not compare with Garrison. It produces a melon up to 50 pounds, but will have an average weight of 30-35 pounds. Relatively resistant to hollow-heart and develops less "gourdneck" and blossom end rot than most long melons. Fairfax is resistant to both Fusarium Wilt and anthracnose. The vines are vigorous and hardy and show considerable capacity to withstand adverse weather.

**Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00**  
**5 to 25 lbs. \$1.90 per lb.**



*The Congo (U.S.D.A. 46-40) (Anthracnose-Resistant) Watermelon*

Plant 1 pound of seed per acre.

In the garden plant 1 packet to 75 feet of row for a family of 3 to 4 people.

If interested particularly in watermelon production write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida for a free copy of Circular 96, Revision A, entitled "Watermelon Production Guide." Similarly, if interested in production of any other vegetable crop, write the Experiment Station for a free copy of their production guide on that crop, being sure to indicate the crop on which you desire guidance.

**CONGO—FLORIDA STATE CERTIFIED — (Anthracnose Resistant).** (90 days.) This State Certified strain of Congo comes from fields inspected and certified for trueness to type and freedom of mixture by the Florida State Department of Agriculture. **THIS CERTIFIED SEED IS SOLD ONLY IN 1 LB. SEALED BAGS** with the official Florida State Department of Agriculture certification tag attached to each sealed bag.

**1 lb. \$2.25; 5 to 25 lbs. \$2.15 per lb.**

**CONGO (U.S.D.A. 46-40) (Anthracnose Resistant).** (90 days.) This outstanding disease-resistant variety was developed and introduced by the U.S.D.A. Southeastern Vegetable Breeding Laboratory, Charleston, S. C. It is resistant to both Fusarium Wilt and Anthracnose diseases. The vines remain green longer than other varieties to protect the melons from sun burning. In order to be sure Congo retains its resistance to anthracnose, it is very important to get seed produced in Florida where anthracnose is prevalent every year. It is of exceptional importance not only because of its anthracnose-resistance, but also because of its exceedingly hard, tough rind which resists breakage and bruising in transit. In addition to these two good qualities, it possesses many other desirable features. It is a heavy yielder of large size melons, under good growing conditions ranging from 32 to 40 pounds. Similar in size and shape to Garrison but having a darker green rind with darker green stripes. It is free of the tendency to produce gourd necks. Flesh of attractive bright red color, of excellent quality and flavor, ripening well out

to the rind, with small white seeds having light tan smears, and very attractive interior appearance. This variety does not need a heavy fertilization as other varieties, and little, if any, side dressing with soda or nitrogen fertilizer should be used because heavy nitrogen may cause hollow heart or separation of the flesh. Melons of this variety should be harvested just before full maturity to avoid hollow hearts. This is one of the best varieties of watermelon for shipment yet introduced, and it has a big future.

**Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c;**  
**1 lb. \$1.75**  
**5 to 25 lbs. \$1.65 per lb.**



WE HAVE SPECIALIZED IN  
WATERMELON SEED FOR  
MANY YEARS

OUR MELON SEED IS PRODUCED  
FROM SPECIALLY SELECTED  
STOCK SEED



*Florida Giant (Black Diamond or Cannon Ball). A standard, high-yielding variety for Florida*

**FLORIDA GIANT (BLACK DIAMOND or CANNON BALL).** (92 days.) The vines are of vigorous growth and produce large, nearly round, solid dark green-skinned melons, with thick rind and firm, rather coarse grained and fibrous, sweet, deep red, attractive flesh with small black seeds. Very hardy, and withstands poor growing conditions better than most other varieties. Because of the dark green rind, it is in great demand, always having a fresh appearance. Has grown in favor among commercial melon growers in Florida. Also a desirable home garden variety.

Pkt. 10c; oz. 15c;  $\frac{1}{4}$  lb. 45c; 1 lb. \$1.50  
5 to 25 lbs. \$1.40 per lb.

**CHARLESTON GRAY (51-27) (Wilt and Anthracnose resistant).** (New.) (92 days.) This is one of the most promising of the new melon varieties recently introduced. Developed by the U.S.D.A., Southeastern Breeding Station at Charleston, S. C., it is highly resistant to both Fusarium wilt and Anthracnose. It makes a long, grey melon very slow to "sunburn", with a tough but not particularly thick rind, very striking in appearance when cut, with red meat and black seed. It is of delicious flavor and quality. A good yielder and very uniform in size, averaging under good growing conditions about 28 to 35 lbs. Unlike other varieties it can stay in the field in good condition for some days after it is ripe. This, along with its high resistance to the two most serious diseases of watermelons, makes it especially valuable for home garden use and to hold for a strong market. This new variety offers much promise as a high quality, disease-resistant, good shipping type melon for Florida.

Pkt. 10c; oz. 45c;  $\frac{1}{4}$  lb. \$1.40; 1 lb. \$5.00  
5 to 25 lbs. \$4.90 per lb.

**NEW HAMPSHIRE MIDGET (ICEBOX TYPE) (New).** (65 days.) This small icebox type melon, weighing 3 to 5 lbs. is attracting attention in Florida for home use, roadside stands, and local markets. It is the earliest of all watermelons making ripe fruits in from sixty to seventy days from seed. The vines lack vigor, but are prolific, setting many small melons, round to slightly elongated in shape with a light grey mottled rind, which is very thin and tender. Flesh is a beautiful

deep red with small black seeds. This has been found to be the best of the "icebox type" melons for Florida.

Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 50c; 1 lb. \$1.75  
5 to 25 lbs. \$1.65 per lb.

**BLACK DIAMOND—OKLAHOMA CERTIFIED.** (92 days.) Our certified Black Diamond or Cannon Ball is produced for us by one of the best watermelon seed growers in Oklahoma and the fields are thoroughly inspected and certified by the Oklahoma Crop Improvement Association. No finer strain of seed of this popular variety can be secured anywhere at any price. Not a melon is shipped from our Oklahoma seed growing fields, and seed is saved from only the earliest maturing, largest sized and best shaped melons. **This certified seed is sold only in 1-lb. sealed bags** with the official Oklahoma Crop Improvement Association Certification tag attached to each sealed bag.

1 lb. \$2.50; 5 to 25 lbs. \$2.40 per lb.

**SUPER BLACK DIAMOND (Yellow Belly Strain).** (92 days.) Our seed of this strain was harvested from only the largest, most uniform melons in the field. Melons are round in shape, like the regular Black Diamond, but of a darker blue-green color bottom of the melon being a deep yellow. Seeds blackish brown little darker than the regular Black Diamond, and smaller. Rind somewhat thinner, and a little tougher. Meat very red, larger heart, and better quality. Anthracnose does not bother this type as much as the regular Black Diamond melon. The Super Black Diamond is a heavy bearer. Florida melon growers will be highly pleased with this new strain because of its attractive dark green rind and superior edible quality.

Pkt. 10c; oz. 25c;  $\frac{1}{4}$  lb. 75c; 1 lb. \$2.50  
5 to 25 lbs. \$2.40 per lb.

**GARRISON.** (95 days.) We offer the genuine strain of this prized variety. Fruits are uniformly cylindrical in shape, very large, long, with a thin, tender rind. The color is white and green striped, the green stripes usually being broken. Delicate flavor, sweet and tender meat, and grows to weigh 40 to 50 pounds, if the vines are pruned to one or two good shaped melons per plant, pulling off all other melons at least twice a week. Because of its high edible quality and large size it is popular and commands premium prices. It is not a shipping melon because of the brittle, easily broken rind, but is grown extensively for local markets.


Pkt. 10c; oz. 20c;  $\frac{1}{4}$  lb. 60c; 1 lb. \$2.00  
5 to 25 lbs. \$1.90 per lb.



# FIELD AND ROASTING EAR CORN

(For Sweet Corn, See Page 18.)

Plant 7 pounds ( $\frac{1}{2}$  peck) per acre or 56 pounds (1 bushel) to 8 acres.



*Dixie 18 Hybrid Corn*

Kilgore's field corn seed is grown in isolated fields from the highest yielding pedigreed strains. Only large ears with deep grain, straight rows and good tip and butt are saved. Each ear is hand-tipped and butted, shelled and carefully graded to remove cracked oversized and undersized grains.

To increase yields with field corn in Florida two things are essential, namely closer spacing and ample fertilizer. On good land 20 to 26 inch spacing in 38-inch rows is best with 500 lbs. per acre of fertilizer at planting time followed by side dressings of nitrate of soda up to 300 lbs. per acre about 5 weeks after planting.

The number of days indicated after each item represents the time from seed planting to harvest under average spring conditions in Florida.

*Write for special prices on larger quantities.*

*All prices quoted are subject to change without notice.*

**CUBAN YELLOW FLINT.** (96 days.) This special strain is deep rooted, therefore can be given closer spacing than other varieties. If given close spacing it will yield as much as any prolific corn with two ears on 20% to 25% of the plants. Produces larger ears, longer grains, and heavier yields than other strains of this variety. Weevil resistant due to the long, tight, heavy and very tough husks. Seed is lemon to orange-yellow in color, with about 95% white cobs, 5% red cobs. Produces well in wet or dry seasons, and is recommended in all parts of Florida where extreme resistance to weevil damage is desired.

$\frac{1}{4}$  lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid

Not prepaid, 15 lbs. (pk.) \$1.75; 56 lbs. (bu.) \$6.50

**IMPROVED GOLDEN DENT.** (86 days.) A wonderful yielder, and does splendidly here in Florida. It is planted extensively for early use, but is too soft for a general crop. Has large, deep, broad grains, and is desirable for early feed.

$\frac{1}{4}$  lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid

Not prepaid, 14 lbs. (pk.) \$1.50; 56 lbs. (bu.) \$5.50

## ABOUT HYBRID SEED CORN

Hybrid corn is the result of crossing certain inbred, or self-pollinated lines of corn in various combinations. Maximum benefits are derived from the first generation of hybrid seed. Later generations are less vigorous and uniform. Hybrid corn seed should be obtained each year from producers who are specialists in hybrid seed corn production. Seed saved from a field of hybrid corn will always produce disappointing results.

**DIXIE 18 HYBRID (Certified).** (93 days.) This yellow hybrid field corn was developed at the Coastal Plain Experiment Station in south Georgia in cooperation with the U. S. Department of Agriculture. Dixie 18 is the best yellow hybrid corn for central and north Florida and south Georgia; a heavy yielder, averaging 2 ears per plant. Yields of 80 to 100 bushels per acre have been reported. The plants are relatively short, sturdy, and very vigorous, with a deep root system, resisting wind damage, lodging and drouth. The ears are uniform and borne low down on the plant. It makes a semi-hard corn and is very desirable for "hogging off". The grain is yellow and intermediate in texture being harder than Whatley. Has good resistance to weevil.

$\frac{1}{4}$  lb. 15c; 1 lb. 50c; 2 lbs. 85c; postpaid

Not prepaid, 14 lbs. (pk.) \$3.00; 56 lbs. (bu.) \$11.50

**FUNK'S G-737A HYBRID.** (95 days.) This new yellow corn has given excellent results in numerous commercial trials in the Everglades muck and offers fine possibilities for other sections of the state. It consistently makes high yields (two ears per stalk) of attractive, medium large ears, well covered to the tips with hard, semi-flint yellow kernels, with extra heavy shuck, very weevil resistant and of excellent keeping quality. Makes strong, sturdy stalks which stand drouth better than many others.

$\frac{1}{4}$  lb. 25c; 1 lb. 60c; 2 lbs. \$1.00; postpaid. Not prepaid, 14 lbs. (pk.) \$3.25; 56 lbs. (bu.) \$11.90

25 bu. \$11.40 per bu.



*Typical ears of Funk's G-737 Hybrid showing heavy husks and excellent resistance to weather and weevil damage.*



All prices quoted are subject to change without notice.

## CORN - Roasting Ear

(For Sweet Corn, See Page 18)

Plant 7 pounds ( $\frac{1}{2}$  peck) per acre. In the garden make two plantings at different times. Plant  $\frac{1}{4}$  pound to 200 feet of row.

Write for special prices on larger quantities.

**Culture.** Same as for sweet corn (see page 18) except given a little wider spacing—in rows 3 to  $3\frac{1}{2}$  feet apart, and 18 to 24 inches in the row. Grows more vigorously and is a little later in maturing than most varieties of sweet corn. For control of corn earworm see page 54.

The number of days after each variety represents the time from seed planting to the first harvest under average conditions.

**OKLAHOMA SILVERMINE.** (82 days.) A heavy yielder of large ears. The cobs are covered with 14 to 18 rows, and run from 11 to 12 inches in length. Very resistant to worm damage. An excellent variety for shipping, with a heavy dark green husk that holds its color in shipment and on the market. This is the standard roasting ear variety grown in central Florida.


$\frac{1}{4}$  lb. 15c; 1 lb. 50c; 2 lbs. 85c, postpaid

Not prepaid, 14 lbs. (pk.) \$2.00; 56 lbs. (bu.) \$7.00

**SOUTHERN SNOWFLAKE.** (80 days.) The ears are large, 10 to 12 inches long, well filled with 16 rows of large, broad kernels, and of very attractive appearance. The long tight husks protect the ears from worms.

$\frac{1}{4}$  lb. 15c; 1 lb. 50c; 2 lbs. 85c, postpaid

Not prepaid, 14 lbs. (pk.) \$1.75; 56 lbs. (bu.) \$6.50



Oklahoma Silvermine

## Lawn and Golf Course Grasses for Florida

All grass seed prices quoted are f.o.b. Plant City except lb. lots which are postpaid. Write for prices on larger quantities.

*Kilgore's Lawn Grasses are especially adapted to Florida conditions. Our grass seeds are recleaned, well-bagged, and of highest quality stocks.*

All grass seed prices quoted are subject to change without notice. Bitter Blue-Stemmed St. Augustine and Centipede Grass are used only for lawns. The other grasses listed and described on this page are used for lawns, golf courses and for pastures. For information on "When to plant grass seeds in Florida" see page 5. For lawn and golf course fertilizers see page 67. For lawn grass pesticides and supplies see page 57 and 74.

### BERMUDA GRASS

**Unhulled**—For lawns sow 2 to 4 pounds per 1000 square feet. For pasture, sow 25 pounds per acre.

**Hulled**—For lawns sow  $1\frac{1}{2}$  to 2 pounds per 1000 square feet. For pastures sow 15 to 25 pounds per acre.

Bermuda grass is generally considered best for maintaining a beautiful and perfect lawn. It is of fine texture, can be maintained green practically the year around, is a rapid grower and produces a lush, green carpet of turf though it requires careful attention in maintenance. Frost will turn it brown for a short period, but it soon regains its green color. It prefers loam or clay soil and will grow well on sand or muck. 20 to 30 pounds per 1000 square feet of a complete fertilizer such as Vigoro or Vertagreen should be applied twice a year. Frequent weeding is desirable. Although

Bermuda grass will respond to watering, it will stand considerable drouth. A thorough watering every 5 to 7 days during periods of drouth should be sufficient. The grass may be clipped closely ( $\frac{3}{4}$  to 1 inch) and frequently. Bermuda will not grow in shade. It spreads by surface runners and by underground root stalks. Seed can be planted from March to November.

### UNHULLED BERMUDA GRASS SEED

1 lb. 80c; 10 lbs. \$4.80; 25 lbs. \$11.50; 100 lbs. \$45.00

F.O.B. Plant City, except pound lots which are prepaid.

### HULLED BERMUDA GRASS SEED

The almost watertight husks of the seed have been removed, facilitating quicker, surer results, the seed germinating more uniformly and in a much shorter time with proper moisture conditions.

1 lb. 90c; 10 lbs. \$5.50; 25 lbs. \$13.25; 100 lbs. \$52.00

F.O.B. Plant City, except pound lots which are prepaid.

### PENSACOLA BAHIA GRASS FOR LAWNS

For Pasture See Page 41.

For lawns sow 100 to 150 pounds per acre or 2 to 3 pounds per 1000 square feet. For pastures sow from 10 to 15 pounds per acre.

This variety of pasture grass is sometimes used for low maintenance lawns. It is of fine texture and will remain green most of the winter (longer than Centipede grass). Will tolerate dry or infertile soils, but makes

(Continued on page 40)

For further information on starting, fertilizing and care of Florida lawns write for a free copy of "Kilgore's Flower Guide for Florida." This valuable book also contains complete information on growing flowers in Florida. Also write the Florida Agricultural Experiment

Station, Mailing Department, Gainesville, Florida for a free copy of Bulletin 518 entitled "Lawns in Florida" and for Circular S-42 entitled "Insects and other Pests of Lawns and Turf."





*Beautiful lawns can be grown in Florida to enhance the appearance of your Florida home.*

#### LAWN AND GOLF COURSE GRASSES FOR FLORIDA—(Con'd.)

an open, generally less attractive turf or lawn than regular lawn grasses. It is somewhat difficult to mow, especially when rapid growing seed shoots develop during the summer, but should not be mowed closer than 1½ to 2 inches. It requires fertilization in mid-spring and late summer, with a complete fertilizer such as Vigoro or Vertagreen. It is susceptible to damage from salt spray, but is relatively free of insect and disease troubles. Spring and early summer are the best periods for planting. Seeds are somewhat slow to germinate, sometimes requiring several weeks. The seedbed should be kept moist until germination occurs. For best distribution of seed, it should be mixed with dry sand. Sow seed at the rate of 2 to 3 pounds per square feet. "Common" Bahia should never be used for lawns.

1 lb. 75c; 10 lbs. \$4.30; 25 lbs. \$10.25; 100 lbs. \$40.00

#### ITALIAN RYE GRASS

*For Lawns Sow 250 to 300 Pounds Per Acre.*

*Or 6 to 8 Pounds Per 1000 Square Feet.*

*For Temporary Winter Pasture Sow 50 Pounds Per Acre.*

This is a valuable and desirable fall and winter grass for Florida, but it does not make a permanent lawn. It dies out in warm weather of late spring and early summer. For quick covering of lawns and golf courses in winter it is unsurpassed. Do not plant in Florida before October or later than March. It will make a vigorous, luxuriant growth throughout the fall, winter and early spring but must have plenty of water. A common practice in Florida is to seed Italian Rye grass on other brown lawns or in brown unsightly spots of old lawn in the fall and winter for a velvety green all-winter lawn. Plant ½ lb. to 100 square feet right on top of old lawn after loosening the surface soil. Keep soil moist until seed sprouts.

1 lb. 50c; 10 lbs. \$1.55; 25 lbs. \$3.40; 100 lbs. \$12.50

#### CARPET GRASS

*For Lawns Sow 100 to 150 Pounds Per Acre or 2 to 4 Pounds Per 1000 Square Feet.*

*For Pastures 25 Pounds Per Acre.*

Carpet grass is particularly well adapted to low moist soils and does not require much fertilization, but needs plenty of moisture. This grass is not suited to drouthy conditions or to sweet soils with a high pH. It turns brown in the winter, is not at all salt tolerant nor will it stand much shade. Keep mowed to a height of 2 inches. Plant in the spring from March to July, or in fall from October to December.

1 lb. 80c; 10 lbs. \$4.80; 25 lbs. \$11.50; 100 lbs. \$45.00

#### BITTER BLUE-STEMMED ST. AUGUSTINE GRASS

*Use one bushel of sprigs per 200 to 300 square feet.*

This is one of the most popular lawn grasses for Florida. It is quick to establish, rather coarse in texture, but easy to mow, and remains green the year around. It should be cut at a height of 2 inches. Severe cutting back is quite harmful. It should be fertilized two or three times a year at the rate of 20 pounds per 1000 square feet with a complete fertilizer such as Vigoro or Vertagreen. St. Augustine is our best shade grass and will grow on alkaline as well as slightly acid soils. It is very salt tolerant and can be grown along the beaches where a good soil base is provided. It can be grown by vegetative propagation only. Sprigs should be planted in well prepared soil, setting about 6 x 9 inches apart. For chinch bug control in St. Augustine lawns see page 57.

1 bu. \$2.50; 5 bu. \$2.40 per bu.; 10 bu. \$2.25 per bu.

*Not prepaid. Available April to November. Shipped only by express direct from a Florida nursery.*

#### CENTIPEDE GRASS

*Use one bushel of sprigs per 300 to 400 square feet.*

This is considered the best low maintenance grass for Florida. It is adapted to a wide range of soils including high sand hills, and will tolerate infertile acid soils, can get by on a minimum of fertilizer and water, and is relatively free of insect pests and diseases. It will not do well on alkaline or sweet soils but makes best growth on acid soils. A light spring and/or fall application of a complete fertilizer such as Vigoro or Vertagreen—10 to 15 pounds per 1000 square feet, should be sufficient for best results. **Do not over-fertilize.** The fertilizer should be watered in. In periods of extended drouth a thorough watering every 5 to 7 days should be sufficient, wetting the soil 4 to 6 inches deep each time. It is quick to establish and easy to mow to a height of 1½ inches. However, it turns brown during the winter, does not tolerate shade and salt spray as well as St. Augustine, and does not grow well in alkaline soils or in dense shade.

Centipede sprigs or root runners 3 to 5 inches long should be planted in rows 10 to 12 inches apart, 4 to 5 inches apart in the rows. Insert sprigs in the ground, leaving about an inch showing. Give frequent watering until plants are rooted firmly.

##### Centipede Grass Seed

1 oz. \$1.15; ¼ lb. \$4.25; ½ lb. \$8.00; 1 lb. \$15.00

*Postpaid*

##### Centipede Grass Sprigs or Roots

1 bu. \$2.25; 5 bu. \$2.15 per bu.; 10 bu. \$2.00 per bu.

*Not prepaid. Available April to November. Shipped only by express direct from a Florida nursery.*

*Write Mail Order Department, Kilgore Seed Co., Plant City, Fla., for instructive folder on "Centipede Grass Lawns."*



# PASTURE GRASSES and CLOVERS

All prices quoted are f.o.b. Plant City except 1 lb. and 2 lb. lots which are postpaid.  
Write for prices on larger quantities.  
All prices quoted are subject to change without notice.

To get better growth and increase soil nitrogen, it is essential to inoculate seed of all leguminous crops—clovers, Southern peas (cowpeas), crotalaria, hairy Indigo, lupines, peanuts, sesbania, etc. (See page 69.) Bermuda, Carpet and Italian Rye Grass are used for pastures as well as lawns. (See pages 39 and 40).

## BAHIA GRASSES FOR PASTURES

Sow 10 to 15 pounds broadcast per acre for pasture.

Bahia grass is established with seed planted  $\frac{1}{4}$  to  $\frac{1}{2}$  inch in depth, during any month of the year in Florida. Seeds often germinate slowly being covered by tough, waxy coat, which does not allow water to enter for germination. Exposure to sun for a few days before planting hastens germination. Widely adapted. Will grow in dry soils because of deep root system. Makes one of Florida's best permanent pasture grasses.

**Common Bahia.** A low growing perennial, spreading by short, stout, woody runners which root heavily with large fibrous roots, forming tough sod, even on drouthy, sandy soils. Leaf blades shorter and coarser than Pensacola but usually less than  $\frac{1}{2}$  inch wide.

1 lb. 60c; 10 lbs. \$2.30; 25 lbs. \$5.25; 100 lbs. \$20.00

**Pensacola Bahia.** Narrow leaf, tall, erect-growing strain of Bahia grass, very hardy, claimed to stand cold weather conditions better than Common. Pensacola variety seed usually germinates quicker and stronger than Common. Succeeds on wide range of soils. Has heavy root system and withstands drouth better than other pasture grasses. For best results fertilize with 500 lbs. per acre of a complete fertilizer. Prevents erosion and provides very palatable pasturage. Withstands heavy grazing and is very resistant to trampling by livestock.

1 lb. 75c; 10 lbs. \$4.30; 25 lbs. \$10.25; 100 lbs. \$40.00

**Argentine Bahia.** Sow 2 to 5 pounds per acre. Has attracted widespread attention as cattle pound-producer grass, all over Florida. Makes a semi-erect grass generally more vigorous than other Bahias, with leaves wider than Pensacola but slightly narrower than Common. Adapted to all kinds of soils, from high dry sand to low wet areas. Just as frost-resistant as Pensacola. For best results sow seed in late fall or early winter one-half to one inch deep. In numerous tests Argentine Bahia exceeded other varieties of Bahia in forage produced per acre. As nutritious and claimed to be more palatable to cattle than other Bahias. Ideal forage grasses for late summer grazing.

1 lb. 80c; 10 lbs. \$5.00; 25 lbs. \$12.00; 100 lbs. \$47.00

## BUFFEL GRASS

Sow 1 to 2 pounds of seed per acre

This perennial bunch grass widely adapted to soils and climates. Stands hot dry weather well. Due to massive root system extending as much as 8 feet deep, it is unequalled for erosion control and soil building. Provides excellent, quick growing hay and grazing. Palatable to cattle and possesses high protein content. Grows 3 to 4 feet high, even 6 feet under ideal conditions. Sow preferably in early spring through middle of summer, 1 to 2 pounds of seed per acre in rows 36 to 42 inches apart,  $\frac{1}{4}$  to  $\frac{1}{2}$  inch deep, packing soil firmly over seed. Fertilize with a complete fertilizer high in nitrogen.

1 lb. \$1.75; 10 lbs. \$17.25; 25 lbs. \$42.75; 100 lbs. \$170.00

## SWEET SUDAN GRASS (Certified)

Sow 10 pounds in drills or 20 pounds broadcast per acre.

Resembles common Sudan in plant habit and yield, but is sweet and juicy, more productive and resistant to several foliage diseases which attack common Sudan. Grazing tests have indicated that cattle prefer Sweet Sudan to common Sudan. Sow from March to Sept.

1 lb. 50c; 10 lbs. \$1.70; 25 lbs. \$3.75; 100 lbs. \$14.00

## BEGGARWEED (Florida Clover)

Sow 10 pounds per acre.

Perfectly adapted to Florida sandy soil. Makes fine pasture, nutritious hay. For best quality hay, cut crop when 3 to 4 feet high, or at beginning of blooming

period. Sow from April to August. Cover seed lightly, as it is small, and if put in ground too deep it cannot germinate. Inoculate seed with Nitragin. (See page 69.)

1 lb. \$1.15; 10 lbs. \$7.80; 25 lbs. \$19.00; 100 lbs. \$75.00

## BENNE (SESAME)

Sow 3 to 5 pounds per acre drilled.

Produces flowers, followed by seed pods, which scatter oily seeds in profusion, attracting quail and other game birds. Valuable for poultry feed. Plant from March to June, drilling seed thinly in rows 3 to 4 feet apart. We have only the native giant or tall variety growing to a height of from 4 to 6 feet, and rather spreading.

$\frac{1}{4}$  lb. 25c; 1 lb. 80c; 10 lbs. \$4.30; 25 lbs. \$10.25;  
100 lbs. \$40.00

## ALFALFA-HAIRY PERUVIAN

Sow 20 pounds per acre.

A valuable forage and hay crop for Florida. Does well in central and south Florida where it is used extensively by dairy and cattle farmers. Does best on moist, dark sandy soils (not subject to flooding). Also does well on muck soil. Should be tested before starting seedbed and apply sufficient limestone several months in advance of seeding worked deep into soil to correct pH. The pH should be brought up to 6.5 or 6.8. Extra lime can be applied after the seedbed has been prepared if pH is too low. After plowing, firm seedbed by disking and cultivating. Broadcast 1000 pounds per acre of fertilizer high in phosphoric acid and potash plus 20 pounds of borax ahead of seeding. Inoculate seed with double recommended rate of inoculant. Use cultipacker type of seeder, and seed from September to January at rate of 20 pounds per acre, when soil is moist. After seeding apply 500 pounds of a phosphoric acid-potash fertilizer broadcast in 60 to 90 days.

1 lb. 75c; 10 lbs. \$4.50; 25 lbs. \$10.75; 100 lbs. \$42.00

## ALYCE CLOVER

Sow 10 to 15 pounds per acre.

Used as cover crop, soil builder, for hay, and as pasture plant. Best time for planting is during spring. Must be planted on well drained soil free of contamination with root knot. Seed should be planted on well prepared ground and covered very lightly. Pack soil after seeding. Makes palatable hay. Cut for hay when it begins to flower. A good soil builder. Most economical fertilizer application is 200 to 400 pounds per acre of 0-16-16 or 0-14-14 mixture.

1 lb. 50c; 10 lbs. \$1.90; 25 lbs. \$4.25; 100 lbs. \$16.00

## ANNUAL SWEET CLOVER

Sow 8 to 12 pounds per acre of inoculated seed alone,

or 3 to 5 pounds per acre in mixed clover plantings.

A most valuable annual legume crop for grazing and soil improvement. May be sown in Florida from Sept. through Jan., although from Oct. 15 to Nov. 15 sowing is best. Makes a rapid growth, and widely adapted, growing best on moist soils not subject to flooding. Adapted to higher, drier ground than Louisiana White Dutch, requires less moisture. Does not tolerate very close grazing. Grazing reaches peak during March and April. Liming soil heavily and applying 400 to 600 pounds of a fertilizer high in Phosphoric Acid and Potash at planting time is recommended.

**HUBAM (Sweet).** The best and most promising commercial variety for Florida.

1 lb. 60c; 10 lbs. \$2.50; 25 lbs. \$5.75; 100 lbs. \$22.00

**FLORANNA (Sweet).** This new variety developed by the Florida Agricultural Experiment Station from hardy strains found volunteering is best adapted to Florida soil and climatic conditions. It outyields and comes through drouth and cold weather better than other varieties.

1 lb. 80c; 10 lbs. \$4.80; 25 lbs. \$11.50; 100 lbs. \$45.00



# CLOVERS and SOUTHERN PEAS (Cowpeas)

All prices quoted are subject to change without notice.  
Prices quoted are f.o.b. Plant City except 2 lb. lots or less which are postpaid.  
Write for prices on larger quantities, indicating variety and quantity desired.

*Clovers (continued from page 41)*

## WHITE DUTCH CLOVER (Louisiana grown)

Sow 4 to 6 pounds per acre alone, or 2 to 3 pounds per acre in a mixture.

Recommended principally for mixture in perennial pastures. Furnishes abundant grazing throughout the year in the South. Also useful for lawns, especially in mixtures. Does well with Bermuda Grass. Succeeds best on moist ground and during wet seasons. Sow in the late fall and early winter. Fertilize with 400 to 500 lbs. per acre of 0-14-10 mixture.

1 lb. \$1.25; 10 lbs. \$8.80; 25 lbs. \$21.50; 100 lbs. \$85.00

## LADINO CLOVER (Certified)

Sow 4 to 6 pounds per acre.

This is a large form of White Dutch Clover. It grows rapidly and spreads by creeping stems that root at the nodes. Does not flower so profusely as White Dutch. Usually a perennial, but may die if too dry or under long duration of very high temperatures. Sow in late fall or early winter. Used for soil improvement, pasture and hay. Does best on fertile, moisture retaining soils.

1 lb. \$1.15; 10 lbs. \$7.80; 25 lbs. \$19.00; 100 lbs. \$75.00

## CERTIFIED KENLAND RED CLOVER

Sow 4 to 6 pounds per acre.

This clover is rapidly gaining in popularity in Florida as a luxuriant hay and pasture crop for cattle and hogs. It has a longer active growing season than some other legumes. Fertilize with 400 to 600 lbs. per acre of a 0-14-10 or 0-10-10 fertilizer mixture just before planting time. Seed from September to December. Grows well with Bermuda and Bahia grass. For hay cut when in full bloom. It makes good growth in April, May and June when White Dutch and Crimson clover don't do so well.

1 lb. \$1.00; 10 lbs. \$6.30; 25 lbs. \$15.25; 100 lbs. \$60.00

## CULTURE OF SOUTHERN PEAS

Peas may be sown any time from March to September. Field varieties are always sown broadcast. Table varieties are always sown in drills three to four feet apart. The amount of seed per acre depends upon the size of seed, which varies with varieties. Fertilize up to 300 lbs. per acre with 3-8-5 mixture. Inoculate seed with Nitragin. (See page 69.) It is important to treat pea seed with Spergon at the rate of 2 oz. per bushel when planted before the ground gets warm in spring. (See pages 52 and 62.) For insect control see page 54.

Sow 20 to 45 pounds per acre in drills with table varieties.

In the garden  $\frac{1}{2}$  to  $\frac{3}{4}$  pounds to 150 foot row.

Sow 1 to  $1\frac{1}{2}$  bushels per acre broadcast with field varieties.

A bushel weighs 60 pounds.



*Thin Shelled Purple Hull (Dixie Queen)*

## TABLE VARIETIES OF SOUTHERN PEAS

**IMPROVED BUSH CONCH.** (62 days.) This variety of Conch pea is very early. A bunch type, the stocky vigorously upright growing plants making a bush about 24 inches tall. Some short runners or tendrils may develop as with most table peas. The amount of runner growth is dependent largely upon the moisture available. Pods are oval, medium green, straight, and closely filled with light green, tender peas. Seeds are small, smooth, ivory white.

$\frac{1}{4}$  lb. 20c; 1 lb. 60c; 2 lbs. \$1.00  
15 lbs. (pk.) \$4.65; 60 lbs. (bu.) \$18.00

**CALIFORNIA BLACKEYE NO. 5 (Early Wilt-Resistant Ramshorn Blackeyes).** (65 days.) This strain most closely resembles the old California Blackeye in its small, erect, early maturing vines and quality of seed. It is resistant to nematodes (root knot) and cowpea wilt. Very early and prolific, making a great number of 10 to 12 inch long pods on an upright vine. Uniform in size, growth and maturity, and flavor. Our seed of this variety is California grown and is at least 98 per cent free of noxious weed seeds and mixture, and of high germination.

$\frac{1}{4}$  lb. 20c; 1 lb. 50c; 2 lbs. 85c  
15 lbs. (pk.) \$2.90; 60 lbs. (bu.) \$11.00

**GIANT WILT-RESISTANT RAMSHORN BLACK-EYES.** (70 days.) A vigorous heavy yielding pea. About 10 days later than the Early Wilt-Resistant Ramshorn Blackeye. The peas are much larger than the regular Blackeyes and can be used both green and dried.

$\frac{1}{4}$  lb. 20c; 1 lb. 50c; 2 lbs. 85c

*Write for prices on larger quantities.*

**ALABAMA CROWDER.** (70 days.) The very vigorous bush is of the semi-runner type. The large, straight pods average about 8 to 9 inches in length and are of very attractive appearance, which has created a demand on the market for this variety, selling for higher prices than other Crowder peas. A sure cropper, heavy yielder, and well adapted to Florida growing conditions.

$\frac{1}{4}$  lb. 20c; 1 lb. 60c; 2 lbs. \$1.00  
15 lbs. (pk.) \$4.00; 60 lbs. (bu.) \$15.00

**THIN SHELLLED PURPLE HULL (Dixie Queen).** (75 days.) Vines make vigorous branching growth, and do best when planted in three-foot rows. Very productive. Pods ripen to a deep purple color, are very large, long, slightly curved, well filled with large seeds of high edible quality.

$\frac{1}{4}$  lb. 20c; 1 lb. 60c; 2 lbs. \$1.00  
15 lbs. (pk.) \$4.15; 60 lbs. (bu.) \$16.00

## FIELD VARIETIES OF COWPEAS

*Due to the unstable condition of the market, we are unable to make price quotations on field varieties of cowpeas.*

*Write for prices*

**BRABHAM.** The Brabham Cowpea is resistant to wilt and root knot which attack some other varieties, makes fine, vigorous growth of vines, is early, and, when planted in March, April, or May good cuttings of hay can be made. Also a very good soil builder.

**MIXED.** The advantage claimed for sowing Mixed Cowpeas is that they produce an exceedingly heavy growth of vines for forage or soil improvement.



# FIELD SEED

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.

For information on "When to Plant Field Seed in Florida" see page 5.

## CHUFAS

Plant 1 to 1½ pecks (10 to 15 lbs.) per acre.

(90 days.) This is a species of ground nut, easily grown, a useful hog fattening crop. Plant from April to August. Cultivation same as for bunch peanuts. Can be left in the ground until time to turn in hogs who do harvesting. On sandy loam soil best adapted for Chufas, plant in 2-ft. rows with plants 12 inches in the row, or in 3-ft. rows with plants 6 inches in the row.

1 lb. 50c; 2 lbs. 90c; 10 lbs. (pk) \$2.25  
40 lbs. (bu.) \$8.00

## CHICKEN CORN. (Shallu or Egyptian Wheat)

Sow 8 to 10 pounds per acre for grain.  
25 to 30 pounds for cover crop.

This crop is being grown extensively and very successfully as a summer green manure or cover crop in many sections of Florida, especially in muck soils. Grows rapidly and makes a great amount of vegetable matter to plow under in a relatively short time. Not a legume. If allowed to go to seed for a grain crop like other sorghums, makes large, loose, bushy heads covered thickly with small grains, which if left standing, shatter, and chickens gather them. Grain is of high feeding value and can be fed to chickens, hogs and cattle as desired. For grain sow seed thinly in 3 to 4 foot rows, leaving 3 to 4 plants every 3 feet of row. Plant from March to August.

1 lb. 50c; 10 lbs. \$1.55; 25 lbs. \$3.40; 100 lbs. \$12.50

## CROTALARIA

Sow 10 to 20 pounds of seed per acre.

Crotalaria is particularly desirable as a cover crop for groves, truck and farm lands, does well on all soils except muck. Is a legume, thus adding a large amount of nitrogen to soil. Is especially valuable because it is immune to nematodes causing root knot disease. Makes profuse growth, furnishing abundance of green material for humus.

Sow broadcast March to June, on well prepared land.

**Crotalaria—Early Spectabilis.** This vigorous growing, long leaved variety is not so fibrous as others, the growth is much heavier, and the stems are hollow. As a result, it rots much quicker when plowed under. Spectabilis does well on high ground, but is best adapted to the better grades of sandy soils. Leaves, stems and seed of this Crotalaria plant, green or dry, are poisonous to poultry and livestock.

Write for prices.



Crotalaria—Early Spectabilis

Valuable for adding organic matter and nitrogen to the soil

**HEGARI (Higear)—See Sorghum, page 46**

## EARLY HAIRY INDIGO (Indigofera hirsuta)

Sow three to five pounds per acre in 30 inch drills,  
6 to 10 pounds broadcast.

A three purpose legume, used as a summer cover crop, for hay and for grazing. Used extensively as a cover crop by citrus growers in south Florida, as a grazing crop or for hay by cattlemen and as a cover crop by general farmers. Its feeding value compares fairly well with alfalfa, making a palatable and nutritious hay. It makes an ideal summer cover crop for citrus groves, harboring no troublesome insects, and reseeds itself year after year. It has a deep tap root, making it very resistant to drouth. The plants grow four to seven feet in height, and produce an abundance of hay and grazing. If grazed or harvested for hay before the stems become woody a second or third growth may be expected.

Seed may be planted not over an inch deep, from the middle of March to the latter part of May, but early seedings are preferable. Heavier seedings are desired for green manure or forage, lighter seeding for seed. On poor soil 300 to 500 lbs. per acre before seeding of a 0-10-10 or 0-14-10 fertilizer mixture is desirable. On extremely acid soils apply 1000 lbs. of lime per acre. It will grow on a wide range of soils, but well drained sandy loam soils are best. It is sensitive to cold, is killed by the first hard frost and is best suited to the section of Florida from Gainesville south, although the early strain can be grown successfully in north Florida and as far north as central Georgia. Highly resistant to root knot, but in general it is reasonably free from disease and insect injury, and is resistant to pumpkin bugs.

Write for prices.

## BLANKET INDIGO

Sow 3 to 5 pounds of seed per acre.

This is a summer annual legume, different from Hairy Indigo in growth—ordinarily 18 to 24 inches. Excellent for sowing with other permanent pasture grasses. Also recommended for sowing in groves. Plant from early spring to the end of June. Offers great possibilities as a pasture or hay crop in Florida—especially for hay in late summer—September or early October. Prefers soils of low fertility, especially low in nitrogen. A good soil builder. Will not stand excessive moisture. It is highly palatable and will stand close grazing.

Write for prices.

## LUPINES

50 to 60 pounds with yellow and 50 to 90 pounds with blue per acre broadcast, or preferably in drills with a grain drill.

Lupines are adapted to north and central Florida as a winter legume cover crop. They are not affected by cold and because of their heavy yield make an ideal green manure crop. Plant from late September to December, preferably in late September or October, in well prepared soil, using between 250 and 300 lbs. of Superphosphate per acre.

**Bitter Blue Lupine.** This variety has rapidly grown in popularity as a soil builder. Normal growth is about waist high and normal weight of green matter 20-30 tons per acre. It produces about three times as much growth as Winter Peas, Hairy Vetch, etc. It protects the soil from erosion and restores humus and plant food removed by peanuts and other crops. Peanut and corn yields following Bitter Blue Lupine have been greatly increased. A good winter cover or green manure crop, but poisonous to livestock.

1 lb. 40c; 10 lbs. 95c; 25 lbs. \$1.85; 100 lbs. \$6.50

(Continued on next page)



# FIELD SEED (Continued)

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.

## LUPINES (Continued)

**SWEET BLUE LUPINE.** This variety is the pasture and feed counterpart of Bitter Blue Lupine. Both plant and leaf are sweet to the taste and are relished by all kinds of livestock. Makes about the same growth and tonnage as Bitter Blue Lupine. Stock can be turned in on Sweet Blue Lupine as soon as it is about a foot high. The plants have a high protein value averaging approximately 18% of the dry matter.

1 lb. 50c; 10 lbs. \$1.70; 25 lbs. \$3.75; 100 lbs. \$15.00

**SWEET YELLOW LUPINE. (White seeded.)** Seems to be more palatable and therefore is more readily grazed by cattle than the Sweet Blue strains, and will recover more rapidly than the Blue after grazing. The sweet strains of Yellow Lupine tend to be more tender and less resistant to excessive cold than the Blue, and thus the Yellow Lupine is more particularly adapted to the southern portion of the Lupine belt than to the more northern areas.

1 lb. 60c; 10 lbs. \$2.30; 25 lbs. \$5.25; 100 lbs. \$20.00

## PEARL or CATTAIL MILLET

Sow 8 to 10 pounds per acre in rows 3 feet apart.

Sow 20 to 25 pounds per acre broadcast for pasturage.

This is one of the best yielders of green forage and hay, and continues to grow and produce through the entire season if cut frequently enough for hay to prevent its going to seed. Relished by all kinds of stock, and they eat it greedily. Plant in early spring as soon as frost danger is past in March, and can be planted from then on until September 1. Sow thinly in rows three feet apart or for pasturage, sow broadcast 20 to 25 lbs. per acre.

1 lb. 50c; 10 lbs. \$1.80; 25 lbs. \$4.00; 100 lbs. \$15.00



Southland Oats

## OATS

Sow 5 pecks (40 lbs.) per acre in drills; broadcast 2 to 2½ bu. per acre. A bushel weighs 32 pounds.

Plant in drills 1½ feet apart and cover seed 2 to 3 inches deep. If the seed is broadcast, use tooth harrow. The best time to plant oats in Florida is October and November, but can be planted to January. Land should receive 300 to 400 lbs. per acre of complete fertilizer at planting, followed by a side-dressing of 150 to 200 lbs. Nitrate of Soda or its equivalent after the crop is half grown.

**SOUTHLAND (Disease Resistant).** (125 days.) Is a vigorous growing oat with broad, coarse leaves and a strong upright growth habit. Produces a large amount of very early forage. A spring type oat and is not cold-hardy, and where not injured by cold it produces an abundance of green pasture. Adapted to the Gulf Coast region and to north central Florida. Resistant to Victoria blight and to most races of Crown rust, it is susceptible to two races of this rust which have now become prevalent in Florida. It suffers little from stem rust, but is quite susceptible to Culm rot. Yields of forage and grain have been high under good cultural conditions. Grain is heavy and plump, free of hairs, white in color and of high quality. A good general purpose oat for Florida.

1 lb. 35c; 2 lbs. 50c; 8 lbs. (pk.) 60c; 32 lbs. (bu.) \$1.75

**FLORILAND (new) (Disease resistant).** (115 days.) The outstanding characteristic of this new variety, selected and increased at the north Florida Experiment Station, is its very high resistance to Crown rust, the most important oat disease in Florida. An early intermediate to upright habit of growth which gives good yields of forage and grain. More resistant to Culm rot than Southland, and it is also resistant to Victoria blight. Has better winter hardiness than Southland and is well adapted to north and west Florida. Kernels are red and quite hairy. The grain quality is only fair, recommended chiefly for pasture.

Lb. 35c; 2 lbs. 50c; 8 lbs. (pk.) 70c; 32 lbs. (bu.) \$2.00

**SEMINOLE (new) (Disease resistant).** (110 days.) This new, early maturing oat has been increased on Experiment Station Farms in Florida, and seed is now available. It is resistant to different races of Crown rust and to smut. Due to its earliness it escapes damage from stem rust. A short variety with strong straw and not subject to lodging. The grayish-white grain is larger, plumper and of higher quality than Floriland. Makes a very early, upright growth of medium large leaves. Produces an abundance of early forage and is good for early and mid-winter forage. Also excellent in grain yield. Because of its earliness it is well adapted to central and south Florida.

Lb. 40c; 2 lbs. 65c; 8 lbs. (pk.) 80c; 32 lbs. (bu.) \$2.50

## PEANUTS

Plant 1 bushel of seed per acre.

A bushel weighs 25 lbs., except Valencia 22 lbs. per bu.

In addition to the crop of nuts for hogs, peanuts yield a large quantity of nutritious dry forage. Dixie Runner peanuts are by far the best when grown to fatten hogs, as they can be left in the ground until eaten.

It is better to shell the seed that is to be planted. If planted in corn, a bushel will plant 4 acres, but where planted solid it requires a bushel of seed per acre, in 3-foot rows, dropping seeds in hills a foot apart. Plant from March to July. For best results inoculate peanuts with Nitragin. (See page 69.)

Most shelled peanuts are treated so they will keep better and longer, and also to protect them from rot and disease after they are planted and to increase the stand. Do not use Nitragin inoculation with treated shelled peanuts.

For Insect and Disease Control See Page 57.

**VALENCIA.** (100 days.) This variety is adapted for growing in light sandy soils. It fills the demand for a peanut that is large and contains more nuts than the common varieties. Pods are large, well filled, usually containing 3 or 4 nuts.

Lb. 55c; 5½ lbs. (pk.) \$1.75; 22 lbs. (bu.) \$6.50


(Continued on next page)



# FIELD SEED

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.

## PEANUTS (Continued)



*Dixie Runner*

**DIXIE RUNNER.** (120 days.) This variety produces high yields like Spanish peanuts, in four years' tests averaging 53 per cent higher yields than the Florida Runner, and yet remains dormant after maturity like the regular Runners, thus being especially suited to "hogging off." It is a vigorous grower with more prostrate branches, and therefore is more apt to fill the spaces left by skips in the stand. Kernels are much like Florida Runner in size, shape, markings and color. Pods are somewhat lighter and brighter in color, shorter, thicker at the waist and more cylindrical in shape. In addition to high yield and dormancy characters, this variety possesses a valuable quality for shellers, as it shows less than 1 per cent of concealed damage, or decay of the interior of the kernels, which is a serious disadvantage of some varieties going to the shelling plants, and may average as high as 20 per cent concealed damage.

Lb. 45c; 6¼ lbs. (pk.) \$1.40; 25 lbs. (bu.) \$5.00

**Shelled and treated Dixie Runners.** One bushel of good Dixie Runners should yield about 16 lbs. shelled peanuts. Plant 25 lbs. per acre.

Lb. 65c; 10 lbs. \$3.30; 25 lbs. \$7.75; 100 lbs. \$30.00

## DWARF ESSEX RAPE

Sow 4 pounds in drills or 8 to 10 pounds broadcast per acre.

For early spring, fall or winter planting to give quick green forage or grazing for hogs and poultry. While mostly planted in fall and early winter, it may be planted in early spring, but it will not stand much warm weather. It makes a quick succulent growth. It can be planted in rows like turnips, or broadcast covering by a light harrowing. It is ready for grazing in eight to ten weeks after planting. After cutting, or grazing, it keeps coming out again. Rape can be sown with fine results any month from September to March. It is a splendid green feed and forage crop relished by all kinds of stock and especially by milk cows.

Pkt. 10c; 1 lb. 50c; 10 lbs. \$2.30; 25 lbs. \$5.25  
100 lbs. \$20.00

## RYE

Sow 3 pecks in drills; broadcast 1 to 1¼ bushels per acre. A bushel weighs 56 pounds.

Rye can be planted from September to January but October, November and December are the best months in which to plant for seed. Earlier planting is recommended for winter and early spring grazing. Fertilization is the same as for oats. (See page 44.)

**FLORIDA BLACK.** A standard, upright, tall growing variety. Northern and western varieties of rye have a tendency to fall over on the ground and are not suitable for grazing or cutting in Florida. Florida Black is upright in growth and is exceptionally well suited for early grazing, making pasturage quickly, and it withstands cold very well. Highly recommended by the Florida Agricultural Experiment Station.

1 lb. 50c; 2 lbs. 85c; 14 lbs. (pk.) \$1.40  
56 lbs. (bu.) \$5.00

**ABRUZZI (certified).** One of the best for Florida, producing a heavier tonnage of green feed and grain than most other varieties. Tall and upright, and a heavy yielder.

1 lb. 45c; 2 lbs. 80c; 14 lbs. (pk.) \$1.25  
56 lbs. (bu.) \$4.50



*Abruzzi Rye*

## RICE

Sow ½ bushel (22 lbs.) per acre.

Rice makes fine feed for both stock and poultry. Prepare the ground thoroughly, drill ½ bushel per acre in 2½-foot rows and cultivate a few times. Should be planted in March or April so that it may mature at a time to miss the rice-bird season. If planted in June and July it will also miss the bird season.

**GOPHER.** A popular variety for Florida, generally known as Upland Rice, but will also do well on low land. Grains heavy, broad, and flat.

Write for prices.

**PEARL (White).** Grown extensively in Florida. It produces a heavier, taller, more vigorous stalk with heavier heads than other varieties of this type. Does not shatter so badly as other varieties. Produces long, slender grains. Best adapted for low wet land.

Write for prices.



# FIELD SEED (Cont'd)

All prices quoted are subject to change without notice and are f.o.b. Plant City except 2 lb. lots or less which are postpaid. Write for prices on larger quantities.



Sorghum—Texas Seeded Ribbon Cane

## SORGHUM

Amount of seed needed per acre depends upon the purpose for which this crops is grown as indicated below.

Sow 8-10 lbs. to 1 bushel (50 lbs.) per acre.

The uses of Sorghum are many; green feed, hay, pasturage, silage, syrup, etc.

It is common practice to sow Sorghum and Cowpeas together for hay and pasturage purposes using about ½ bushel Sorghum and 1 bushel of Cowpeas per acre, sown broadcast. Planted in drills alone in 3- to 3½-foot rows use 1 peck or 12 to 15 lbs. per acre; where planted for making syrup, in 4-foot rows, use 8 to 10 lbs. per acre, thinning plants 8 to 10 inches apart; or where sown broadcast for forage, use 1 bushel of seed per acre. For grain Sorghums apply 300 to 500 lbs. of a complete fertilizer mixture per acre. Plant from March to August.

**TEXAS SEEDED RIBBON CANE.** (For syrup, ensilage, forage and green feed.) The plants grow ten to twelve feet tall, and the thick stalks contain an abundance of juice and also produce plenty of leaves for fodder. Not only valuable for syrup-making, but will make a large amount of forage and green feed for livestock. The tallest and heaviest producing Sorghum for both forage and syrup.

1 lb. 60c; 10 lbs. \$2.30; 25 lbs. \$5.25; 100 lbs. \$20.00

**HEGARI (Higear). Early Dwarf Type.** (For grain and forage.) (100 days.) This crop, which is quite similar to Kaffir corn, has become very popular in the south-east because of its quick growth and heavy yield of grain, sometimes running over a ton to the acre. Unlike Kaffir, the stalks are very sweet and juicy. Grows 2 to 5 feet depending on type of soil, with large heads of white grains which do not shatter. Stands dry weather exceptionally well. Sow ten pounds per acre for grain, one bushel (50 lbs.) per acre broadcast for forage.

1 lb. 40c; 10 lbs. \$1.10; 25 lbs. \$2.25; 100 lbs. \$8.00

**HEGARI (Higear). Regular Type.** (For forage and cover crop.) (115 days.) This strain is slower to seed but makes more forage quicker than the Early Dwarf Type, and also is better for cover crop. Sow 30 to 40 lbs. per acre broadcast for forage or for a cover crop. On the lower east coast this strain is used extensively as a cover crop and a common practice is to cut the stand just before seed heads form; the plants will then sprout out and make similar height as before, thus giving a double cover crop to turn under.

1 lb. 40c; 10 lbs. \$1.10; 25 lbs. \$2.25; 100 lbs. \$8.00

**SHALLU OR EGYPTIAN WHEAT.** See "Chicken Corn"—page 43.

## SESBANIA

Sow 30 to 40 pounds per acre.

Sesbania is a most valuable summer cover crop to grow on land that is more or less covered with water. Unlike other cover crops, Sesbania will grow in water, and will grow ideally well in the hot weather of mid-summer on low, wet land. It is used extensively as a summer cover crop in low, wet muck lands, as well as in other vegetable and commercial flower-growing lands. It is valuable not only as a green manure cover crop, but also as a means of killing out Bermuda grass, nut grass and other weeds, because it grows so rapidly. In addition, being a leguminous crop, forming nodules on the roots, it fixes atmospheric nitrogen in the soil for seed inoculant. (See page 69.)

We recommend scarified Sesbania seed for quicker and higher germination. Seed is broadcast at the rate of thirty to forty pounds per acre, even a little heavier if weeds are bad. Usually in late May, June and July, dense, tall growth is obtained in about two months, at which time, just before blooming, the crop may be plowed under and the stalks will rot or decay rapidly. In some sections Sesbania is grown for pole bean sticks. For this purpose it should be sown 12 to 15 lbs. per acre.

Write for prices.

## SUNFLOWER

Sow 6 pounds per acre.

**MAMMOTH RUSSIAN.** The Mammoth Russian makes three to four times as much seed as ordinary varieties. It is used extensively as a windbreak for beans. Plant from March to September. Plant seed in drills three feet apart, and hills one foot apart in the drill, thinning out to one stalk in the hill when four to six inches high.

1 lb. 60c; 10 lbs. \$2.80; 25 lbs. \$6.50; 100 lbs. \$25.00

## VELVET BEANS

Sow 2 pecks (30 lbs.) per acre.

Velvet Beans are good for groves and truck lands in building up the humus; good pasturage for cattle and hogs; and fine land renovators. They are specially valuable because they are resistant or practically immune to nematodes causing root knot disease. Plant from March to August. Use Nitragin. (Page 69.)

**90-DAY.** (90 to 100 days.) This variety makes a very luxuriant growth and an abundant amount of foliage for pasturage and soil improvement. It matures quicker than other varieties.

Write for prices on Velvet Beans, indicating quantity desired.



A typical field of Hegari (Higear) grown for grain



# STRAWBERRY PLANTS

All prices quoted are subject to change without notice.

We supply Maryland, Tennessee and Arkansas grown strawberry stock plants of the Missionary and Florida 90 variety from nurseries in those states. These northern grown stock plants are set out in multiplication beds in Florida from January through March. The young plants are usually set in the permanent field from September through November or early December, depending on weather conditions. The northern grown plants are usually available about November 1st and when Florida grown plants are not available are used by commercial growers for November setting and are also used to a great extent for home gardens. Care should be exercised never to allow the roots to dry out.

Strawberry plants may be set in rows three feet apart for single rows, or 4 to 5 feet apart for double rows, allowing 16 inches between double rows on each bed with 12 to 14 inches between plants in the row.

In setting strawberry plants, holes should be made deep enough to set the roots straight. Plants should be set so that the surface soil comes just to the crown.

Soil should be well firmed around the roots, and during dry weather each plant should be watered daily for several days after setting, preferably in the late afternoon.

If the land to be set to berries is very sour or acid, apply 500 lbs. lime or 1200 lbs. hardwood ashes per acre before setting plants. For control of insects and diseases see page 56.

After the plants are well established, 3 or 4 successive applications of high grade commercial fertilizer, such as 4-7-5 mixture, should be given the plants at about monthly intervals, being sure to make one heavy application of a 3-8-8 or 4-8-8 mixture when the plants begin

|                           | 100<br>plants | 250<br>plants | 500<br>plants | 1000<br>plants |
|---------------------------|---------------|---------------|---------------|----------------|
|                           | Postpaid      |               | f.o.b.        | Nursery        |
| <b>Missionary Variety</b> |               |               |               |                |
| Maryland plants           | \$2.00        | \$3.25        | \$4.25        | \$7.50         |
| Tennessee plants          | 2.00          | 3.50          | 5.00          | 8.00           |
| <b>Florida 90 Variety</b> |               |               |               |                |
| Maryland plants           | 2.25          | 3.75          | 5.25          | 9.00           |
| Tennessee plants          | 2.00          | 3.50          | 5.00          | 8.00           |

## ROSE BUSHES FOR FLORIDA

### Planting and Care of Rose Bushes.

Plant bushes 1½ to 2 feet apart each way. Dig holes sufficiently deep and wide to permit roots to be set in natural position, not doubled up. Put two to three pounds of pulverized sheep manure in each hole and mix well with a half pound of Vigoro or Vertagreen fertilizer. (See page 67). Then pour in a liberal supply of water, set bush, press the soil tightly around the roots, leaving no air spaces. Roses need lots of water, and should be fertilized every six to eight weeks with Vigoro or Vertagreen fertilizer, using about one-half pound per plant, and an occasional application of sheep manure will help. Spray with Nicotine Pyrox or dust thoroughly with Kilgore's Rose Dust (See page 63) at least once a week or as soon as diseases or insects appear.

All orders will be shipped direct from Texas nursery.

Delivery made in 12 to 15 days after receipt of order. Extra strong, No. 1 grade bushes—bloom approximately 2 months after setting out.

|  | Postpaid to any point in Florida |        |         |         |         |
|--|----------------------------------|--------|---------|---------|---------|
|  | 5                                | 10     | 25      | 50      | 100     |
|  | \$4.50                           | \$8.50 | \$20.00 | \$37.50 | \$70.00 |

No orders accepted for less than five bushes. However this does not apply to patented roses.

Plant any time after November 1 to the end of March.

### VARIETIES OF BUSH ROSES

- RED—  
Red Radiance, Poinsettia, Etoile De Hollande, Red Talisman.
- PINK—  
Pink Radiance, Editor McFarland, Briarcliff.
- WHITE—  
White American Beauty (Frau Karl Druschki), Caledonia.
- YELLOW—  
Golden Dawn Soeur Therese.
- TWO-TONE—  
Talisman—Orange and copper tones, suffused with pink.

### VARIETIES OF CLIMBING ROSES

- American Beauty (pink), Paul's Scarlet (scarlet).  
Mrs. Pierre S. DuPont (rich golden yellow).

### PATENTED AND NOVELTY ROSES

- CHARLOTTE ARMSTRONG—Plant Pat. 455. Beautiful cerise pink \$2.25  
MIRANDY—Plant Pat. 632. Beautiful bright red 2.25  
PEACE—Plant Pat. 591. Golden yellow etched with pink 2.50  
FORTY-NINER—Plant Pat. 792. Bi-color in contrasting red and yellow 2.50  
BRAVO—Plant Pat. 983. Red 2.25  
BLANCHE MALLERIN—Plant Pat. 494. White 2.00

### FLORIBUNDA ROSE

- FASHION—Plant Pat. 789. Peach 2.00

## The Kilgore Seed Company

General Offices and Mail Order Dept., Plant City, Fla.  
FOURTEEN KILGORE SEED STORES  
SERVING FLORIDA

A Florida Institution Maintained for Service to  
Florida Growers

With over 40 years experience specializing in vegetable, flower and field crop seeds and growers' supplies for Florida, and with fourteen Kilgore seed stores we are best qualified to serve Florida growers.

"Everything for the Florida Grower" is a Kilgore motto. Seeds, insecticides, fungicides, sprays, dusts, spraying and dusting equipment, fertilizers, cans and canning equipment as well as poultry and beekeepers' supplies are always carried in stock in Kilgore's Plant City warehouses and fourteen Florida stores.

High quality with prompt and efficient service at reasonable prices are foundation stones on which this pioneer Florida seed company has been built.



We now have a group of technically trained field men thoroughly familiar with local conditions, who are available to discuss your insect, disease and seed problems with you at any time.

## GLADIOLUS BULBS

### Rainbow Mixed


Enjoy beautiful cut flowers from your garden. For continuous blooms plant bulbs every 10 days to two weeks. In north and central Florida plant from August to October, and January to May. In South Florida plant from September to May. Set bulbs 4 inches deep and 6 inches apart in well-prepared and well-drained beds.

|   |                 |
|---|-----------------|
| Large, Exhibition size Gladiolus bulbs (2-2½ inches.) |                 |
| 1 dozen bulbs   | \$ .85 postpaid |
| 25 bulbs  | 1.50 postpaid   |
| 100 bulbs   | 4.00 postpaid   |
| 1000 bulbs  | 35.00 postpaid  |

Disease-free, well cured, graded extra large bulbs of good color assortment. No finer Glads for Florida can be secured.







Sunburst Mixture Zinnias

Grow flowers to make  
your home a more inviting  
and pleasant place  
to live.



Send for your free copy today.



Kilgore's Superb Mixture Petunias

## FLOWER SEEDS FOR FLORIDA

Recommended by

### The Kilgore Seed Company, Florida's Flower Garden Headquarters

Surround Your Florida Home with Beauty. Do It with Flowers.  
For larger quantities, we shall be glad to quote special prices.

Rate of sales tax—10c no tax; 11 to 35c - 1c tax; 36 to 65c - 2c tax; 66 to \$1.00 - 3c tax.

We recommend the following kinds and varieties of flowers for Florida as the result of many years of experimental work and testing under Florida conditions. For information on "When to sow flower seeds in Florida" see page 5.

**Ageratum** or **Floss Flower**, Purple Perfection, Dwarf Blue Cap

**African Daisy** (Arctotis Grandis)

**Alyssum**—Little Gem, Royal Carpet

**Aster**—Imperial mixture, Crego assorted colors

**Baby's Breath**—Pure White

**Bachelor's Button**—Sky Blue, Dainty mixture

**Balsam**—Double Camellia-flowered, Selected mixture

**Bean**—Scarlet Runner

**Blanket Flower** or **Gaillardia**—Double gaiety mixture

**Blue-eyed African Daisy** or **Arctotis**

**Blue Lace Flower** (Didiscus)—Heavenly Blue

**Calendula**—Campfire Improved, Lemon Queen, Excellent mixture, Art Shades mixture.

**California Poppy**—Gorgeous mixed. (Single flowering)

**Calliopsis** or **Annual Coreopsis**—Choice mixture

**Candytuft**—White Hyacinth-flowered, Border Group

**Carnation**—Charm mixture—Chaband's giant improved strain

**Celosia** (Cockscomb)—Tall Royal Velvet—Comb type. Pampas Plume—Plume type.

**Chinese Forget-Me-Not** or **Cynoglossom**—Sky Blue

**Cosmos**—Fiesta Semi-Double, Choice mixture, Single Orange Flare

**Dahlia**—Unwin's Bedding mixture

**Delphinium** (Perennial Larkspur)—Belladonna Hybrids mixed

**Floss Flower** or **Ageratum**—Dwarf Blue Cap, Purple Perfection

**Globe Amaranth** (Gomphrena)—Assorted colors

**Gourd** (Ornamental)—Small varieties mixed

**Hollyhock**—Indian Spring (Annual)—Semi-double, Rose and Pink shades

**Larkspur**—Regal finest mixed

**Linaria** (Baby Snapdragon) Northern Lights

**Lupine**—Assorted Colors, Dwarf Pixie Delight

**Marigold**—Gigantea Sunset Giants mixture, Yellow Supreme Dwarf French Double Harmony, Glitters (Chrysanthemum flowered type), Crown of Gold (Collarette Type), Dwarf Gigantea Pot o'Gold, Extra-Dwarf Double Mixed

**Morning Glory** (Improved Ipomea)—Scarlet O'Hara (Red), Pearly Gates (White), Heavenly Blue Improved (Blue)

**Mourning Bride** or **Scabiosa**—Sunrise mixture

**Nasturtium**—Semi-double Golden Gleam, Semi-double Scarlet Gleam, Semi-double Gleam Hybrids, Semi-double Dwarf Gem mixture

**Pansy**—Englemann Giant mixture

**Petunia**—Rosy Morn, Heavenly Blue, Howard Star Improved, Elk's Pride, Flaming Velvet, Snow Queen, Fire Chief (All-American Selection), Superb mixture, Fluffy Ruffles mixture

**Phlox**—Peninsular mixture, Dwarf Nan Compacta mixture

**Pinks** or **Dianthus**—Sweet Wivelsfield Double Mixed, Medal or Imperator mixture double flowered, Single Gaiety mixed

**Queen Anne's Lace**

**Rose Bushes**—different colors and varieties. (See page 47)

**Salvia**—Bonfire

**Shasta Daisy**—Single and Semi-Double Giants

**Snapdragon** or **Antirrhinum**—Giant mixture (rust resistant), Half-tall mixture (rust resistant), Giant Ruffled "Tetraploid" mixture

**Statice**—Single colors of Blue, Rose, Lavender, White, Yellow, also assorted colors

**Strawflower** or **Helichrysum**—Many colors

**Sunflower** or **Helianthus**—Sunshine mixture

**Sun Plant** or **Portulaca**—Double mixed

**Sweet Peas** (Early-flowering Spencer)—Mrs. Herbert Hoover, Mars, Shirley Temple, Hope, Lavanda, Ball's Orange Improved, Rainbow mixture, Zoolanek's multiflowered mixture

**Tithonia** or **Florida Sunflower**—Torch (All-America winner)

**Verbena**—Master mixture

**Vinca** or **Periwinkle**—Pink and White mixed

**Zinnia**—Blaze—All-America winner for 1954, Giant Hybrid mixture, Special Crown o' Gold mixture, Fantasy mixture, Giants of California special mixture. Pompon or Lilliput mixture, Crimson Monarch, Wonder Rose, Polar Bear, Sunburst mixture, Peppermint Stick, Persian Carpet

For further and complete information on growing flowers in Florida write the Florida Agricultural Experiment Station, mailing department, Gainesville, Fla., for a free copy of Florida Agricultural Extension Service Bulletin 133, entitled "Annual Flowers." Also write The Kilgore Seed Company, Plant City, Florida, for a free copy of "Kilgore's Flower Guide for Florida."



# THE KILGORE SEED COMPANY

A FLORIDA INSTITUTION MAINTAINED FOR SERVICE TO FLORIDA GROWERS  
FOURTEEN KILGORE SEED STORES AND MAIL ORDER DEPARTMENT SERVING FLORIDA



BELLE GLADE—304-12 S. Main St.



GENERAL OFFICES, WAREHOUSE  
AND MAIL ORDER DEPARTMENT  
214 S. Collins St.



FORT MYERS—1117 Anderson Ave.



MIAMI—600 N.W. Fifth St.



PALMETTO—710 - 13th St.



SANFORD—300-2 W. First St.



PAHOKEE—313-19 Belle Glade Rd.



WEST PALM BEACH—1310 Belvedere Rd.



WAUCHULA—702-6 W. Main St.



OCALA—909 N. Magnolia St.



GAINESVILLE—203 S.E. First Ave.



PLANT CITY—101 N. Evers St.



VERO BEACH—1109 20th St.



POMPANO BEACH—228 N. Flagler St.



HOMESTEAD—370 S. Flagler Ave.

Visit your nearest Kilgore Store. You will find many useful farm supplies there that are not listed in our catalog. "Everything for the Florida grower" is a Kilgore slogan.





# PESTICIDE SECTION

(Insecticides, Fungicides, Fumigants, Vermicides, Weed Killers,  
Spreaders, Stickers, Emulsifiers—Pages 50 to 66.)

The sources of information and recommendations in connection with insect pests and diseases given in this catalog are the State Experiment Stations, The Kilgore Seed Company Laboratory, our technically trained field men, and many members of our store personnel who have had years of close contact with agriculture.

Every attempt has been made to provide such in-

formation in a plain and simple form for the benefit of our growers. As, however, constant changes are taking place in the field of pesticidal problems because of new insect pests and diseases and the introduction of new pesticides, it is difficult to keep an annual catalog strictly up-to-date. We therefore invite you to contact our field and store men. They will be glad to assist you whenever you are facing new problems.

The use of pesticides is now just as necessary as that of fertilizers and various modern cultural practices. Many factors are involved in the use of pesticides in relation to results desired, such as the choice of proper pesticides for the control of different diseases and insects, and the time and method of application. One cannot expect satisfactory control with poor or wrongly timed applications. Too often the grower waits until the disease or insect has too much of a start before spraying or dusting. This usually is more expensive than starting off with a regular spray or dust program and keeping it up to prevent diseases and insects. He not only has to use excessive amounts of sprays or dusts to get control, but he also suffers considerable crop damage. As a rule, it is much cheaper to use a combination spray or dust when one must contend with

both insects and diseases, as one application may do both jobs. Most insecticides and fungicides are compatible and can be used together in sprays or dusts. Although we manufacture numerous combination dusts, only the more important are listed in this catalog. If you do not find the combination listed which you want, write to The Kilgore Seed Company, Plant City, Florida, or call at any of our stores.

With the aid of newly installed mixing and blending equipment in our Plant City factory, you can be assured that you will receive fresh and perfectly blended dusts formulated to meet your particular requirements. Only high-grade technical materials are used by us in producing the various pesticidal compositions that you may desire.

**SANITATION**—Serious losses may result from neglect of a few simple rules. Diseases will spread readily from sick to healthy plants by various means, depending upon the nature of the disease. Spores of many species of fungi are air- or wind-borne, others may be scattered over fields in rain or irrigation water. Insects, particularly those of the sucking type, such as aphids and thrips disseminate diseases. Certain cultural operations, such as pruning, picking, transplanting, etc., may scatter diseases, especially those caused by bacteria and viruses (mosaic being a specific example of the latter).

Considering methods of infection, it becomes obvious that the source of the disease should be destroyed. Affected plants, as well as decayed fruit, should be removed from the field and burned. Cultivation should be clean and frequent. Mechanical injury to plants by plowing, spraying or picking operations should be avoided as much as possible. Crop rotation is essential, provided that no other crop affected by the responsible diseases is grown for several years. This method often proves cheaper than an attempt to control the disease chemically. As weeds serve as host plants to many insects, which may be carriers of fungi, viruses and other undesirable organisms, the necessity for their eradication certainly becomes evident. Whenever practical, soil moisture should be kept fairly constant. Corrective measures should be adopted where soils are either too acid or too alkaline.

**MOSAIC**—Mosaic disease is caused by virus organisms in the sap of the affected plants. The viruses may be transmitted in various ways from diseased to healthy plants. Insects undoubtedly play an important part in the transmission of the disease, although it may be spread by other means, such as pruning operations, bruising of plants by workers or mechanical equipment, picking fruit, the use of tobacco and by other means.

The symptoms of mosaic vary slightly in different plants, but in general, certain broad characteristic indications can be noted in most all infected plant types. There is usually a stunting or dwarfing with accompanying pattern of light-yellowish areas merging with green patches, giving a "mottled" effect. Often there is a roughening of the surface, producing, in some cases, a warty appearance. Edges of leaves may cup downward. The fruit also may be mottled and warted.

Mosaic has become costly and destructive on many crops, especially peppers, tomatoes, cucumbers and squash. This disease apparently does not carry over in the soil, provided cover crops are turned under long enough to be decomposed before planting. It is carried from season to season in wild host plants (weeds) in and around the field, and is transmitted to the plants in the field by insects, particularly aphids and thrips.

Pepper and tomato seedbeds should have clean cultivation two or three weeks before planting the seed. The seedbeds should not be close to ditch banks or fence rows, unless the weeds near the beds are destroyed. Smoking or other use of tobacco should not be permitted while working in the seedbeds or handling plants. Tobacco mosaic has been known to be carried on tobacco. If plants become affected with mosaic in the seedbeds, do not set them in the field. Once the plant gets the disease there is no known cure.

The State Experiment Stations recognize the severity of the problem and they are now very much engaged in the attempt to develop effective control measures by chemical or biological means. A serious effort is also being manifested in the development of resistant varieties of various vegetables commonly susceptible to virus diseases. In the event of success, such new varieties would hardly be available commercially for several years, and we would again like to stress the importance of sanitary practices and effective insect and weed eradication as the only means of crop damage reduction at the present time.



# Safe Combinations of INSECTICIDES AND FUNGICIDES

| Insecticides               | Trade Names       | Can be used with   |
|----------------------------|-------------------|--|
| PARATHION .....            | (Vapophos) .....  | DDT, DDD, methoxychlor, chlordane, benzene hexachloride, lindane, toxaphene, aldrin, dieldrin, Dithane, "Parzate," "Zerlate," "Fermate," "Manzate," and sulphur. Most copper compounds can be used with parathion and malathion for spraying mixtures if mixed freshly and used immediately. |
| MALATHION .....            | .....             |  |
| DDT .....                  | (Kilcide) .....   | Each other, and also with nicotine, sulphur, copper, Dithane, "Parzate," "Zerlate," "Fermate," and "Manzate."  |
| DDD .....                  | (RHothane) .....  |  |
| METHOXYCHLOR .....         | ("Marlate") ..... |  |
| CHLORDANE .....            | (Kilchlor) .....  |  |
| BENZENE HEXACHLORIDE ..... | (Gamtox) .....    |  |
| LINDANE .....              | (Isotox) .....    |  |
| TOXAPHENE .....            | (Alltox) .....    |  |
| ALDRIN .....               | .....             |  |
| DIELDRIN .....             | .....             |  |
| TEPP .....                 | (Vapotone) .....  | DDT, DDD, methoxychlor, chlordane, benzene hexachloride, lindane, toxaphene, sulphur.  |
| NICOTINE .....             | .....             | DDT, toxaphene, sulphur, copper, Dithane, "Parzate."   |
| KRYOCIDE .....             | .....             | Sulphur, copper, Dithane, "Parzate."   |

## DONT'S

**A Reference to this table may prevent serious crop losses**

### DON'T USE:

### ON THE FOLLOWING CROPS:

|                            |  |
|----------------------------|--|
| DDT (Kilcide) .....        | Cucurbits (cucumbers, squash, melons, etc.).                             |
| DDD (RHothane) .....       | Cucurbits.   |
| TOXAPHENE (Kilphene) ..... | Cucurbits, young tomatoes*, tobacco, lettuce and escarolle**, avocados.  |
| (Alltox) .....             |  |
| B. H. C. (Gamtox) .....    | Cucurbits, corn, young tomatoes, potatoes and other tuber or root crops. |
| (Kilhex) .....             |  |
| LINDANE (Isotox) .....     | Corn, young cucurbits when heavy dew is present, potatoes and other      |
| (Kilgam) .....             | tuber or root crops.   |
| CHLORDANE (Kilchlor) ..... | Cucurbits, tobacco seedbeds and young tobacco plants in field.           |
| KRYOCIDE .....             | Corn, Asparagus plumosus.  |
| COPPER COMPOUNDS .....     | Corn.  |
| TEPP (Vapotone XX) .....   | Tomatoes.  |
| (Vapotone Dust) .....      |  |
| ALDRIN .....               | Don't use on any crops not approved by USDA. Consult local authorities.  |
| DIELDRIN .....             |  |

\* As toxaphene may prove toxic to young tomato plants in some local areas, local experiment stations should be consulted before treating.

\*\* Do not treat lettuce and escarolle within a period of three weeks from seeding or transplanting.

*Avoid combinations of emulsions and wettable powders. Many are not compatible physically.*



## SEED TREATMENT

(For Disease Control)

The following recommendations are suggested:  
Treatment for the prevention of damping-off fungi:

|                                 | Oz. per Bu.               | Teasps. per lb. |
|---------------------------------|---------------------------|-----------------|
| Beans, Lima .....               | Spergon                   | 2 1/3           |
| Beans, Snap and Pole .....      | Spergon                   | 2 1/3           |
| Peas, English .....             | Spergon                   | 2 1/3           |
| Cowpeas .....                   | Spergon                   | 2 1/3           |
| Corn .....                      | Arasan or Ortho Seedguard | 2 1/4           |
| Broccoli .....                  | Arasan                    | 1 1/2           |
| Cantaloupe .....                | Spergon or Semesan        | 1               |
| Carrot .....                    | Spergon                   | 3/4             |
| Celery .....                    | Spergon or Semesan        | 3/4             |
| Cabbage, Cauliflower, etc. .... | Arasan or Semesan         | 1 1/2           |
| Cucumber .....                  | Spergon or Semesan        | 1               |
| Eggplant .....                  | Semesan                   | 3/4             |
| Escarolle .....                 | Semesan                   | 1               |
| Lettuce .....                   | Semesan                   | 1               |
| Pepper .....                    | Semesan                   | 1 1/2           |
| Potato .....                    | Orthocide 50-W            | 4               |
| Spinach .....                   | Spergon or Arasan         | 1 1/2           |
| Tomato .....                    | Semesan                   | 1               |
| Watermelon .....                | Spergon or Semesan        | 1               |

The seed and seed treatment material should be thoroughly mixed together in a rotary drum mixer until all the seed is uniformly coated. In most cases this will take from three to five minutes at a speed of forty revolutions per minute. If Arasan is used on beans and peas, use one-half as much as the Spergon recommendation. Do not use more material than recommended. Screen off excess dust.

### Bichloride of Mercury (Corrosive sublimate) treatment affecting organism on surface of seeds:

Cucumbers: Dilute 1:1000, immersing seed for 10 minutes.

Less hard seeds: Dilute 1:1000, immersing seeds for 5 minutes.

For small quantities of seed dissolve 1 tablet in 1 pint of water.

For larger quantities dissolve 1 oz. of the powder form in 7 1/2 gallons of water. It is advisable to dissolve the powder in a small quantity (a pint or quart) of boiling water first.

Place seeds in loosely woven cloth bags (about half full). Submerge bags in solution and stir with a stick to loosen adhering bubbles. After soaking the proper length of time, remove bags and drain thoroughly. Seed should be washed for 15 minutes in several changes of water. Then spread in thin layers to dry.

Do not prepare solution of bichloride of mercury in metal containers. Only glass, earthenware or wooden vessels should be used. In working with the compound, the greatest caution should be exercised, as it is extremely poisonous.

### HOT WATER TREATMENT FOR CONTROL OF SEED-BORNE DISEASES

**BLACK ROT.** Cabbage seed which is not known to be free from the black rot organism should be soaked in water held at 122° F. for 25 minutes (cauliflower and broccoli, 18 minutes). Use an accurate thermometer and hold the temperature within 1° F. of that specified to control the disease without injuring the seed. Fill a cheesecloth bag two-thirds full of seed, tie the top, immerse it in the hot water and then stir it enough to remove the air. After treatment, dip the seed in cold water, drain and spread in a thin layer to dry.

Use new land for seedbeds or old land in which cabbage or other related crops have not been grown for two years.

Do not use plants from a plant bed in which black rot has been found.

## DILUTION TABLE FOR INSECTICIDES AND FUNGICIDES

Average amounts needed to make spray

| Material                               | 50 Gals.      | 4 Gals.    | 1 Gal.     |
|--|---------------|------------|------------|
| Aldrin 2 Spray .....                   | 1/2 pt.       | 8 tbs.     | 2 tbs.     |
| Alltox 40-W .....                      | 2 lbs.        | 3 3/4 tbs. | 4 tsp.     |
| Alltox Spray No. 48 .....              | 1 1/2 pts.    | 10 tbs.    | 2 1/2 tbs. |
| Arsenate of Lead .....                 | 1 1/2 lbs.    | 8 tsp.     | 2 tsp.     |
| Black Leaf 40 .....                    | 1 pt.         | 3 tbs.     | 2 1/4 tsp. |
| Ceresan .....                          | 1 lb.         | 3 tbs.     | 2 1/4 tsp. |
| C-M-Z Spray .....                      | 3 lbs.        | 8 tbs.     | 2 tbs.     |
| C-M-Z Garden Spray .....               | 2 1/2 lbs.    | 8 tbs.     | 2 tbs.     |
| Copper-A Compound .....                | 2 lbs.        | 8 tbs.     | 1 1/2 tsp. |
| Cuprocide—Yellow .....                 | 1 lb.         | 5 tbs.     | 4 tsp.     |
| DDT 25% Emulsion .....                 | 1 qt.         | 8 tbs.     | 2 tbs.     |
| DDT 50% Wettable .....                 | 1 1/4 lbs.    | 5 tbs.     | 4 tsp.     |
| Dieldrin Emulsion (1.5) .....          | Consult label | 5 tbs.     | 4 tsp.     |
| Dithane D-14 .....                     | 1 qt.         | 3 tsp.     | 3/4 tsp.   |
| (Zinc Sulphate) .....                  | 6 oz.         | 5 tbs.     | 4 tsp.     |
| Dithane M-22 .....                     | 1 lb.         | 5 tbs.     | 4 tsp.     |
| Dithane Z-78 .....                     | 1 lb.         | 5 tbs.     | 4 tsp.     |
| Fermate .....                          | 1 lb.         | 9 1/3 tbs. | 7 tsp.     |
| Fish Oil Soap .....                    | 1 1/2 lbs.    | 4 tbs.     | 1 tbs.     |
| Gamtox 10 Wettable .....               | 1 1/4 lbs.    | 8 tbs.     | 2 tbs.     |
| Isotox Wettable No. 25 .....           | 1/2 lb.       | 4 tbs.     | 1 tbs.     |
| Karathane WD .....                     | Consult label | 6 1/4 tbs. | 4 tsp.     |
| Kilchlor 40-W (Chlordane) .....        | 1 1/4 lbs.    | 8 tbs.     | 2 tbs.     |
| Kilcop "45" .....                      | 1 1/2 lbs.    | 10 tbs.    | 2 1/2 tbs. |
| Kilcop "53" .....                      | 2 lbs.        | 13 tbs.    | 3 1/3 tbs. |
| Kryocide .....                         | 3 lbs.        | 5 tbs.     | 4 tsp.     |
| Malathion "50" Spray .....             | 1 qt.         | 7 tbs.     | 5 tsp.     |
| Malathion 25% Wet. ....                | 1 lb.         | 6 2/3 tbs. | 5 tsp.     |
| Manganese Sulphate .....               | 1 1/2 lbs.    | 3 1/4 tbs. | 4 tsp.     |
| Manzate .....                          | 3/4 lb.       | 1 lb.      | 1/4 lb.    |
| Nicotine Pyrox .....                   | 14 lbs.       | 5 tbs.     | 4 tsp.     |
| Nicotrol .....                         | 1 qt.         | 10 1/2 oz. | 5 1/2 tbs. |
| Oil Emulsion (Fico) .....              | 1 gal.        | 12 tbs.    | 3 tbs.     |
| Orthocide 50-W .....                   | 2 lbs.        | 7 1/2 tbs. | 5 1/2 tsp. |
| Orthol Garden Spray .....              | 1 1/2 qts.    | 8 tbs.     | 2 tbs.     |
| Ortho-Klor 44 Spray (Chlordane) .....  | 1 1/2 qts.    | 8 tbs.     | 2 tbs.     |
| Ortho-Klor 8 Spray (Chlordane) .....   | Consult label | 3 oz.      | .....      |
| Orthophos 4 Spray (Parathion) .....    | 3 oz.         | .....      | .....      |
| Vapophos 15 Wettable (Parathion) ..... | 1/2 lb.       | 5 tbs.     | 4 tsp.     |
| Parzate .....                          | 1 lb.         | 5 tbs.     | 4 tsp.     |
| Penetrol .....                         | 1 qt.         | 3 tbs.     | 2 1/4 tsp. |
| Phygon-XL .....                        | 1 lb.         | 8 tsp.     | 2 tsp.     |
| Red Arrow Insect Spray .....           | 1 pt.         | 8 tbs.     | 2 tbs.     |
| RHothane WP-50 .....                   | 1 1/4 lbs.    | 2 pts.     | 1/2 pt.    |
| Scalecide .....                        | 3 1/3 gals.   | 11 tbs.    | 8 tsp.     |
| Spergon Wettable .....                 | 1 1/2 lbs.    | 5 tbs.     | 4 tsp.     |
| Sulfocide .....                        | 1 qt.         | 3/4 tsp.   | 20 drops   |
| Triton B-1956 .....                    | 3 tbs.        | .....      | .....      |
| Vapotone-XX (TEPP) .....               | 4 oz.         | 20 tbs.    | 5 tbs.     |
| Volck Isotox Spray .....               | 1 gal.        | 3 tbs.     | 2 tsp.     |
| Zinc Sulphate .....                    | 1 lb.         | .....      | .....      |

### BUREAU OF STANDARDS HOUSEHOLD MEASUREMENTS

|                       |                |                |
|-----------------------|----------------|----------------|
| Tsp.—level teaspoon   | Gal.—gallon    | 1 Cup—8 ounces |
| Tbs.—level tablespoon | 3 Tsp.—1 tbs.  | 2 Cups—1 pint  |
| Pt.—pint              | 2 Tbs.—1 ounce | 2 Pts.—1 quart |
| Qt.—quart             | 16 Tbs.—1 cup  | 4 Qts.—1 gal.  |
| Oz.—ounce             |                |                |

Sprays or dusts required per acre depend chiefly upon stage of growth, variety of crop and severity of insect infestation or disease infection. Spray for small plants require from 75 to 100 gals. per acre, for larger plants and heavy growth 100 to 150 gals. per acre. Dust for small plants 25 to 30 lbs. per acre, for large plants and heavy growth 35 to 50 lbs. per acre.



# INSECT AND DISEASE RECOMMENDATIONS

## BEANS—

### INSECTS

|                      |                             | Coined Name | Common Name        |
|----------------------|-----------------------------|-------------|--------------------|
| Leaf miners          | Kilphos Dust No. 2          | Parathion   | 20-30 lbs., acre   |
|                      | Vapophos Wettable 15%       | Parathion   | 1 lb., 100 gals.   |
|                      | Alltox Wettable 40          | Toxaphene   | 2½ lbs., 100 gals. |
| Bean leaf hoppers    | Kilphos Dust No. 2          | Parathion   | 25 lbs., acre      |
| Bean leaf rollers    | Kilcide 5-Sulphur Dust      | DDT         | 20-35 lbs., acre   |
| Thrips               | DDT Wettable—50%            |             | 2 lbs., 100 gals.  |
|                      | Kilphene 5-Sulphur Dust     | Toxaphene   | 20-30 lbs., acre   |
| Army worms           | Kilchlor-Kilphene Poison    | Chlordane   | 15-30 lbs., acre   |
|                      | Bait No. 11                 | Toxaphene   |                    |
| Cutworms             | Kilphene Poison Bait No. 10 | Toxaphene   | 15-30 lbs., acre   |
|                      | Kilchlor Poison Bait No. 8  | Chlordane   | 15-30 lbs., acre   |
| Mexican bean beetles | Kilphos Dust No. 2          | Parathion   | 20-30 lbs., acre   |
|                      | Rotenone Dust 1%            |             | 20-30 lbs., acre   |



Adult

BEAN LEAF HOPPER


### DISEASES

|                |                  |                       |
|----------------|------------------|-----------------------|
| Rust           | Sulphur Dust     | 20-40 lbs., acre      |
| Powdery mildew | Sulphur—Wettable | 10-16 lbs., 100 gals. |

## BROCCOLI, CABBAGE, CAULIFLOWER—

### INSECTS

|               |                             |           |                    |
|---------------|-----------------------------|-----------|--------------------|
| Aphids        | Kilphos Dust No. 3          | Parathion | 20-30 lbs., acre   |
|               | Kilgam Dust 1½%             | Lindane   | 20-30 lbs., acre   |
|               | Vapotone XX Spray           | TEPP      | 1 pt., 100 gals.   |
|               | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.   |
|               | Orthophos 4-Spray           | Parathion | 1/3 pt., 100 gals. |
| Cabbage worms | Kilchlor Dust—5%            | Chlordane | 25-35 lbs., acre   |
|               | Kilphene 10 Dust            | Toxaphene | 25-35 lbs., acre   |
|               | Kryocide 30-70 Dust         |           | 25-35 lbs., acre   |
|               | 25% DDT Emulsion            |           | 1 qt., 100 gals.   |
|               | Kilchlor 40-W               | Chlordane | 2½ lbs., 100 gals. |
| Leaf miners   | Kilphos Dust No. 2          | Parathion | 25-35 lbs., acre   |
|               | Kilchlor-Sulphur Dust No. 3 | Chlordane | 25-35 lbs., acre   |
|               | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.   |
| Cutworms      | Kilchlor-Kilphene Poison    | Chlordane | 15-30 lbs., acre   |
| Army worms    | Bait No. 11                 | Toxaphene |                    |
| Grasshoppers  | Kilphene Poison Bait No. 10 | Toxaphene | 15-30 lbs., acre   |
| Mole crickets | Kilchlor Poison Bait No. 8  | Chlordane | 15-30 lbs., acre   |



CABBAGE LOOPER



Adult

HARLEQUIN  
CABBAGE BUG

### DISEASES

|                    |                      |   |                                    |
|--------------------|----------------------|---|------------------------------------|
| For Seedbeds       | Spergon Dust No. 1   | Consult local Experiment Station<br>for rate of application |                                    |
|                    | Downy mildew         | Spergon Wettable  | 4 lbs., 100 gals.                  |
| For Crops in Field | Alternaria leaf spot | Dithane D-14 plus zinc                                      | Nabam 2 qts. plus ¾ lb., 100 gals. |
|                    | Downy mildew         | Dithane Z-78  | Zineb 2 lbs., 100 gals.            |

## CELERY—

### INSECTS

|                         |                             |           |                   |
|-------------------------|-----------------------------|-----------|-------------------|
| Green climbing cutworms | Alltox Wettable 40          | Toxaphene | 3 lbs., 100 gals. |
| Cutworms                | Alltox Wettable 40          | Toxaphene | 3 lbs., 100 gals. |
| Army worms              | Kilphene Poison Bait No. 10 | Toxaphene | 15-30 lbs., acre  |
| Celery leaf tiers       | Kilphene-10 Dust            | Toxaphene | 20-30 lbs., acre  |
| Celery loopers          | 25% DDT Emulsion            |           | 1 qt., 100 gals.  |
| Aphids                  | Kilphos Dust No. 1          | Parathion | 20-30 lbs., acre  |
|                         | 25% DDT Emulsion            |           | 1 qt., 100 gals.  |
|                         | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.  |

### DISEASES

|              |                        |        |                              |
|--------------|------------------------|--------|------------------------------|
| Early blight | Dithane D-14 plus zinc | Nabam  | 2 qts. plus ¾ lb., 100 gals. |
| Late blight  | Kilcop "53"            | Copper | 4 lbs., 100 gals.            |
|              | Copper A Compound      |        | 6 lbs., 100 gals.            |
|              | Manzate                |        | 1½ lbs., 100 gals.           |



IMPORTED CABBAGE WORM



CELERY LEAF-TIER



## INSECT AND DISEASE RECOMMENDATIONS (Continued)

### CORN (SWEET)—


#### INSECTS

#### Coined Common Name

|                    |                             |           |   |
|--------------------|-----------------------------|-----------|---|
| Corn ear worms     | Kilphos Dust No. 4          | Parathion | 30-35 lbs., acre                              |
|                    | Kilphene-10 Dust            | Toxaphene | 30-35 lbs., acre                              |
|                    | Kilcide 10% Dust            | DDT       | 30-35 lbs., acre                              |
|                    | Vapophos Wettable 15%       | Parathion | 2 lbs., 100 gals.                             |
|                    | Alltox Wettable 40          | Toxaphene | 2½ lbs., 100 gals.                            |
| Fall army worms    | DDT Wettable—50%            |           | 2 lbs., 100 gals.                             |
|                    | 25% DDT Emulsion            |           | 1 qt., 100 gals.                              |
|                    | Kilphos Dust No. 4          | Parathion | 30-35 lbs., acre                              |
| Corn lantern flies | Kilphene-10 Dust            | Toxaphene | 30-35 lbs., acre                              |
|                    | Kilchlor Dust 5%            | Chlordane | 30-35 lbs., acre                              |
|                    | Vapophos Wettable 15%       | Parathion | 2 lbs., 100 gals.                             |
|                    | Alltox Wettable 40          | Toxaphene | 2½ lbs., 100 gals.                            |
|                    | Kilchlor 40-W               | Chlordane | 2½ lbs., 100 gals.                            |
| Wire worms         | Kilphene Poison Bait No. 10 | Toxaphene | 15-30 lbs., acre                              |
|                    | Chlordane 50%               |           | 4 lbs., 100 gals.                             |
|                    | Aldrin Emulsion             |           | Broadcast 2 gals., acre<br>in 100 gals. water |
| Corn leaf blight   | Dithane Dust No. 2          | Zineb     | 25-30 lbs., acre                              |
|                    | Dithane Z-78                | Zineb     | 2 lbs., 100 gals.                             |
|                    | Dithane D-14 plus zinc      | Nabam     | 2 qts. plus ¾ lb., 100 gals.                  |



CORN EARWORM  
(TOMATO FRUITWORM)



WIRE WORM

### COWPEAS (SOUTHERN PEAS)—

#### INSECTS

|             |                         |           |                  |
|-------------|-------------------------|-----------|------------------|
| Pod weevils | Kilphene 5—Sulphur Dust | Toxaphene | 25-35 lbs., acre |
|             | Kilcide 5—Sulphur Dust  | DDT       | 20-35 lbs., acre |

#### DISEASES

|                      |                        |        |                              |
|----------------------|------------------------|--------|------------------------------|
| Scab and leaf blight | Kilcop Dust No. 4      | Copper | 25-35 lbs., acre             |
|                      | Dithane D-14 plus zinc | Nabam  | 2 qts. plus ¾ lb., 100 gals. |



Adult  
STRIPED CUCUMBER  
BEETLE

### CUCUMBER, SQUASH, CANTALOUPE—

#### INSECTS

|                  |                             |           |                   |
|------------------|-----------------------------|-----------|-------------------|
| Aphids           | Kilphos Dust No. 1          | Parathion | 25-35 lbs., acre  |
|                  | Kilgam Dust 1½%             | Lindane   | 25-35 lbs., acre  |
|                  | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.  |
|                  | Vapotone XX                 | TEPP      | 1 pt., 100 gals.  |
|                  | Isotox Wettable No. 25      | Lindane   | 1 lb., 100 gals.  |
| Leaf miners      | as above except Vapotone XX |           |                   |
| Cucumber beetles | Kilgam Dust—1½%             | Lindane   | 25-35 lbs., acre  |
|                  | Kryocide Dust 30-70         |           | 25-35 lbs., acre  |
| Pickle worms     | Kilphos Dust No. 1          | Parathion | 25-35 lbs., acre  |
|                  | Isotox Wettable No. 25      | Lindane   | 1 lb., 100 gals.  |
| Melon worms      | Kryocide Wettable           |           | 6 lbs., 100 gals. |
|                  | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.  |

#### DISEASES

|                   |                        |        |                              |
|-------------------|------------------------|--------|------------------------------|
| Downy mildew      | Dithane Dust No. 1     | Zineb  | 25-35 lbs., acre             |
|                   | Fermate Dust No. 1     | Ferbam | 25-35 lbs., acre             |
| Powdery mildew    | Dithane D-14 plus zinc | Nabam  | 2 qts. plus ¾ lb., 100 gals. |
|                   |                        |        |                              |
| Angular leaf spot |                        |        |                              |
| Anthracnose       | Orthocide 50-W         |        | 4 lbs., 100 gals.            |



SQUASH BUG

### EGGPLANT—

#### INSECTS

|             |                         |           |                    |
|-------------|-------------------------|-----------|--------------------|
| Aphids      | Kilphos Dust No. 2      | Parathion | 20-60 lbs., acre   |
|             | Kilhex-Sulphur Dust—1½% | BHC       | 20-60 lbs., acre   |
|             | Vapophos Wettable 15%   | Parathion | 1 lb., 100 gals.   |
|             | Vapotone XX             | TEPP      | 1 pt., 100 gals.   |
|             | Gamtox Wettable No. 10  | BHC       | 2½ lbs., 100 gals. |
| Red spiders | Dusting Sulphur         |           | 20-60 lbs., acre   |
|             | Vapotone XX             | TEPP      | 1 pt., 100 gals.   |
|             | Kilphos Dust No. 1      | Parathion | 20-60 lbs., acre   |
| Cutworms    | Kilchlor-Kilphene       | Chlordane | 15-30 lbs., acre   |
|             | Poison Bait No. 11      | Toxaphene |                    |

### LETTUCE, ESCAROLLE—

#### INSECTS

|          |                    |           |                  |
|----------|--------------------|-----------|------------------|
| Cutworms | Kilchlor-Kilphene  | Chlordane | 15-30 lbs., acre |
|          | Poison Bait No. 11 | Toxaphene |                  |



SQUASH VINE BORER



# INSECT AND DISEASE RECOMMENDATIONS (Continued)

## OKRA—

### INSECTS

|                         |                              | Coined Common Name |                    |
|-------------------------|------------------------------|--------------------|--------------------|
| Aphids .....            | Vaportone XX .....           | TEPP .....         | 1 pt., 100 gals.   |
|                         | Kilphos Dust No. 1 .....     | Parathion .....    | 25-35 lbs., acre   |
| Leaf miners .....       | Kilphene 10% Dust .....      | Toxaphene .....    | 25-35 lbs., acre   |
| Pumpkin bugs .....      |                              |                    |                    |
| Okra caterpillars ..... | Alltox Wettable No. 40 ..... | Toxaphene .....    | 2½ lbs., 100 gals. |



GREEN PUMPKIN BUG  
(SOUTHERN STINK BUG)

## ONION—

### INSECTS

|              |                              |           |                    |
|--------------|------------------------------|-----------|--------------------|
| Thrips ..... | Gamtox Wettable No. 10 ..... | BHC ..... | 2½ lbs., 100 gals. |
|              | DDT 25% Emulsion .....       |           | 1 qt., 100 gals.   |

## PEPPER—

### INSECTS

|                       |                              |                 |                    |
|-----------------------|------------------------------|-----------------|--------------------|
| Aphids .....          | Kilphos Dust No. 1 .....     | Parathion ..... | 20-30 lbs., acre   |
|                       | Vapophos Wettable 15% .....  | Parathion ..... | 1 lb., 100 gals.   |
|                       | Black Leaf 40 .....          |                 | 1 qt., 100 gals.   |
| Pepper weevils .....  | Kilhex Dust 1½% .....        | BHC .....       | 25-30 lbs., acre   |
| Fall army worms ..... | DDT 25% Emulsion .....       |                 | 1 qt., 100 gals.   |
| Blister beetles ..... | Gamtox Wettable No. 10 ..... | BHC .....       | 2½ lbs., 100 gals. |



PEPPER WEEVIL

### DISEASES

|                            |                              |              |                           |
|----------------------------|------------------------------|--------------|---------------------------|
| Cercospora leaf spot ..... | Kilcop Dust No. 6 .....      | Copper ..... | 30-35 lbs., acre          |
|                            | Kilcop "53" .....            | Copper ..... | 5 lbs., 100 gals.         |
| Bacterial leaf spot .....  | Kilcop Dust No. 6 .....      | Copper ..... | 30-35 lbs., acre          |
| Phytophthora blight .....  | Dithane D-14 plus zinc ..... | Nabam .....  | 2 qts. + ¾ lb., 100 gals. |
|                            | Dithane Z-78 .....           | Zineb .....  | 2 lbs., 100 gals.         |
| Damp-off .....             | Orthocide 50-W .....         |              | 4 lbs., 100 gals.         |
|                            | Yellow Cuprocid .....        | Copper ..... | 2 lbs., 100 gals.         |

## POTATOES—

### INSECTS

|                               |  |                 |                    |
|-------------------------------|--|-----------------|--------------------|
| Aphids .....                  | Kilphos Dust No. 1 .....                   | Parathion ..... | 20-40 lbs., acre   |
| Army worms .....              | Kilphene-5 Dust .....                      | Toxaphene ..... | 20-40 lbs., acre   |
| Cutworms .....                | Vapophos Wettable 15% .....                | Parathion ..... | 1 lb., 100 gals.   |
| Colorado potato beetles ..... | Alltox Wettable No. 40 .....               | Toxaphene ..... | 2½ lbs., 100 gals. |
|                               | DDT Emulsion—25% .....                     |                 | 1 qt., 100 gals.   |
| Leaf miners .....             | Same as above except omit DDT Emulsion 25% |                 |                    |
| Pumpkin bugs .....            |  |                 |                    |
| Leaf-footed plant bugs .....  |  |                 |                    |

### DISEASES

|                    |                              |              |                           |
|--------------------|------------------------------|--------------|---------------------------|
| Late blight .....  | Dithane Dust No. 2 .....     | Zineb .....  | 25-35 lbs., acre          |
|                    | Kilcop Dust No. 6 .....      | Copper ..... | 25-35 lbs., acre          |
| Early blight ..... | Dithane D-14 plus zinc ..... | Nabam .....  | 2 qts. + ¾ lb., 100 gals. |
|                    | Dithane Z-78 .....           | Zineb .....  | 2 lbs., 100 gals.         |
|                    | Kilcop "53" .....            | Copper ..... | 4 lbs., 100 gals.         |



Adult  
COLORADO POTATO BEETLE

## POTATOES (SWEET)—

### INSECTS

|                    |                              |                 |                    |
|--------------------|------------------------------|-----------------|--------------------|
| Caterpillars ..... | Kilphene-10 Dust .....       | Toxaphene ..... | 25-35 lbs., acre   |
| Gold bugs .....    |                              |                 |                    |
| Grasshoppers ..... | Alltox Wettable No. 40 ..... | Toxaphene ..... | 2½ lbs., 100 gals. |

## RADISH, TURNIP, MUSTARD—

### INSECTS

|              |                         |                 |                  |
|--------------|-------------------------|-----------------|------------------|
| Aphids ..... | Pyrocide Dust—7½% ..... | Pyrethrum ..... | 15-35 lbs., acre |
|              | Vaportone XX .....      | TEPP .....      | 1 pt., 100 gals. |



Wingless Adult  
APHID



# INSECT AND DISEASE RECOMMENDATIONS (Continued)

## STRAWBERRIES—

### INSECTS


|                  | Coined Common Name                        |                            |
|------------------|---|----------------------------|
| Thrips           | Malathion Dust No. 1                      | 20-25 lbs., acre           |
|                  | Kilphene-5 Sulphur Dust                   | 20-25 lbs., acre           |
| Pameras          |   |                            |
| Ants             |   |                            |
| Flea beetles     |   |                            |
| Cucumber beetles | Kilchlor-Sulphur Dust No. 3               | Chlordane 20-25 lbs., acre |
| Leaf rollers     |   |                            |
| Grasshoppers     |   |                            |
| Mole crickets    | Kilchlor-Sulphur Dust No. 3               | Chlordane 20-25 lbs., acre |
| Corn borers      | Kilchlor Poison Bait No. 8                | Chlordane 15-25 lbs., acre |
| Cutworms         | Kilchlor-Kilphene Poison                  | Chlordane                  |
| Field crickets   | Bait No. 11                               | Toxaphene 20-25 lbs., acre |
| Red spiders      | Dusting Sulphur                           | 20-30 lbs., acre           |
|                  | Malathion Wettable 25-W                   | 2 lbs., 100 gals.          |
|                  | Malathion Dust No. 1                      | 20-30 lbs., acre           |
|                  | Note: Use Malathion before fruiting only. |                            |



RED SPIDER

### DISEASES

|             |                    |        |                    |
|-------------|--------------------|--------|--------------------|
| Anthracnose | Kilcop Dust No. 2  | Copper | 20-30 lbs., acre   |
| Leaf spots  | Kilcop "53"        | Copper | 4 lbs., 100 gals.  |
|             | Dithane Dust No. 1 | Zineb  | 20-30 lbs., acre   |
|             | Dithane M-22       | Maneb  | 1½ lbs., 100 gals. |



CUTWORM

## TOMATOES—

### INSECTS

|                        |                             |           |                   |
|------------------------|-----------------------------|-----------|-------------------|
| Aphids                 | Kilphos Dust No. 1          | Parathion | 25-35 lbs., acre  |
| Thrips                 | Kilhex Dust 1½%             | BHC       | 25-35 lbs., acre  |
|                        | Kilgam Dust 1½%             | Lindane   | 25-35 lbs., acre  |
|                        | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.  |
|                        | Gamtox Wettable No. 10      | BHC       | 2 lbs., 100 gals. |
|                        | Isotox Wettable No. 25      | Lindane   | 1 lb., 100 gals.  |
| Army worms             | Kilphos Dust No. 1          | Parathion | 25-35 lbs., acre  |
| Cutworms               | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.  |
| Tomato fruit worms     | DDT 25% Emulsion            |           | 1 qt., 100 gals.  |
| Pin worms              | DDT 50% Wettable            |           | 2 lbs., 100 gals. |
|                        | RHothane WP-50              | TDE—DDD   | 2 lbs., 100 gals. |
|                        | Kilchlor Poison No. 9       | Chlordane | 15-25 lbs., acre  |
| Leaf miners            | Kilchlor-Kilphene Poison    | Chlordane | 20-25 lbs., acre  |
| Pumpkin bugs           | Bait No. 11                 | Toxaphene |                   |
| Leaf-footed plant bugs | Kilphos Dust No. 1          | Parathion | 25-35 lbs., acre  |
|                        | Vapophos Wettable 15%       | Parathion | 1 lb., 100 gals.  |
| Horn worms             | Kryocide-Sulphur Dust 30-70 |           | 35-45 lbs., acre  |
|                        | RHothane 25% Emulsion       | TDE—DDD   | 1 qt., 100 gals.  |



TOMATO PINWORM



Nymph  
THRIPS

### DISEASES

|                           |                            |        |                           |
|---------------------------|----------------------------|--------|---------------------------|
| Late blight               | Dithane-Sulphur Dust No. 4 | Zineb  | 30-40 lbs., acre          |
| Early blight              | Kilcop-Sulphur Dust No. 5  | Copper | 30-40 lbs., acre          |
| Gray leaf-spot (Red rust) | Dithane D-14 plus zinc     | Nabam  | 2 qts. + ¾ lb., 100 gals. |
|                           | Kilcop "53"                |        | 4 lbs., 100 gals.         |
|                           | Dithane Z-78               | Zineb  | 2 lbs., 100 gals.         |
|                           | Phygon XL                  |        | ¾ lb., 100 gals.          |

## WATERMELON—

### INSECTS

|                                       |                       |           |                  |
|---------------------------------------|-----------------------|-----------|------------------|
| Pumpkin bug                           |                       |           |                  |
| Leaf-footed plant bug                 |                       |           |                  |
| Squash bug                            | Kilphos Dust No. 1    | Parathion | 20-35 lbs., acre |
| Other insects same as Cucumbers, etc. | Vapophos Wettable 15% | Parathion | 1 lb., 100 gals. |
| Page 54                               |                       |           |                  |

### DISEASES

|                   |                    |        |                   |
|-------------------|--------------------|--------|-------------------|
| Downy mildew      | Dithane Dust No. 1 | Zineb  | 30-40 lbs., acre  |
| Anthracnose       | Kilcop Dust No. 5  | Copper | 30-40 lbs., acre  |
| Gummy stem blight | Dithane Z-78       | Zineb  | 2 lbs., 100 gals. |
|                   | Kilcop "53"        | Copper | 4 lbs., 100 gals. |



LEAF-FOOTED PLANT BUG



# INSECT AND DISEASE RECOMMENDATIONS (Continued)

## TOBACCO—

### INSECTS

|               |                             | Coined Common Name |                   |
|---------------|-----------------------------|--------------------|-------------------|
| Aphids        | Kilphos Dust No. 1          | Parathion          | 10-35 lbs., acre  |
|               | Vapophos Wettable 15%       | Parathion          | 1 lb., 100 gals.  |
|               | Vapotone XX                 | TEPP               | 1 pt., 100 gals.  |
| Bud worms     | RHothane 5-Dust             | TDE-DDD            | 15-35 lbs., acre  |
|               | DDT 50% Wettable            |                    | 2 lbs., 100 gals. |
|               | RHothane WP-50              | TDE-DDD            | 2 lbs., 100 gals. |
| Horn worms    | RHothane 5-Tobacco Dust     | TDE-DDD            | 25-30 lbs., acre  |
|               | Kryocide Dust 30-70         |                    | 15-35 lbs., acre  |
|               | RHothane WP-50              | TDE-DDD            | 2 lbs., 100 gals. |
| Mole crickets | Kilchlor Poison Bait No. 8* | Chlordane          | 15-25 lbs., acre  |

\* CAUTION: Do not use directly on plants.  
Scatter in paths or walk ways.



Adult  
MOLE CRICKET

### DISEASES—PLANT BEDS

|           |                       |        |                              |
|-----------|-----------------------|--------|------------------------------|
| Blue mold | Dithane Dust No. 2    | Zineb  | 1½ to 4 lbs. to 100 sq. yds. |
|           | Fermate Dust No. 2    | Ferbam | as required by size of plant |
|           | Dithane Z-78 Wettable | Zineb  | 4 lbs., 100 gals.            |
|           | Fermate Wettable      | Ferbam | 3 lbs., 100 gals.            |

## PEANUTS—

### INSECTS

|                          |                             |     |                  |
|--------------------------|-----------------------------|-----|------------------|
| Velvet bean caterpillars | Kilcide 5—Sulphur Dust      | DDT | 25-35 lbs., acre |
|                          | Kryocide—Sulphur Dust 30-70 |     | 25-35 lbs., acre |

### DISEASES

|           |                 |  |                  |
|-----------|-----------------|--|------------------|
| Leaf spot | Dusting Sulphur |  | 15-25 lbs., acre |
|-----------|-----------------|--|------------------|



Adult  
THRIPS

## GLADIOLUS—

### INSECTS

|            |   |           |                                    |
|------------|---|-----------|------------------------------------|
| Thrips     | Kilcide-Kilhex Dust No. 3*                      | DDT-BHC   | 25-35 lbs., acre                   |
|            | DDT 50% W in combination with Gamtox W No. 10*  | BHC       | 2 lbs. + 1¼ lbs. Gamtox, 100 gals. |
| Army worms | DDT 50% Wettable                                |           | 4 lbs., 100 gals.                  |
|            | Dieldrin Emulsion (1.5 lbs. technical per gal.) |           | 3 pts., 100 gals.                  |
| Red spider | Kilphos Dust No. 2                              | Parathion | 25-35 lbs., acre                   |
|            | Vapophos Wettable 15%                           | Parathion | 1 lb., 100 gals.                   |
|            | Wettable Sulphur                                |           | 12 lbs., 100 gals.                 |

### DISEASES

|             |                        |       |                           |
|-------------|------------------------|-------|---------------------------|
| Curvularia  | Dithane D-14 plus zinc | Nabam | 2 qts. + ¾ lb., 100 gals. |
| Botrytis    |                        |       |                           |
| Stemphylium |                        |       |                           |

\* CAUTION: Kilhex and Gamtox Wettable may give spikes a musty odor if sprayed just prior to shipping.



Adult  
CHINCH BUG

## LAWNS—

### INSECTS

|               |                      |  |
|---------------|----------------------|--|
| Chinch bugs   | Kilgore's Kil-Chinch | Treat infested areas with 2 lbs., 100 sq. ft.  |
| Army worms    |                      |  |
| Sod web worms | Saba Chinch          | Treat infested areas with 4 lbs., 1000 sq. ft. |
| Mole crickets |                      |  |
| Ants          |                      |  |

NOTE: See our 1954-55 Annual Flower Guide for Florida, pages 27, 28, 29.

## PASTURES—

### INSECTS

|              |                          |           |                  |
|--------------|--------------------------|-----------|------------------|
| Aphids       | Vapotone XX              | TEPP      | 1 pt., 100 gals. |
|              | Vapophos 15%             | Parathion | 1 lb., 100 gals. |
| Army worms   | RHothane 5-Sulphur Dust* | TDE-DDD   | 20-30 lbs., acre |
| Grasshoppers | Kilphene 10-Dust*        | Toxaphene | 20-30 lbs., acre |

\* CAUTION: Keep cattle off pastures for 2 weeks after applying these insecticides. Do not use in pastures for dairy cows.



Nymph  
CHINCH BUG

The Kilgore Seed Company processes many combinations of insecticides and fungicides to suit local conditions which are too numerous to recommend in our General Catalog. If you do not find the combination you want, write our General Office in Plant City or call at your nearest Kilgore Store; it is likely we will have a dust in stock to fill your requirements.



## INSECT AND DISEASE RECOMMENDATIONS (Continued)

### SOIL BORNE PESTS—

|  |   |  |
|--|---|--|
| Nematodes<br>(Root knot)<br>Wire worms | D.D. Soil fumigant.....                               | 22 gals., acre in 12 inch rows.<br>5 to 6 gals., acre in 4 foot rows.<br>Apply 15 days before planting.  |
|  |   |  |
|  | Soilfume 80-20 .....                                  | 30 gals., acre in 12 inch rows.<br>8 to 10 gals., acre in 4 foot rows.<br>Apply 15 days before planting. |
| Wire worms only .....                  | Aldrin Emulsion<br>(2 lbs. technical per gallon)..... | 2 gals., acre in 100 gals. water   |
|  | Ortho-Klor 8 .....                                    | 2 qts., acre in 100 gals. water  |




WIRE WORM

## Termite Control

Coined Common  
Name

|                |                          |                |                    |
|----------------|--------------------------|----------------|--------------------|
| Termites ..... | Ortho-Klor 8 Spray.....  | Chlordane..... | 5 qts., 100 gals.  |
|                | Ortho-Klor 44 Spray..... | Chlordane..... | 2 gals., 100 gals. |



SUBTERRANEAN TERMITE  
WING ADULT QUEEN

Apply to infected soil areas with low pressure spraying equipment. Thoroughly soak area. Where outside trenching operations next to foundation walls are desired, dig trench 1 to 2 ft. deep and 8" to 12" wide. Apply at the rate of one gallon per linear foot of trench. Also treat soil as trench is being refilled. If above recommendations are followed, present results show that the protection period should be at least five years.

## Livestock Pests

### DAIRY CATTLE—

Coined Common  
Name

|             |                                |                                  |
|-------------|--------------------------------|----------------------------------|
| Flies ..... | "Marlate" 50-Insecticide ..... | Methoxychlor. 8¾ lbs., 100 gals. |
| Lice .....  | "Marlate" 50-Insecticide ..... | Methoxychlor. 25 lbs., 100 gals. |

### DAIRY BARNs AND MILK SHEDS—

|                   |   |                                |                                  |                         |
|-------------------|---|--------------------------------|----------------------------------|-------------------------|
| Flies .....       | { | Isotox Dairy Spray.....        | Lindane.....                     | 10 lbs., 100 gals.      |
|                   |   | "Marlate" 50-Insecticide ..... | Methoxychlor. 25 lbs., 100 gals. |                         |
|                   |   | Ortho Fly Killer.....          |                                  | Follow label directions |
| Screw worms ..... |   | See Range Cattle.....          |                                  |                         |

**CAUTION:** All feed troughs, feed and utensils should be covered before spraying.



ADULT CATTLE  
FEVER TICK

### RANGE CATTLE—

#### SPRAYS

(Use one of following)

|                  |  |                                 |
|------------------|--|---------------------------------|
| Horn flies ..... | DDT 50% Wettable.....                            | 8 lbs., 100 gals.               |
|                  | Gamtox Wettable No. 10.....                      | BHC..... 3¾ lbs., 100 gals.     |
|                  | Isotox Dairy Spray.....                          | Lindane..... 1½ lbs., 100 gals. |
|                  | "DuPont Livestock Spray<br>and Dip No. 30" ..... | 11 lbs., 100 gals.              |

(Use one of following)

|                   |                             |                             |
|-------------------|-----------------------------|-----------------------------|
| Mange mites ..... | Kilchlor 40-W .....         | 5 lbs., 100 gals.           |
|                   | Gamtox Wettable No. 10..... | BHC..... 2½ lbs., 100 gals. |

#### DIPS\*

(Use one of following)

|                              |                             |                             |
|------------------------------|-----------------------------|-----------------------------|
| General purpose<br>dip ..... | DDT 50% Wettable.....       | 8½ lbs., 100 gals.          |
|                              | Gamtox Wettable No. 10..... | BHC..... 2½ lbs., 100 gals. |

Combination of: .....

|   |                                 |
|---|---------------------------------|
| Ortho Healthy Herd<br>Wettable Powder ..... | BHC & DDT... 10 lbs., 100 gals. |
|---|---------------------------------|

|  |  |
|--|--|
| "DuPont Livestock Spray<br>and Dip No. 30" ..... | 11 lbs., 100 gals.                                   |
| Ortho Kleen Stock Spray .....                    | Lindale & See label for<br>Toxaphene..... directions |

**\*CAUTION:** In using dipping vats, all traces of arsenicals, oils or sludge from such materials must be thoroughly removed before charging with Gamtox, Isotox, "Marlate" or DDT. A mixture of arsenicals or oil solutions with Gamtox, Isotox, "Marlate," or DDT is highly toxic to animals and may cause death. Directions on all containers should be carefully followed.

|                   |  |                         |
|-------------------|--|-------------------------|
| Screw worms ..... | "Smearex," Screw Worm Preparation..... | Follow label directions |
|                   | "Ortho 1038 Screw Worm Control".....   |                         |
|                   | Ortho EQ 335 Screw Worm Remedy.....    |                         |



ADULT  
HORN FLY



# PESTICIDES

For household and garden insecticides, see Page 63

When ordering pesticides allow enough for postage because all prices quoted are f.o.b. Plant City or any Kilgore store unless otherwise specified. Inquire of your postmaster for proper parcel post rates.

All pesticides not used for commercial agricultural purposes are subject to 3% State Sales Tax.

All prices quoted are subject to change without notice.

## INSECTICIDES

**ALDRIN 2 SPRAY**—A liquid formulation containing 2 lbs. aldrin per gallon. 1 gal. \$4.65; 5-gal. can \$22.25; 30 gals. \$130.50; 50 gals. \$207.50.

**ARAMITE**—15-W. An effective miticide. Used on citrus, ornamentals and vegetables. Controls red spider mites. Use 1½ lbs. per 100 gals. Price 4-lb. bag \$2.88; 50 lbs. \$34.50.

**ARSENATE OF LEAD POWDER.** Use 6 lbs. to 100 gals. water. Pound canister 55c; 4 lbs. \$1.40.

**BENZENE HEXACHLORIDE PREPARATIONS.**

**GAMTOX WETTABLE**—No. 10 (Contains 10% gamma isomer.) Use 2½ lbs. per 100 gals. water. 5-lb. bag \$1.60; 50-lb. bag \$14.50.

**KILHEX DUST**—1½% (Code No. G-3)—(Gamtox) Contains 1½% gamma isomer. Lb. 10c; 25 lbs. \$1.90; 50 lbs. \$2.75; 100 lbs. \$5.50.

**KILHEX-SULPHUR DUST**—(Code No. G-4)—(Gamtox) Contains 1½% gamma isomer. Lb. 10c; 25 lbs. \$2.15; 50 lbs. \$3.30; 100 lbs. \$6.60.

**GAMTOX WETTABLE**—No. 10. (Contains 10% gamma isomer.) Use 2½ lbs. per 100 gals. water. 5-lb. bag \$1.60; 50-lb. bag \$14.50.

### LINDANE PREPARATIONS


**ISOTOX SPRAY**—No. 200 (Contains 20% gamma isomer). Use 1 pt. per 100 gals. water. 4-oz. bottle 95c; 16-oz. bottle \$2.35; 1 gal. \$15.25.

**ISOTOX**—No. 25-W (Contains 25% gamma isomer). Use 1 lb. per 100 gals. water. For cucumbers, squash, melons, etc. 1-lb. can \$2.40; 4-lb. can \$9.40; 20 lbs. \$41.00.

**KILGAM DUST**—1% (Code No. Q-4)—(Isotox) Contains 1% gamma isomer. Lb. 15c; 25 lbs. \$3.00; 50 lbs. \$5.00; 100 lbs. \$10.00.

**KILGAM DUST**—1½% (Code No. Q-5)—(Isotox) Contains 1½% gamma isomer. Lb. 20c; 25 lbs. \$3.90; 50 lbs. \$6.85; 100 lbs. \$13.70.

**KILGAM-SULPHUR DUST**—No. 6 (Code No. Q-6) (Isotox) Contains 1% gamma isomer. Lb. 15c; 25 lbs. \$3.25; 50 lbs. \$5.50; 100 lbs. \$11.00.



**BLACK LEAF 40.** For the control of aphids, thrips. Use from 1 to 2 pts. per 100 gallons water. Is effective as a spray against poultry lice. 1 oz. 39c; 5 oz. \$1.19; 1-lb. \$2.69; 2 lbs. \$4.79; 5 lbs. \$7.60; 10 lbs. \$14.70. Prices quoted are prepaid.

**CALCIUM ARSENATE POWDER.** 1-lb. can 45c; 4-lb. bag 65c.

### CHLORDANE PREPARATIONS

**KILCHLOR 40-W.** (Contains 40% technical chlordane). Use 2½ lbs. per 100 gals. of water. 1-lb. can 75c; 5-lb. bag \$3.00; 10-lb. bag \$6.00; 25 lbs. \$13.75; 50 lbs. \$25.00.

**ORTHO-KLOR "44"**—(Containing 44% liquid chlordane). Readily emulsifiable in water. For control of ants, mole crickets, cutworms, army worms, grasshoppers, roaches and many other insects. 4 oz. 90c; 8 oz. \$1.45; 16 oz. \$2.45; 32 oz. \$4.25.

**ORTHO-KLOR-8 SPRAY.** (Contains 72% chlordane) 1 gal. \$10.75; 5 gals. \$52.50. Use 2 qts. per 100 gallons for soil applications.

**KILCHLOR 5% DUST** (Code No. H-1)—(Contains 5% chlordane). Lb. 15c; 25 lbs. \$2.50; 50 lbs. \$3.95; 100 lbs. \$7.90.

**KILCHLOR-SULPHUR DUST NO. 3** (Code No. H-3)—(Contains 5% chlordane). Lb. 15c; 25 lbs. \$2.75; 50 lbs. \$4.50; 100 lbs. \$9.00.

**CORN EARWORM OIL (KILGORE'S).** An effective material for individual treatment of ears. Does not contain poisonous substances. 1 qt. 85c; 1 gal. \$2.75; 5 gals. \$13.00; 55-gal. drum \$137.50. (1 gal. of corn earworm oil will treat approximately 7,000 ears.)

### DDT PREPARATIONS

**DDT 50% WETTABLE POWDER.** Use 2 lbs. to 100 gals. water. 1-lb. canister 70c; 3-lb. bag 90c; 4 lbs. \$1.20; 6-lb. bag \$1.80; 50-lb. bag \$13.50.

**DDT 25% EMULSION.** Use 2 qts. per 100 gals. water. 5 gals. \$12.50; 30-gal. drum \$67.50; 50 gals. \$107.50.

**KILCIDE 5 DUST**—(Code No. D-3)—(Contains 5% DDT). Lb. 10c; 25 lbs. \$1.65; 50 lbs. \$2.30; 100 lbs. \$4.60.

**KILCIDE 5-SULPHUR DUST**—(Code No. D-10)—(Contains 5% DDT). Lb. 10c; 25 lbs. \$1.85; 50 lbs. \$2.75; 100 lbs. \$5.50.

**KILCIDE 10 DUST**—(Code No. D-4)—(Contains 10% DDT). Lb. 10c; 25 lbs. \$2.20; 50 lbs. \$3.35; 100 lbs. \$6.70.

**KILCIDE 3-MANGANESE-SULPHUR DUST**—(Code No. DX-1)—(Contains 3% DDT). Lb. 10c; 25 lbs. \$1.80; 50 lbs. \$2.60; 100 lbs. \$5.20.

**KILCIDE 5-MANGANESE-SULPHUR DUST**—(Code No. DX-2)—(Contains 5% DDT). Lb. 10c; 25 lbs. \$2.00; 50 lbs. \$3.00; 100 lbs. \$6.00.

**DIELDRIN EMULSION.** (Contains 1.5 lbs. technical dieldrin per gal.) 1 gal. \$8.00; 5 gals. \$38.75; 50 gals. \$375.00.

**KARATHANE.** 3-lb. bag \$5.70.



# INSECTICIDES (Continued)

**KRYOCIDE (Natural Cryolite).** A fluorine spray for chewing insects. Use 3 pounds Kryocide to fifty gallons of water. 1-lb. can 45c; 4-lb. bag \$1.00; 50-lb. bag \$9.75.

**KRYOCIDE DUST 30-70 (Code No. K-6)**—(Contains 30% Kryocide). A fluorine dust especially recommended for worms on cucumbers and squash, pinworms, and horn worms on eggplants and tomatoes. Lb. 15c; 25 lbs. \$2.45; 50 lbs. \$3.90; 100 lbs. \$7.80.

**KRYOCIDE-SULPHUR DUST 30-70 (Code No. K-4)**—(Contains 30% Kryocide.) Lb. 15c; 25 lbs. \$2.70; 50 lbs. \$4.40; 100 lbs. \$8.80.

**MALATHION EMULSION.** (Contains 5 lbs. technical malathion per gal.) Use 1 qt. per 100 gals. water. 1 oz. 49c; 5 oz. \$1.39; 1 pt. \$2.98; 1 qt. \$5.29; 1 gal. \$12.35; 5 gals. \$60.25.

**MALATHION. 25% Wettable Powders.** Use 2 lbs. per 100 gals. water. 1 lb. 79c; 5 lbs. \$3.35; 50 lbs. \$31.50; 100 lbs. \$62.00.

**LIME-SULPHUR SOLUTION.** (32 to 34 degrees Baume.) Dilute with water at 1 to 60. 1-gal. can 65c; 5-gal. can \$2.75. Write for special prices on 55-gallon barrels.

**OIL EMULSION 60.** For the control of scales, mealy bugs, red spiders, etc. Dilute with water 1 to 60. Gal. 80c; 5 gals. \$3.25; 55 gals. \$22.00. Deposit on containers.

**ORTHOL GARDEN SPRAY.** A modernized oil emulsion containing nicotine and DDT for the combined control of aphids, red spiders, mealy bugs, thrips and many other garden insects. Use at the rate of 6 table-spoonfuls to 1 gal. of water. 4 oz. 35c; 16 oz. 75c; 32 oz. \$1.25; 1 gal. \$2.85.

**PARA-SCALECIDE.** Effective and safe liquid treatment for peach borers in young and old trees. Dilute 1 gal. Para-Scalecide with 7 gals. water. This will treat from 50 to 100 trees depending on the size. Write for descriptive pamphlets on Para-Scalecide. Qt. \$1.45; gal. \$3.25; 5 gals. \$12.50.

**PARATHION PREPARATIONS.** Parathion products are highly toxic to man and warm blooded animals. It is urged that all precautions and recommendations stated on labels affixed to all of our parathion products be carefully followed.

**VAPOPHOS WETTABLE 15%.** A wettable powder containing 15% parathion. Use at the rate of 1 lb. to 100 gals. of water. 1 lb. 56c; 3-lb. bag \$1.53; 4-lb. bag \$2.04; 5-lb. bag \$2.55; 100-lb. drum \$50.00.

**ORTHOPHOS 4 SPRAY.** (Contains 42% parathion per gal.) Use 1/3 pt. per 100 gallons of water. 1 gal. \$22.00; 5 gals. \$108.75.

**KILPHOS DUST NO. 1.** (Code No. E-1)—(Contains 1% parathion. 50 lbs. \$3.00; 100 lbs. \$6.00.

**KILPHOS DUST NO. 2.** (Code No. E-2)—Contains 1% parathion and wettable sulphur. 50 lbs. \$3.55; 100 lbs. \$7.10.

**KILPHOS DUST NO. 4.** (Code No. E-4)—Contains 2% parathion. 50 lbs. \$4.35; 100 lbs. \$8.70.

**KILPHOS-KILPHENE-SULPHUR DUST No. 2** (Code No. EI-2)—Containing 75% parathion and 5% toxaphene. 50 lbs. \$4.25; 100 lbs. \$8.50.

**PARIS GREEN.** 1/4 lb. 30c; 1 lb. 80c; 5 lbs. \$3.45.

**POISON BAITS.** Ready for use to control mole crickets, cutworms, grasshoppers and field crickets. Also ideal for airplane application. Do not use any of the poison baits No. 10 and 11 on cucumbers, squash and melons.

**POISON BAIT NO. 8** (Code No. B-8)—(Contains chlordane). 1 lb. 10c; 25 lbs. \$1.95; 50 lbs. \$3.45; 100 lbs. \$6.90.

**POISON BAIT NO. 10** (Code No. B-10)—(Contains toxaphene). 1 lb. 10c; 25 lbs. \$1.90; 50 lbs. \$3.40; 100 lbs. \$6.80.

**POISON BAIT NO. 11** (Code No. B-11)—(Contains chlordane and toxaphene). 1 lb. 15c; 25 lbs. \$2.30; 50 lbs. \$4.15; 100 lbs. \$8.30.

**RED ARROW INSECT SPRAY.** Kills garden aphids, melon aphids, worms, leaf hoppers, white flies, red spiders, caterpillars, thrips, celery leaf tiers. 1 oz. 35c; 4 oz. \$1.00; pint \$2.85; quart \$5.00; gal. \$18.20. Average dilution 1 to 400 parts water.

**RHOTHANE.** A compound closely related to DDT. Its toxicity to most insects is equal to that of DDT and data obtained from experimental stations and other sources strongly indicates less toxicity to humans and warm blooded animals.



**RHOTHANE WP-50.** A wettable form of powder, containing 50% of the technical material. Use at the rate of 2 lbs. to 100 gals. water. 4-lb. bag \$2.00; 50-lb. bag \$23.00.

**RHOTHANE EMULSION CONCENTRATE.** (Contains 25% of the technical material.) 1 gal. \$3.05; 5 gals. \$14.00; 30-gal. drum \$78.00; 50-gal. drum \$125.00.

**RHOTHANE 5-TOBACCO DUST.** (Code No. J-2)—(Contains 5% DDD.) 1 lb. 10c; 25 lbs. \$2.25; 50 lbs. \$3.55; 100 lbs. \$7.10.

**ROTENONE DUST, 1%.** (Code No. R-6)—1 lb. 15c; 25 lbs. \$3.00; 50 lbs. \$5.00; 100 lbs. \$10.00.

**SCALECIDE.** Excellent for citrus. Combined with Sulfocide at the rate of two gallons Scalecide and one gallon Sulfocide to two hundred gallons of water. It takes care of many citrus troubles. Pt. 75c; qt. \$1.15; gal. \$2.75; 5 gals. \$10.75.

## TOXAPHENE PREPARATIONS

**ALLTOX 40W.** (Contains 40% toxaphene.) Use from 3 to 4 lbs. per 100 gals. of water. 6-lb. bag \$1.50; 50-lb. bag \$11.00.

**ALLTOX SPRAY NO. 4.8.** (Contains 50% toxaphene.) Use from 1 to 1 1/2 qts. per 100 gals. of water. Gal. \$4.20.

**KILPHENE 5-DUST** (Code No. I-3)—(Contains 5% toxaphene.) 1 lb. 10c; 25 lbs. \$1.70; 50 lbs. \$2.40; 100 lbs. \$4.80.

**KILPHENE 10-DUST.** (Code No. I-2)—(Contains 10% toxaphene.) Lb. 10c; 25 lbs. \$2.25; 50 lbs. \$3.45; 100 lbs. \$6.90.

**KILPHENE 5-SULPHUR DUST.** (Code No. I-1)—(Contains 5% toxaphene.) Lb. 10c; 25 lbs. \$1.95; 50 lbs. \$2.90; 100 lbs. \$5.80.

**VAPOTONE XX SPRAY (TEPP)** for control of aphids, thrips, red spiders. Use from 1/2 to 1 pt. per 100 gals. water. 4-oz. bottle \$1.00; 16-oz. bottle \$2.75; 1 gal. \$12.00; 5 gal. \$58.50.

**VOLCK OIL SPRAY.** A liquid oil spray and spray carrier, furnished in the following sizes only, 4 oz. 35c; 16 oz. 75c; 32 oz. \$1.25; 1 gal. \$3.00.





# FUNGICIDES

**ARASAN.** An effective treatment for vegetable seed, reduces loss from seed decay and damping-off.  $\frac{3}{4}$  oz. 25c; 8 oz. \$1.00; 2 lbs. \$3.20; 10-lb. bag \$11.50.

**CERESAN (New Improved).** Usually destroys seed-borne diseases by direct contact with the spores or penetrating vapor fumes. Ceresan has been used successfully in treating Gladiolus bulbs for control of Fusarium Rot before planting. Use  $\frac{1}{2}$  lb. Ceresan, 5 tablespoons of DuPont Spreader-Sticker to 25 gallons water. Dip bulbs in solution, let remain 15 minutes. Bulbs can be planted immediately after treatment. 1 lb. \$1.45; 100 lbs. \$93.50.

## COPPER COMPOUNDS

**BLUESTONE** (copper sulphate) Granular or Snowform. 1 to 10 lbs. 18c per lb.; 25 lbs. \$4.05; 50 lbs. \$7.00; 100 lbs. \$12.90.

**BORDEAUX** (powdered). Pound canister 50c; 4 lbs. \$1.35. 1 lb. makes 12 $\frac{1}{2}$  gals. spray.

## COPPER-A COMPOUND PREPARATIONS

**COPPER-A COMPOUND.** An excellent fungicide for spraying purposes, containing 45% metallic copper. Use 4 lbs. per 100 gals. of water. 6-lb. bag \$2.25; 50-lb. bag \$17.50.

**KILCOP DUST NO. 1** (Code No. A-1)—(5.4% metallic copper.) Lb. 10c; 25 lbs. \$2.30; 50 lbs. \$3.55; 100 lbs. \$7.10.

**KILCOP DUST NO. 2** (Code No. A-2)—(6.75% metallic copper.) Lb. 15c; 25 lbs. \$2.60; 50 lbs. \$4.15; 100 lbs. \$8.30.

**KILCOP-SULPHUR DUST NO. 3** (Code No. A-3)—(6.75% metallic copper.) Lb. 15c; 25 lbs. \$2.85; 50 lbs. \$4.70; 100 lbs. \$9.40.

**CUPROCID-YELLOW.** (Contains 80% metallic copper.) A cuprous oxide for the control of various truck crop diseases. Use 2 lbs. per 100 gals. of water. 3-lb. bag \$2.16; 100-lb. drum \$68.00.

## KILCOP "53" PREPARATIONS

**KILCOP "53."** (Contains 53% metallic copper.) Kilcop is extremely fine. This fineness is essential for good coverage and effective control of fungus. Use 4 lbs. per 100 gals. of water. 2-lb. bag 80c; 6-lb. bag \$2.05; 50-lb. bag \$16.03.

**KILCOP DUST NO. 5.** (Code No. O-2)—(7.4% metallic copper.) Lb. 10c; 25 lbs. \$2.20; 50 lbs. \$3.40; 100 lbs. \$6.80.

**KILCOP DUST NO. 6.** (Code No. O-3)—(9% metallic copper.) Lb. 15c; 25 lbs. \$2.40; 50 lbs. \$3.75; 100 lbs. \$7.50.

## DITHANE PREPARATIONS

**DITHANE D-14.** This fungicide has given excellent control on a variety of crop diseases such as late and early blight of potatoes and tomatoes, leaf blights of peppers, curvularia leaf spot of glads, anthracnose of beans and downy mildew of cucumbers, squash and melons. Harmful residue on foliage and fruit is negligible and yields of many crops are often higher than where other fungicides are used. Use 2 qts. Dithane D-14,  $\frac{3}{4}$  lb. zinc sulphate to 100 gals. water. 1 gal. \$2.10; 5 gals. \$9.75; 30 gals. \$52.50.



**DITHANE Z-78** (Wettable). Effective in the control of downy mildew and anthracnose of cucumbers, squash and melons and blue mold on tobacco. Use 2 lbs. to 100 gals. 1-lb. canister \$1.50; 3-lb. bag \$2.55; 100-lb. drum \$82.00.

**DITHANE M-22.** Controls a number of diseases which harm tomatoes, celery, peppers, eggplant and potatoes. Particularly effective where manganese deficiency exists. Use 2 lbs. per 100 gals. water. 3-lb. bag \$3.36; 25-lb. drum \$27.50.

**DITHANE DUST NO. 1** (Code No. M-1)—(Contains 4% zineb). Lb. 15c; 25 lbs. \$2.45; 50 lbs. \$3.90; 100 lbs. \$7.80.

**DITHANE DUST NO. 2** (Code No. M-2)—(Contains 6% zineb). Lb. 15c; 25 lbs. \$3.05; 50 lbs. \$5.05; 100 lbs. \$10.10.

**DITHANE-SULPHUR DUST NO. 4** (Code No. M-5)—(Contains 4% zineb). Lb. 15c; 25 lbs. \$2.75; 50 lbs. \$4.50; 100 lbs. \$9.00.

**FERMATE**, a fungicide for the control of tobacco blue mold, damping-off in celery seedbeds, downy mildew of cabbage, etc. Use 2 lbs. per 100 gals. of water. 3-lb. bag \$1.95; 50-lb. bag \$31.00.

**FERMATE DUST NO. 1** (Code No. F-1)—(Contains 11.4% ferbam). Lb. 20c; 25 lbs. \$3.70; 50 lbs. \$6.40; 10 lbs. \$12.80.

**MANZATE.** A new fungicide demonstrating good control of blights and related diseases. Use 1 $\frac{1}{2}$  lbs. per 100 gallons of water. 3-lb. bag \$3.36; 25-lb. drum \$27.50.

**ORTHOCIDE 50-W.** For control of damping-off in pepper seedbeds and other fungus diseases. For Irish potato seed treatment see page 52. Use 4 lbs. per 100 gals. water. 4-lb. bag \$3.44; 50-lb. drums \$41.50.

**PARZATE.** An effective control of blue mold of tobacco and anthracnose and downy mildew of cucumbers, squash and melons. Use 2 lbs. per 100 gals. of water. 3-lb. bag \$2.55; 50-lb. bag \$41.00.

**PARZATE LIQUID.** An effective control of blue mold of tobacco and anthracnose and downy mildew of cucumbers, squash and melons. Use 2 qts. plus  $\frac{3}{4}$  lb. Zinc Sulphate to 100 gals. water. 5-gal. can \$9.75; 30-gal. drum \$52.50.

**PHYGON-XL** (Wettable Grade). A fungicide for the control of leaf blights of tomatoes, beans and various other plant diseases. Use  $\frac{1}{2}$  to 1 lb. per 100 gals. of water. 2 oz. \$1.00; 1 lb. \$3.00; 4 lbs. \$6.60; 100 lbs. \$158.00.

**SEMESAN.** An effective treatment for vegetable and flower seeds to control damping-off fungi in the soil. Also repels ants. Directions on package. 4-oz. pkg. \$1.20; 12 ozs. \$3.40; 25 lbs. \$87.20.



# FUNGICIDES (Continued)

**SEMESAN BEL.** For treatment of potatoes, controlling certain diseases of this crop. Directions on package. 3 oz. \$1.05; 1 lb. \$3.00; 4 lbs. \$9.70; 40 lbs. \$90.75.

**SPERGON.** Very valuable for treating seeds, especially beans and peas, to control soil diseases for better stands and increased yields. Use 2 ounces per bushel of seed with beans and peas. 1 oz. 50c; 5 ozs. \$1.75; 1 lb. \$2.50; 10 lbs. \$20.00; 50 lbs. \$95.00.

**SPERGON (Wettable Grade).** Valuable for spraying to control diseases—especially downy mildew in cabbage seedbeds. Use at the rate of 3 lbs. to 100 gals. of water. 4 ozs. 50c; 1 lb. \$2.25; 10 lbs. \$11.40; 50 lbs. \$48.50.

**SPERGON DUST NO. 1. (Code No. Y-1)**—Contains 12% Spergon—Wettable. For the control of downy mildew on cabbage. Lb. 25c; 25 lbs. \$5.35; 50 lbs. \$9.65; 100 lbs. \$19.30.

**SULFOCIDE.** A vegetable spray which is also valuable for citrus fruit. Checks mildew, rust and anthracnose on beans. Also used in washing water before shipping beans, tomatoes, eggplants and peppers to check rots and other diseases while in transit. Use 2 qts. to 100 gals. of water. ½ pt. 65c; pt. \$1.00; qt. \$1.50; gal. \$3.00; 5 gals. \$11.50.

**SULPHUR (Dusting).** Lb. 5c; 25 lbs. 90c; 50 lbs. \$1.45; 100 lbs. \$2.85; ton \$56.50.

**SULPHUR (Wettable).** 50 lbs. \$1.55; 100 lbs. \$3.10; ton \$61.50.

**TERSAN 75.** A lawn and turf fungicide. Use 2 tablespoons full to 1½ gals. water to 60 sq. ft. 5 1/3-oz. cans \$1.00; 3-lb. bag \$6.75.

**ZERLATE.** Specific in the control of leaf blights. Use 2 lbs. to 100 gals. of water. 3-lb. bag \$1.95; 50-lb. bag \$31.00.

## Insecticide and Fungicide Combinations

Prices quoted are f.o.b. Plant City or Kilgore stores

Prices subject to change without notice



The Kilgore Seed Company processes many combinations of insecticides and fungicides which are too numerous to list in our General Catalog. If you do not find the combination you want, write our General Office in Plant City or call at your nearest Kilgore Store and it is likely we will have a dust in stock which will fill your requirements.

**DITHANE-KILGRAM DUST NO. 1 (Code No. MQ-1)**—(Contains 1% lindane and 4% Zineb.) This combination is recommended for purposes where Dithane Z-78 has been proved more effective than copper. Refer to Kilgam-Kilcop Dust No. 1 for further important details. 1 lb. 20c; 25 lbs. \$4.50; 50 lbs. \$7.75; 100 lbs. \$15.50.

**DITHANE-KRYOCIDE DUST. (Code No. MK-1)**—(Contains 4% Zineb and 30% Kryocide.) This combination is recommended for purposes where Dithane Z-78 has been proved more effective than copper. Lb. 20c; 25 lbs. \$3.85; 50 lbs. \$6.65; 100 lbs. \$13.30.

**KILCIDE-DITHANE DUST NO. 1 (Code No. DM-1)**—(Contains 5% DDT and 4% Zineb.) For use where DDT is desirable for general insect control with Dithane Z-78. 1 lb. 15c; 25 lbs. \$3.05; 50 lbs. \$5.05; 100 lbs. \$10.10.

**KILCIDE-KILCOP DUST NO. 1 (Code No. DA-1)**—(Contains 3% DDT plus 5.4% metallic copper.) Use where DDT is needed with a neutral copper. Lb. 15c; 25 lbs. \$2.60; 50 lbs. \$4.20; 100 lbs. \$8.40.

**KILPHOS-DITHANE NO. 1. (Code No. EM-1)**—(Contains 1% parathion and 6% Dithane.) Can be used on most crops for most insect and disease control. 50 lbs. \$6.85; 100 lbs. \$13.70.

**KILGAM-KILCOP DUST NO. 1 (Code No. QA-3)**—(Contains 1% lindane (essentially pure gamma isomers of benzene hexachloride) and 5.4% metallic copper.) An excellent combination designed for use on cucumbers, squash, and melons which cannot tolerate ordinary BHC compounds (Kilhex), as well as for cabbage, celery, tomatoes, peppers and eggplants at periods when dusts containing ordinary BHC (Kilhex) cannot be used because of the possible contamination of edible portions due to odor or flavor. 1 lb. 20c; 25 lbs. \$4.20; 50 lbs. \$7.40; 100 lbs. \$14.80.

**KILHEX-KILCOP DUST NO. 1 (Code No. GA-1)**—(Contains 1% gamma isomer and 6.75% metallic copper.) A practical insect and disease control combination at a minimum cost for cabbage, celery, tomatoes, peppers and eggplants for application before fruits or edible portions are well developed. 1 lb. 15c; 25 lbs. \$3.15; 50 lbs. \$5.25; 100 lbs. \$10.50.

**KILPHENE-KILCOP DUST NO. 2. (Code No. IA-2)**—A convenient insect and disease control combination containing 5% toxaphene and 6.75% metallic copper. (Under no circumstances use this material on cucumbers, squash, or melons.) 1 lb. 15c; 25 lbs. \$3.15; 50 lbs. \$5.25; 100 lbs. \$10.50.

**KILCIDE-KILCOP-MANGANESE-SULPHUR DUST NO. 1. (Code DAX-1)**—(Containing 5% DDT, 5.4% metallic copper, manganese sulphate and sulphur) a combination dust for control of insects, disease and nutritional deficiencies. 1 lb. 15c; 25 lbs. \$3.15; 50 lbs. \$5.30; 100 lbs. \$10.60.

**KRYOCIDE-KILCOP DUST NO. 3. (Code No. KA-1)**—(Contains 30% Kryocide plus 5.4% metallic copper.) A good insecticide and fungicide combination for most vegetable crops. Lb. 20c; 25 lbs. \$3.65; 50 lbs. \$6.30; 100 lbs. \$12.60.



# HOUSEHOLD AND GARDEN PESTICIDES

**BLACK LEAF 40.** For the control of aphids, thrips. Use 1 qt. in 100 gallons water. 1 oz. 39c; 5 oz. \$1.19; 1 lb. \$2.69; 2 lbs. \$4.79. Prices quoted are prepaid.

**KILGORE'S GENERAL PURPOSE DUST.** (Code No. P-6.) Contains pyrethrins. A non-poisonous dust, which can be used in or around the house to kill ants, roaches, fleas and many garden insects. The ideal dust for use on garden vegetables up to picking time, Dust Gun is refillable and can be used over and over provided the gun is kept in a dry place when not in use. Dust gun complete with dust 75c; refills for dust gun (dust only), 1 lb. 30c.

**IMPROVED ISOTOX GARDEN SPRAY-M**—An effective spray containing 5% lindane, for control of insects on roses, camellias, azaleas, ixoras, hibiscus, bougainvillea, and other ornamentals, house and garden plants. This spray is marked by the absence of odor and visible residue. Use four teaspoons per gallon of water. Price: 4-oz. bottle \$1.00; 8-oz. \$1.75; 16-oz. \$2.95.

**KILCHLOR 40-W.** A chlordane concentrate for the control of chinch bugs, army worms, sod worms and ants by spraying. 1-lb. can 75c; 5-lb. bag \$3.00; 10-lb. bag \$6.00; 25-lb. bag \$13.75; 50-lb. bag \$25.00.

**KILCOP "45"**—(Contains 45% metallic copper.) Kilcop is extremely fine which is essential for good coverage and effective control of fungi. Use 4 lbs. per 100 gals. of water. 1-lb. can 70c.

**C-M-Z GARDEN SPRAY.**—A combined nutritional and fungicidal spray for shrubs, fruit trees and vegetables. Use two tablespoons per 1 gallon of water on Avocados, Mangos and shrubs, 1¼ tablespoons on citrus and 2½ tablespoons on vegetables. No wetting agent required. 1 lb. can 75c.




**KILGORE'S SHRUB DUST.** (Code No. C-1.) Safe and effective material for the control of a wide variety of insects on shrubs and other ornamentals. Kills aphids, thrips, practically all types of caterpillars, grasshoppers, and many other chewing and sucking types of insects. Not effective against scale insects, mealy bugs, white flies and red spiders. Much less toxic to human beings and warm blooded animals than most insecticides. Dust gun complete with dust 75c. Refills for dust gun (dust only) 1-lb. can 40c; 5-lb. bag \$1.30.

**DRY FLY BAITS.**—For use against house flies and their maggots in out-buildings, kennels, around garbage cans, etc. Easy to apply—scatter by hand. No spraying equipment needed. Kills within a few minutes. Lasts for several days when applied to dry ground. Do not use inside of dwellings.

**FLY FLAKES**—Apply 4 oz. per 100 sq. ft. 1 lb. shaker can \$1.15; 5-lb. bag \$1.50; 10 lbs. \$2.75; 25-lb. drum \$6.25.

**ORTHO FLY KILLER (Dry Bait)**—Apply 4 oz. to 500 sq. ft. 1 lb. can \$1.25; 5-lb. bag \$1.98.

**KILGORE'S KIL-CHINCH.** Granulated. (Code No. V-2.) A recently developed material for satisfactory chinch bug control in St. Augustine type of grasses. Furnished in a granular form, similar to coarse sea sand. The toxic ingredient is malathion. Apply at the rate of two pounds per 100 sq. ft. 5-lb. bag \$1.50; 10-lb. bag \$2.50; 50-lb. bag \$10.00; 100-lb. bag \$20.00.



**ORTHO LAWN CHINCH BUG SPRAY.** Controls chinch bugs, cutworms, sod webworms, army worms and ants. One qt. per 1,000 sq. ft. in sufficient water to cover area uniformly. 16 oz. \$1.95; 32 oz. \$3.50.

**KILCHLOR.** A residual household spray. Contains 2% chlordane plus isobornyl thiocyno-acetate, for control of roaches, ants, and most other household insects. Pt. 40c; qt. 65c; gal. \$2.15.

**KILSPRAY.** An excellent indoor space spray for flies, mosquitoes, roaches, ants, fleas, and certain other household insects. Not a residual spray. It should be used as necessary whenever insects are present. Qt. 65c; gal. \$2.15.

**MALATHION Emulsion 50%.** For control of scale and many other lawn and garden insects. Use 2 tablespoons per gal. of water. 1 oz. 39c; 5 oz. \$1.19; pt. \$2.69; qt. \$4.79.

**NICOTINE PYROX.** Kills leaf-chewing insects, thrips, leaf-hoppers, and prevents blight and other diseases. Designed for commercial crops as well as home gardens. 20-oz. jar 60c; 5 lbs. \$2.25; 25 lbs. \$8.50; 50 lbs. \$13.50.

**KILCHLOR-KILPHENE POISON BAIT NO. 11.** (Code No. B-11)—(Contains chlordane and toxaphene). For control of mole crickets, cutworms, and many other soil insects. 1 lb. 15c; 25 lbs. \$2.30; 50 lbs. \$4.15; 100 lbs. \$8.30.

**ORTHO-KLOR "44" SPRAY.**—For control of ants, mole crickets, cutworms, army worms, roaches and many other pests. For Termite Control see page 58. 4 oz. 90c; 8 oz. \$1.45; 16 oz. \$2.45; 32 oz. \$4.25.

**ORTHO-KLOR-8 SPRAY (Containing 72% chlordane emulsion).** For Termite Control see page 58. 1 gal. \$10.75; 5 gals. \$52.50; 30 gals. \$307.50.

**ORTHOL GARDEN SPRAY.** An oil emulsion containing nicotine and DDT for the combined control of aphids, red spiders, mealy bugs, thrips and many other insects. Use at the rate of 6 tablespoonfuls to 1 gal. of water. 4 oz. 35c; 16 oz. 75c; 32 oz. \$1.25; 1 gal. \$2.85.

**RED ARROW INSECT SPRAY.** Kills garden aphids, melon aphids, worms, leaf-hoppers, white flies, red spiders, caterpillars, thrips, celery leaf tier. 1 oz. 35c; 4 oz. \$1.00; pint \$2.85; quart \$5.00; gal. \$18.20. Average dilution 1 to 400 parts water.

**KILGORE'S ROSE DUST.** (Code No. PO-4.) A combined insecticide and fungicide, to keep rose bushes free of insects and diseases, by applying at intervals of seven days. An excellent material as a general insecticide and fungicide for garden vegetables. Dust Gun, complete with dust 75c. Refills for dust gun (dust only) 1 lb. 35c.

**SABA CHINCH.** (Code No. S-2.) For control of chinch bugs, army worms, sod webworms, and ants. Use 4 lbs. per 100 sq. ft. 1 lb. 15c; 25 lbs. \$3.10; 50 lbs. \$5.15; 100 lbs. \$10.30.

**SEEDTONE.** Seed Protectant. (Containing 50% captan)—A guard against seed decay, damping off and seedling blight. ¼ oz. pkt. 25c.

**VOLCK OIL SPRAY.** 4 oz. 35c; 16 oz. 70c; 32 oz. \$1.25; 1 gal. \$3.00.

**VOLCK ISOTOX SPRAY.** For use on ornamental plants against red spiders, certain scale insects, mealy bugs, white flies and certain other garden pests. Use 5 tablespoonfuls per 1 gal. of water. Don't use more than three applications per season. 4 oz. 40c; 8 oz. 60c; 16 oz. 95c; 32 oz. \$1.75.





# FUMIGANTS


**CARBON DI-SULPHIDE (Hi-life.)** Recommended for ridding grain of weevils, and killing ants, etc., and for destroying vermin. Sold in gallon can only. Per gallon \$2.00. (Not mailable.)

## Pyrenone

\*Reg. U. S. Pat. Off.

### GRAIN PROTECTANT

25-lb. bag \$6.00; 50-lb. bag \$11.50, f.o.b. Plant City.




(Not mailable.)

**CYANOGAS A-DUST.** For killing the following pests: ants (Garden, Mound Building; Leaf Cutting). Rats (in burrows; along ditch banks; in city dumps; under pavements and tight floors; in lumber piles; on poultry farms; in grain elevators and stockyards). Mice (field mice and pine mice), moles, gophers. 1-lb. tin \$1.25; 5-lb. can \$4.25; 25-lb. can \$13.75.

**CYANOGAS G-FUMIGANT.** Comparable to sea sand in size of particles. Used for fumigating greenhouses, warehouses, buildings and for grain fumigation. 4-oz. can 50c; 5-lb. can \$4.25; 25-lb. can \$13.75; 100-lb. drum \$43.35. (Not mailable.)

**CYANOGAS ANT-KILLER.** Cyanogas Ant-Killer kills ants with the least trouble. It is a gas-producing powder—flows freely through the spout on the can. Not a bait. It's the gas that kills 'em. 4-oz. can 50c. (Not mailable.)



**SOIL FUMIGATION FOR CONTROL OF SOIL-BORNE PESTS,** root knot nematodes, other nematodes, and wire worms.

Nematodes are present in most of our soils, and seriously affect most of our vegetable crops and reduce yields. Nematodes can be controlled by the proper use of soil fumigants, either solid treatment, which is applied with power applicator about 12 inches apart, or row treatment, which is only applied in the row that is to be planted. This method has become very popular, especially with small growers, as cheap handmade applicators with gravity flow, can be used attached on a tractor or a horse drawn plow. The cost of this method of treatment is only about one-third the cost of solid treatment, and has proved very satisfactory. "D-D" and "Soilfume" are the two materials most extensively used in commercial treatment at this time. Information on soil fumigation can be obtained from any of the Kilgore stores or write The Kilgore Seed Company, Plant City, Florida.

**D-D Soil Fumigant.** 1 gal. will treat approximately 1500 square feet of soil area applied in 15 inch squares, 6 inches deep, with hand applicator. 22 gals. will treat approximately one acre applied with a machine in 12 inch rows. Applied only in the row that is to be planted. 1 pt. will treat 260 feet of row, or approximately 5½ gals. per acre in 4 foot rows. Gallon \$2.95; 54 gals. \$91.80.

**SOILFUME 80-20 (ethylene dibromide).** 1 gal. will treat approximately 1400 square feet of soil area applied in 12-inch squares, 10 inches deep with hand applicator. 30 gals. in 12-inch rows applied with a machine will cover one acre. Applied only in the row that is to be planted, 1 pt. will treat approximately 140 ft. of row or 10 gals. per acre in four foot rows. 1 gal. \$1.65; 30-gal. drums \$46.50; 50-gal. drums \$75.00.

**SOILFUME CAPS (ethylene dibromide).** The use of these capsules for nematode control offers an extremely convenient and easy method for soil fumigation in small areas. Capsules are placed in the soil six inches deep and twelve inches apart. Seed or plants should not be put in under two weeks after treatment. Areas immediately adjacent to other plants or grasses should not be treated.

100 capsules \$1.00



Soil fumigants should be applied 6 inches deep in the soil and well covered. Soil should be in good condition and all trash and cover crops decomposed or removed. Moisture should be about right for planting. Wait two weeks after fumigating before planting.

Do not fumigate wet soil. If it rains and stays wet immediately after fumigating, wait three weeks before planting.

Observe all cautions and instructions on the labels in regard to the use and handling of Cyanogas and soil fumigants.

## Spreaders, Stickers, Emulsifiers, Etc.

**DUPONT SPREADER-STICKER.** A combined spreading and sticking agent designed for use with agricultural sprays. Compatible with hard water. Increases the efficiency of weed-killing compounds such as Ammate. 1 pt. \$1.25; 1 gal. \$8.60; 5 gals. \$42.25.

**TRITON B-1956.** An efficient emulsifier and spreader for insecticides and fungicides. A combined spreader, sticker and deposit builder. Soluble in water and oil,

it is widely used as an emulsifier in fruit, truck crop and ornamental sprays for maximum spreading and sticking qualities. Use at the rate of 2 to 4 ounces per 100 gallons. Gal. \$7.70; 5 gals. \$37.75; 50 gals. \$375.00.

**FISH OIL SOAP.** Increases the efficiency of Black Leaf 40 and many other spray materials. Use 1½ to 2 lbs. to 50 gallons as a spreader and sticker. 1 lb. 40c; 1-gal. can \$2.35; 5-gal. can \$8.35.



# RODENTICIDES AND VERMICIDES

**CARBOLINEUM.** A wood preservative and vermicide. Prevents damage by termites. Kills and protects against chicken mites and fowl ticks. ½-gal. can \$1.65; 1 gal. \$2.50; 5 gals. \$10.50; 35 gals. \$59.50; 55 gals. \$88.00, f.o.b. Plant City.

**NO NIB'L RABBIT REPELLENT** is recommended as a rabbit repellent for flowers, shrubs and other ornamentals. When used as directed, will not harm plants or soil and is non-injurious to birds, rabbits and other animals. It is easy to use and is effective either as a dust or as a spray. 6 oz. shaker-top can 98c.

**DIXIE KILLER.** Exterminates rats, mice, roaches, waterbugs, land crabs, cutworms and other destroyers of food and property. 2-oz. jar 35c; 1 lb. 90c; 2 lbs. \$1.50. Prepaid.

**MAGIKIL JELLY ANT BAIT.** Kills ants and is harmless to plants. Ant trap 25c; Garden size 40c; Estate tube \$1.00; 7½ oz. Jumbo size \$1.95; 10 lbs. \$23.00. Prepaid.

**TERRO ANT KILLER.** Will rid your place of ants quickly. 1¼-oz. bottle 35c; 3-oz. bottle 50c. Prepaid.

**GATOR ROACH HIVES** are sanitary, open-end fibre tubes, containing a specially made gum that roaches, silverfish, waterbugs and crickets eat with relish but results in their certain death. Small size (3 baits) 20c; Large size (3 baits) 40c. Prepaid.



**HARRIS ROACH TABLETS.** 2-oz. pkg. 25c; 4-oz. pkg. 50c.

**COLONIAL-42.**—An effective bait containing Warfarin for the control of all species of rats and mice. 4-oz. pkg. 30c; 10 oz. 60c; 1 lb. 85c; 2 lbs. \$1.50.

**KILGORE'S RAT AND MOUSE BAIT.** A Warfarin bait especially designed for the control of rats and mice in fields and groves. 10 lbs. \$3.50; 25 lbs. \$8.00; 50 lbs. \$15.00.

**QUICK DEATH RAT KILLER.** A fortified red squill formula. Safe to use in homes. Price, 25c per package.

## WEED KILLERS

*Note: Read labels carefully—they are put on the package for your protection. Use Weed Killers wisely and you will be using them safely. Be sure to clean spray equipment thoroughly according to approved methods after using weed control materials of any kind. f.o.b. Plant City.*



**AMMATE WEED KILLER.** Ammate can be used for the control of many types of noxious weeds and generally effects a permanent kill. It is not considered poisonous and may be used safely on weed-infested areas without danger to livestock and humans. Ammate usually exerts only temporary soil sterilizing effects, varying with the soil. Treated areas may often be used for growing crops during the same season. 2-lb. bottle \$1.15; 6 lbs. \$2.95; 50-lb. drum \$15.25. (Use 1 to 3 lbs. per 100 sq. ft. area.)

**"KARMEX" W WEED KILLER (Formerly DuPont's CMU).** Kills weeds and grasses. Prevents regrowth for prolonged periods of time. Recommended for use in parking areas, roadways, storage yards, warehouses, fence rows, drainage ditches, etc. Do not apply on or near valuable woody or herbaceous plants or on areas into which their roots may grow. Do not apply to locations (like driveways) from which it may be washed onto areas where roots of desirable plants may come in contact with the herbicide. Apply at rate of from 20 to 40 lbs. per acre. (1 to 2 pounds per 1,000 sq. ft.) Ask for descriptive folder.


2 lbs. \$8.50; 10 lbs. \$39.00; 50 lbs. \$187.50

**POLYBOR-CHLORATE.** A non-selective herbicide for the prolonged control of weeds about brick walks and patios, fence lines, driveways, or tennis courts, around stores, industrial properties, etc. It is non-selective in its action and will kill any plant with which it comes in contact, consequently cautions, as per label instructions should be observed in order to protect valuable shrubs and trees adjacent to the areas to be treated. Dissolved at the rate of four pounds in one or two gallons of water to be applied to a 100 sq. ft. area. Price 5-lb. pkg. \$2.25; 50-lb. bag \$10.25.

**WEED-B-GON—64X.** An amine salt of 2,4-D. Excellent for killing water hyacinth and most broad leaf

plants. Apply as per directions on container. Caution—do not use on or let material drift on plants you do not want to kill. Price, 1 gal. \$4.80; 5 gals. \$22.90; 30 gals. \$130.50; 55 gals. \$235.40.

**TCA WEED KILLER (DuPont's).** TCA is a grass killer for non-selective control in areas not planted to crops. For maximum control of Para grass along ditch banks it is suggested to first use a "pre-weakening" method, such as treating with a relatively cheap contact herbicide, or by flame singeing. TCA can then be applied at fairly low dosages. As an alternate, several applications of TCA in small amounts will produce better results than one application at a higher dosage. For more specific information write The Kilgore Seed Company Laboratory, Box 6085, West Palm Beach, Fla. 10 lbs. \$5.20; 50 lbs. \$22.50; 100 lbs. \$44.50.



**WEEDONE BRUSH KILLER—32.** Contains 2,4,5-T, plus 2,4-D. Kills poison ivy, poison oak stumps and other woody plants. Use 1 to 1½ gals. to 100 gals. water or ½ to 1 pt. to 5 gals. water. Caution—do not use this material on plants you do not want to kill. Price qt. \$3.75; gal. \$7.95; 5 gals. \$37.75.



**NEW IMPROVED WEEDONE.** A selective weed killer containing 2,4-D plus 2,4,5-T. For complete kill, all the foliage must be sprayed as there is little cross-movement within the stems. Weedone is effective only when the leaves are green and actively growing, for the chemical does not move through dormant or dead tissue. Do not use Weedone on St. Augustine type grasses. 8-oz. bottle \$1.00; 1-qt. bottle \$2.75; 1-gal. bottle \$6.75; 5 gals. \$24.75.

*A number of pre-emergence and post-emergence weed control materials are now being offered to truck crop growers. As most of these materials have not been tried sufficiently under Florida conditions we hesitate to make specific recommendations in this catalog. We suggest that growers interested should consult local authorities.*

**WE WILL MAKE EVERY EFFORT TO SUPPLY WEED CONTROL MATERIALS OF THIS NATURE RECOMMENDED BY THE FLORIDA EXPERIMENT STATIONS FOR EACH AREA.**



# LIVESTOCK PREPARATIONS

Prices are subject to change without notice and are f.o.b. Plant City unless otherwise specified.

**DDT 50% Wettable.** For spray on range cattle. To control horn flies, lice and mosquitoes use 20 lbs. per 100 gals. water. Do not use on dairy cattle. 1-lb. canister 70c; 3-lb. bag 90c; 50-lb. bag \$13.50.



**Gamtox Wettable No. 10** (10% Gamma isomer of BHC). For spray on range cattle to control horn flies, lice and mosquitoes use 3¾ lbs. per 100 gal. water. Do not use on dairy cattle. 5-lb. bag \$1.60; 50 lbs. \$14.50.

**Marlate Methoxychlor** (50% Wettable). For spray on dairy and range cattle to control flies and lice. For lice use 25 lbs. per 100 gal. water. For flies use 8¾ lbs. per 100 gal. water. 4-lb. bag \$3.40; 50-lb. bag \$40.00.

**DuPont Dairy Cattle Spray.** (Methoxychlor, 50% wettable). For control of flies on dairy cattle use 1 lb. in 12 gals. water. 1 lb. \$1.25; 2 lbs. \$2.20.

**DuPont Livestock Spray and Dip No. 30.** For control of horn flies, lice and mosquitoes on range cattle. Do not use on dairy cattle. 2-lb. bag \$1.05; 11-lb. bag \$5.75; 55-lb. bag \$25.50.

**Isotox Dairy Spray.** (Contains lindane). Spray for killing flies in dairy barns and milk sheds. Do not spray on dairy cows. 1 lb. \$2.65; 4 lbs. \$10.40.



**Ortho Fly Killer.** You get a quick kill of house flies in dairy barns, hog pens, poultry houses and other places that attract flies. Mix one 4-oz. bottle of Ortho Fly Killer plus ½ pt. of syrup in three gallons of water. Apply with a regular sprinkling can on barn floors, walkways, etc.

Do not apply Ortho Fly Killer directly on animals or feed. Follow label instructions. 4-oz. bottle \$1.25; Full case of 24 bottles \$27.50.

## DRY FLY BAITS

For use against flies and their maggots in dairy and livestock barns, stables, poultry houses, feed rooms, kennels, manure piles, etc. Do not apply where poultry can feed upon the poison.

**Fly Flakes.** For adult flies apply 4 oz. per 100 sq. ft. For maggots, use 1½ lbs. per 100 sq. ft. 1-lb. shaker-top can \$1.15; 5-lb. bag \$1.50; 10-lb. bag \$2.75; 25-lb. bag \$6.25.

**Ortho Fly Killer Dry Bait.** For adult flies use 4 oz. to 500 sq. ft. 1-lb. canister \$1.25; 5-lb. bag \$1.98.



**ORTHO Healthy Herd.** A wettable powder. (Contains 45% DDT and 2.5% gamma isomer of BHC and 12.5% other isomers of BHC.) Use as spray or dip on cattle, horses, hogs, sheep, and goats to control lice, ticks, horn flies and mosquitoes. Use 10 lbs. per 100 gals. water. (Not for use on dairy cattle or in dairy barns.) 5-lb. bag \$3.25; 50-lb. bag \$25.50.

**ORTHO KLEEN Stock.** An emulsifiable liquid containing toxaphene and lindane for the control of ticks, lice, flies, etc. on beef cattle and hogs. One pt. makes 4 to 12½ gals. finished spray depending on use. Directions on container. (Not for use in dairy barns or on dairy cattle.) Price 16-oz. bottle \$1.40; 32-oz. bottle \$2.40; 1 gal. \$6.95; 5 gals. \$27.50.



**Smearex—(Screw Worm Preparation).** ½ pt. 50c; pt. 90c; qt. \$1.75; gal. \$6.00.

**Ortho 1038 Screw Worm Control.** 4-oz. bottle 90c; 16 oz. \$2.75; 32 oz. \$5.00; 1 gal. \$18.00.

**Ortho EQ 335 Screw Worm Remedy.** 4-oz. bottle 70c; 16 oz. \$2.00; 32 oz. \$3.50.



*Note: Follow label directions on all screw worm preparations.*

## BIG STINKY OUTDOOR FLY TRAPS

The action of Big Stinky's Control Fluid with water on the starting bait aids in producing a potent scent which lures flies in the vicinity into the trap. Big Stinky actually lures flies away from houses, barns, stock pens and other buildings where you don't want them—it's irresistible—to filthy flies.

|   |        |
|---|--------|
| 1-gal. size with 8 oz. control fluid .....      | \$4.95 |
| "Little Stinky" with 3 oz. control powder ..... | 2.95   |
| "Wee Stinky" with 1½ oz. control powder .....   | 1.69   |
| 8-oz. bottle control fluid .....                | 1.50   |
| 3-oz. tube control powder .....                 | .89    |

## POULTRY PREPARATIONS

**Walko Tablets.** Use in drinking water for baby chicks. Aids in treating cold in head, simple catarrh, sneezing, wheezing, and roup. Package 60c.

**Tilghman Condition Powder.** Aromatic stimulant and carminative for cattle, horses, sheep, hogs, and chickens. Package 25c. Postpaid.

**Germozone.** For use in drinking water. Valuable as preventive of simple bowel trouble, also ocular roup. 4 fluid oz. 50c; 12 fluid oz. 90c.

**Kilcide 10 Powder.** (Containing 10% DDT.) For treating chickens against lice. Apply by dusting it well into the feathers in order to get the dust down to, or near, the skin surface. 1 lb. 15c; 25 lbs. \$2.35.



# FERTILIZERS

Kilgore's stores carry a complete line of Fertilizers and Fertilizer Material for Truck Crops, and Groves, that is not listed in this catalog or sold through mail order. All prices quoted subject to change without notice.



**VIGORO**—Complete balanced plant food supplies the nutrient elements plants must get from the soil for best growth and development. Vigoro is odorless, sanitary and easy to apply. Made for Florida soils and conditions. Use it on everything you grow. Vigoro 6-9-6; 1 lb. 15c; 5 lbs. 55c; 10 lbs. 95c; 25 lbs. \$1.70; 50 lbs. \$2.55; 100 lbs. \$4.85 f.o.b. Plant City or any Kilgore store. Write for prices on larger quantities.


## CAMELLIA AND AZALEA PLANT FOOD

A special prepared fertilizer for camellias and azaleas. Containing the three primary plant foods, plus secondary elements. Price 5 lbs. 58c; 10 lbs. \$1.05; 25 lbs. \$1.90; 100 lbs. \$5.75. f.o.b. Plant City.

## SOLUBLE PLANT FOODS

**HY-GRO.** A soluble plant food containing the three major plant food elements, nitrogen, phosphoric acid and potash, plus minor elements.

An excellent starter solution for potted plants, seedbeds and transplanting. Use at the rate of two tablespoonfuls dissolved in one gallon water or 1 teaspoonful to quart. Lawns can be easily watered and fertilized at the same time by attaching a Hy-Gro Gat sprayer (see page 71) to your garden hose.



Hy-Gro can be used on growing and fruiting plants as a plant stimulant spray, or in combination with insecticide and fungicide sprays, at the rate of 4 lbs. to 100 gallons. It prolongs the life of the plant, increases the size of the fruit and prolongs the picking period. Hy-Gro is used very extensively by large commercial vegetable growers.

3 oz. 30c; 1 lb. 90c; 2 lbs. \$1.45; 5-lb. bag \$2.15; 10-lb. bag \$4.00; 100 lbs. \$25.00, f.o.b. Plant City.

**GRO-CROP.** A liquid plant food containing nitrogen, phosphorus and potash, plus minor elements needed to produce healthy plant growth. Used as ground application, starter solution and foliage spray.

Compatible with most insecticides and fungicides spray. Excellent for airplane spraying. It is a liquid and won't clog nozzles.

Use 2 tablespoonfuls per gallon water, or 2 qts. per 100 gallons.

8 oz. 59c; 16 oz. 98c; 32 oz. \$1.79; 1 gal. \$4.25; 30-gal. drums \$97.20.

**NUTRI-LEAF.** A soluble fertilizer containing 16% nitrogen, 16% phosphoric acid and 16% potash plus minor elements. Compatible with most insecticides and fungicides as a foliage spray for plant stimulant. Use 6 to 8 lbs. per 100 gal. water. 3-lb. bag 84c.



**VERTAGREEN** is a high quality, scientifically blended plant food, for lawns, flowers, shrubs, trees, vegetables. Contains nitrogen, phosphorus and potash, besides several other important plant food elements. Clean, odorless, easy to apply. Used with outstanding success.

Price per lb. 15c; 5 lbs. 55c; 10 lbs. 95c; 25 lbs. \$1.70; 50 lbs. \$2.55; 100 lbs. \$4.85. f.o.b. Plant City, or Kilgore Stores.



## PULVERIZED SHEEP MANURE

Sheep manure is highly concentrated, with all weed seeds killed. It supplies the humus and natural plant-food so greatly needed by Florida soils.

**For New Lawns** use 10 to 12 pounds per 100 square feet and rake into the surface soil. On old lawns use 8 to 10 pounds scattered broadcast and wet down with a hose.

**For Flowers and Shrubs** mix sheep manure with the soil around each plant, using 1 to 5 pounds according to size.

**For Fruits and Vegetables** use 1000 to 1500 pounds per acre. Apply with a drill in the row or hill. It is a profitable and lasting soil-building fertilizer. 25 lbs. 1.00; 50 lbs. \$1.75; 100 lbs. \$3.50; Ton \$67.50, f.o.b. Plant City.

**VHPF "VIP"**—A soluble fertilizer—containing the three major plant foods, nitrogen, phosphoric acid and potash plus essential minor elements. Used as a starter solution, a nutritional spray, proved very successful in combination with fungicide and insecticide sprays on growing and fruiting plants as a stimulant. Use 1 lb. to 20 gals. or 6 lbs. to 100 gals. water.

Price: 1-lb. can 50c; 3-lb. bag 75c; 30-lb. bag \$6.90, f.o.b. Plant City or Kilgore stores.



**HYPONEX**—Valuable in seedbeds and to start plants after transplanting. Comes in a powder which makes liquid plant food containing all plant food elements plus vitamin B<sup>1</sup>. Dilute 1 oz. in 6 gals. water. 1 oz. 10c; 3 oz. 25c; 7 oz. 50c; 1 lb. \$1.00; 10 lbs. \$8.00, f.o.b. Plant City.

**NUGREEN.** A soluble plant food containing 45% nitrogen in the form of urea. Gives excellent results when quick acting nitrogen is needed. Can be applied as a foliage spray in combination with insecticides and fungicides at the rate of 4 to 5 lbs. per 100 gals. Can also be applied through irrigation or as top dresser. 80-lb. bag \$6.75; ton \$168.75.

**ORTHO-GRO.** A liquid plant food for potted plants, flowers, lawns and gardens. 4-oz. bottle 35c; 16 oz. 90c; 32 oz. \$1.50; 1 gal. \$4.95.


**PLANTABBS FOR HOUSE PLANTS.** Plantabbs are a complete, balanced plant food of the high analysis 11-15-20. Use Plantabbs for all indoor potted plants and flowers, window and balcony boxes. Also for garden flowers and vegetables. Plantabbs with vitamin B<sup>1</sup> make vastly stronger root growth.

10 Tablets 10c; 30, 25c; 75 50c; 200, \$1.00; 1000, \$3.50; 500 in red plastic pot, \$2.00.





# NUTRIENTS



**ACTIVO.** A compost activator. Converts compost quickly and easily into available plant food.

No. 2 Size for 350 lbs. of Compost .....\$1.00  
No. 7 Size for 1200 lbs. of Compost ..... 2.95

**ADCO.**—A clean odorless powder for converting leaves, grass clippings etc. into compost. 5 lbs. makes 200 lbs. organic compost. 5-lb. bag \$1.15; 20-lb. bag \$2.85.

**ES-MIN-EL.** For soil application. Contains many essential mineral elements in proper proportions. Add to ordinary fertilizers at rate of 1 lb. or 2 lbs. to 10 lbs. regular fertilizer or 100 to 200 lbs. per ton of regular fertilizer. 5-lb. bag \$1.00; 100-lb. bag \$10.00; Ton \$190.00, f.o.b. Plant City.


**ES-MIN-EL.** For spray or dust. For spray use 4 lbs. to 100 gals. water. For dust use 16 lbs. Es-Min-El mixed in 84 lbs. of talc, clay or other inert ingredients. Price: 4-lb. bag \$1.50; 50-lb. bag \$13.00, f.o.b. Plant City.

**MANGANESE SULPHATE (70% spray grade).** A quick acting material for applying manganese to plants in the form of a spray when plants are stunted and turning yellow on very alkaline or sweet soils. Apply at the rate of 4 lbs. to 50 gallons of water. 1 lb. 10c; 50-lb. bag \$2.60, f.o.b. Plant City.

**KILGORE'S C-M-Z Spray.** A combined nutritional and fungicidal spray for tomatoes, potatoes, peppers and other vegetables. Do not use on sweet corn. Apply at the rate of 6 lbs. per 100 gals. of water. 6-lb. bag \$1.85.

**SULPHUR-MANGANESE DUST—10%.** (Code No. X-2) —For manganese deficiencies in beans, peppers, etc. 1 lb. 10c; 25 lbs. \$1.50; 50 lbs. \$2.00; 100 lbs. \$4.00.

**KILGORE'S IRON-AID.** (Contains a newly developed chelated iron compound, which is completely available to plants under certain soil conditions)—This is a material designed for the correction of iron chlorosis wherever symptoms are indicated in flowers or shrubs. As the symptoms of iron deficiency vary widely in various types of plants, it is often difficult to decide whether the chlorotic condition is due to an iron deficiency or not. In a general way these symptoms indicate a tendency of paling the areas between leaf veins, while the veins themselves still retain their original green color. In severe cases or the final stages of iron chlorosis, such areas show an almost pure white color. As some other mineral deficiencies show similar symptoms it is rather difficult to determine the actual reason for the chlorotic condition. As KILGORE'S IRON-AID is a relatively inexpensive material we suggest the use of it in all cases where iron chlorosis is indicated as well as in questionable borderline cases. Consult label for specific directions before applying this material to shrubs, flowers or lawns.



1 lb. 60c; 5 lbs. \$1.25; 10 lbs. \$2.00; 25 lbs. \$3.75; 50 lbs. \$5.90; 100 lbs. \$11.80.

**ALUMINUM SULPHATE.** For increasing acidity of soils where aluminum toxicity is no factor. 10c per lb.; 100 lbs. \$4.00.

**LIME, HYDRATED.** Lime is a wonderful land-sweetener. Use at the rate of a half ton to a ton per acre, broadcast, before breaking the ground. If soil is new and sour, the application should be heavier. You can use hydrated with safety. Also used for the neutralization of manganese, zinc and copper sulphates. 10-lb. bag 30c; 50-lb. bag 85c; ton \$30.00; ½ ton at ton rate, f.o.b. Plant City.

**ZINC SULPHATE.** For zinc deficiencies in beans, tomatoes, etc. 100 lbs. \$9.90; less than 100 lbs. 13c per lb.

# HORMONES

**HORMODIN POWDERS.** Hormodin is a hormone-like plant growth substance. Its principal use is to stimulate rapid growth on cuttings. It has wide-spread use in propagating cuttings, whether rooted with ease, or normally rooted with difficulty. Many cuttings formerly impossible to propagate may now be rooted with the aid of Hormodin. Easy to use, consisting of simply dropping moistened stems in the powder and then planting. The cost of treatment is only a fraction of a cent per cutting.

No. 1—For Chrysanthemums, Poinsettias, Roses, etc. ¼ oz. 25c; 1¾ oz. 50c; 1 lb. \$3.00.

No. 2—For semi-woody plants and most common shrubs. 1¾ oz. 75c; 1 lb. \$4.50.

No. 3—For resistant species, such as Evergreens and dormant leafless cuttings. 1 oz. \$1.00; ½ lb. \$4.50.  
Combination Package, powders Nos. 1, 2 and 3, in carton, 75c.

**FRUITONE** is a wettable powder containing several hormones plus an effective wetting agent for securing a thorough coverage. Fruitone when sprayed on plant blossoms and young set fruit will help to prevent premature dropping. Cut flowers and twigs dipped in Fruitone will hold up longer.

Use 1 level teaspoonful to 1 gal.; 2/5-oz. to 5 gals.; 2 oz. to 25 gals.; 8 oz. to 100 gals. of water.

Price: ¾-oz. pkg. 25c; 2-oz. can \$1.00; 12-oz. can \$5.00.

**ROOTONE.** A hormone powder to hasten the rooting of cuttings. It is easy, economical, safe and sure treatment for rooting cuttings. Low cost—1/30 to 1/50 cent per cutting.

¼ oz. 25c; 2 oz. \$1.00; 1 lb. \$5.00; prepaid.

Rootone No. 10 recommended especially for rooting azaleas and camellias.

2 oz. jar \$2.00; 1 lb. jar \$10.00.

2 oz. pkg. will treat up to 3,000 cuttings.

**TRANSPLANTONE** supplies the plant with chemicals which stimulate root formation and growth after transplanting time. ½ oz. makes five gallons solution.

½ oz. 25c; 3-oz. can \$1.00; 1-lb. can \$4.00.



# SEED INOCULANTS

All prices quoted are postpaid.

When planting a legume feed or cover crop, such as clovers, cowpeas, beggarweed, crotalaria, soy beans and velvet beans, first inoculate the seed. This helps the crop to add the maximum amount of valuable nitrogen plant food material from the air to your soil. A



good leguminous cover crop can take 50 to 150 pounds of nitrogen per acre from the air—if inoculated. For your soil's sake, inoculate all legume seed.

|  |  |
|--|--|
| Group A—Alfalfa and Sweet Clover   | 1 bu. size 55c<br>100 lb. size 75c                   |
| Group B—Clovers<br>White Dutch Special   | 1 bu. size 55c<br>100 lb. size 75c<br>1 bu. size 65c |
| Group C—Vetch, Garden and Sweet Peas   | 100 lb. size 50c                                     |
| Group D—Beans—all varieties  | 1 bu. size 55c                                       |
| Group E—Peanuts, Cowpeas, Velvet Beans, Crotalaria, Beggarweed and Alyce Clover                                | 2 bu. size 40c                                       |
| Group L—Lespedeza  | 100 lb. size 55c                                     |
| Lupines—all varieties  | 100 lb. size 50c                                     |
| Group S—Soy Beans  | 2 bu. size 40c<br>5 bu. size 60c                     |
| Special Cultures: Lima Bean Trefoil, Dalea or Woods Clover, Crown Vetch, Chick Peas, Hairy Indigo and Sesbania | 1/8 bu. size 18c<br>1 bu. size 65c                   |
| Garden Size, for Garden Peas, Sweet Peas, All Beans, Soy Beans and Lupines.                                    | Each package will inoculate 6 lbs. seed—15c          |

## INOCULANT MADE IN FLORIDA ESPECIALLY FOR FLORIDA SOILS

|   | Quantity of seed will inoculate (size) | Price |
|---|--|-------|
| <b>Dixie Inoculant for</b>  |  |       |
| Soy Beans   | 5 bushels                              | .60   |
| Lespedeza, Alyce Clover Hairy Indigo                                | 100 pounds                             | .50   |
| Cowpeas, Crotalaria, Lima Beans, Peanuts and Velvet Beans           | 3 bushels                              | .50   |
| All Vetches, Austrian Peas, Wild Winter Peas, Garden and Sweet Peas | 100 pounds                             | .50   |
| Lupines, Special for Sweet, Yellow and Blue                         | 100 pounds                             | .50   |
| Clovers, Red Crimson, Ladino, White Dutch and Louisiana White       | 100 pounds                             | .75   |
| Alfalfa, Hubam, Floranna, Sweet and Bur Clovers                     | 100 pounds                             | .75   |
| Garden Size   | 8 pounds                               | .15   |



|   |                                     |
|---|-------------------------------------|
| Culture Group 1<br>Alfalfa, Bur, Button, Hubam Clovers                                      | 60 lb. size 55c<br>100 lb. size 75c |
| Culture Group 2<br>Clovers—Alsike, Crimson, Red, White, Ladino                              | 60 lb. size 55c<br>100 lb. size 75c |
| Culture Group 3<br>Vetches, English Peas  | 100 lb. size 50c                    |
| Culture Group 4<br>Beans—Snap, Pole, Navy, Pinto, Kidney                                    | 60 lb. size 55c                     |
| Culture Group 5<br>Lupines  | 100 lb. size 50c                    |
| Culture Group 6<br>Soy Beans  | 2 bu. size 40c<br>5 bu. size 60c    |
| Culture Group 7<br>Peanuts, Cowpeas, Velvet Beans, Crotalaria, Beggar Weed and Alyce Clover | 2 bu. size 40c<br>5 bu. size 60c    |

Garden size, inoculates up to 6 lbs. seed. Peas, Beans, Sweet Peas, Lupines, Soy Beans.....Price 15c

## POULTRY SUPPLIES

All prices f.o.b. Plant City or Kilgore Stores and subject to change without notice.

More complete line of poultry supplies carried in Kilgore stores.

### HUDSON HANDY FILL CHICK FOUNTAIN



|                              |             |
|------------------------------|-------------|
| No. A-4—Capacity one gallon  | .....\$1.10 |
| No. A-8—Capacity two gallons | ..... 1.40  |
| C-3—Capacity three gallons   | ..... 2.15  |
| C-5—Capacity five gallons    | ..... 2.90  |

### MASON JAR FOUNTAIN AND FEEDER

|   |             |
|---|-------------|
| B-35 Glass Pan with three point clip to fit Mason jar | .....\$ .15 |
|---|-------------|

### REEL TOP FEEDERS

|                                      |             |
|--------------------------------------|-------------|
| No. 118 Metal Feeder, 18-inch length | .....\$ .35 |
| No. 224 Metal Feeder, 24-inch length | ..... .55   |
| No. 336 Metal Feeder, 36-inch length | ..... .80   |

### HUDSON "KLEEN-EZY" REEL TOP FEEDERS

|  |             |
|--|-------------|
| No. 6360—36-in. length—Reel Top          | .....\$1.35 |
| No. 5480 Feeder, 48-inch length Reel Top | ..... 1.95  |

### GRILL TOP FEEDER

|  |             |
|--|-------------|
| No. 7480 Feeder, 48-in. length Grill Top | .....\$2.60 |
|--|-------------|

**Key Ring Leg Bands.** Colors, red, yellow, green, white and blue. Sizes for baby chick, pigeon, bantam, medium, intermediate and large. Specify size and color. 1c each; 75c per 100.

|                 | Each        | Doz.        |
|-----------------|-------------|-------------|
| Nest Eggs—china | .....\$ .05 | .....\$ .50 |

### THERMOMETERS

Black oxidized brass scales, white filled figures and graduations, mercury tubes.


|   |             |
|---|-------------|
| Incubator thermometer                     | .....\$1.50 |
| Brooder thermometer                       | ..... 1.25  |
| Weather thermometer, 7 in. wood case      | ..... 1.00  |
| Weather thermometer, 8½-in. Metal Utility | ..... 2.00  |



# KNAPSACK AND HAND DUSTERS

**THE CHAMPION NO. 100 DOUBLE ACTION DUSTER** is carried like a knapsack. It fits comfortably on the back and leaves both hands free for efficient operation. The easily operated bellows throws a screen of dust exactly where wanted within a range of 30 feet. The complete duster weighs only 14 pounds empty and will hold as much as 17 pounds of fluffy powder. Can be converted to single action by removing upper intake valve.


**Price \$27.95, Prepaid**



*Champion No. 100  
Double Action Duster*

## ROOT HAND GUN MODEL C-3A


Root Hand Duster C-3A  
with "Y" Attachment  
**\$34.00, Postpaid**



Root Hand Duster C-3B  
(Less "Y" Attachment)

**Price \$31.20, Postpaid**

## CHAMPION HAND DUSTER — MODEL 103



*Champion Hand  
Duster Model 103*


Simplicity of design and ease of operation make the Champion Hand duster ideal equipment for flowers, vegetables, shrubs and small trees. It is used extensively in home gardens and small estates as well as in experimental and show gardens; very efficient and economical for spot dusting small plants in the field such as watermelons, etc. It is of highest quality construction. All parts

are enclosed and the steel tank is lead coated for long life. It is extremely light in weight and will carry up to eight pounds of powder, although it operates effectively with only a fraction of an ounce of powder in the tank.

**Price \$9.85, Prepaid**

## HAND DUSTERS

Made of aluminum sheet and castings for lightest weight. The Hudson Roto-Power "806" is designed for growers who want the easiest turning and lightest carrying duster, one that produces the high velocity and volume of air necessary to break up dust and blast it right into the plant. Holds 14 lbs. average density dust. Applies from 5 to 45 lbs. of dust per acre. High velocity: 1 crank turn produces 60½ fan revolutions.



*Hudson No. 806 Roto-Power Duster (Aluminum)*

**Price \$39.80, Postpaid**

*f.o.b. Plant City or Kilgore stores*

## SMALL HAND DUSTERS

*f.o.b. Plant City*



|  |        |
|--|--------|
| Brown Crystal Duster, Glass Jar, capacity 1 quart..... | \$2.25 |
| Hudson No. 612-A Patrol Duster.....                    | 1.90   |
| Hudson No. 673 Cadet.....                              | .75    |
| Hudson Admiral No. 766-C.....                          | 2.40   |
| Hudson Adjutant No. 674.....                           | 1.25   |
| Root P-1 .....   | 2.50   |
| Root P-2 .....   | 3.50   |
| Dobbins No. 132, 1-lb. capacity, price .....           | 2.35   |



## HUDSON ADMIRAL No. 766-C

An ideal duster for the average home garden, for use on vegetables, flowers, trees, and shrubs.

## RESPIRATORS—

|   |             |
|---|-------------|
| Approved for parathion dusting .....                | each \$5.50 |
| Replacement cartridges, No. 11-A (4 to box).....    | Box 2.00    |
| Replacement filters, No. R-414 (50 to box).....     | Box 2.00    |
| No. R-435 (contains 4 cart. and 12 filters).....    | Box 2.50    |
| Monogoggles for eye protection with respirators.... | 2.80        |

*Postpaid*

## DOBBINS No. 310 Superbilt Flame Sprayer

Dobbins Flame Sprayer has many uses. Develops 2000° F. of easy portable heat in a flame that can be directed at the will of the operator. Can also be converted to a liquid sprayer by using spray attachment No. 225.

|   |         |
|---|---------|
| Dobbins No. 310, 4-gal. capacity Flame Sprayer, price ..... | \$24.80 |
| No. 225 Liquid Spray Attachment, price.....                 | 4.98    |




# POWER AND HAND SPRAYERS

Prices quoted are f.o.b. Plant City unless otherwise indicated

## HUDSON POWER SPRAYER NO. 43030 ILLUSTRATED

High pressure sprayer recommended for estates, small groves, dairy barn and cattle spraying, and landscape gardens. Handles all insecticides, fungicides, weed control chemicals, and white-wash.



|                              |          |
|------------------------------|----------|
| No. 43030, 30-gal. tank..... | \$297.50 |
| No. 43015, 15-gal. tank..... | 215.00   |

## COMPRESSED AIR SPRAYERS

### Prepaid

Compressed air sprayers are strong, efficient and easy to use. Ideal for dozens of spraying jobs in homes, truck gardens, dairy barns, poultry houses, etc.

|   |         |
|---|---------|
| Brown Compressed Air, Galv. Open-Head No. 64-D (4-gal.).....      | \$13.25 |
| Hudson Compressed Air Perfection Sprayer (4-gal.) No. 310-B ..... | 14.25   |
| Hudson Bugwiser No. 220-B (2½-gal.) .....                         | 7.35    |
| Hudson Leader No. 293-B (3½-gal.) .....                           | 9.85    |
| Hudson Climax No. 335A (3-gal.) .....                             | 12.65   |
| Dobbins 44-GS (4 gal.) .....                                      | 14.15   |
| Dobbins No. 21-G (3½-gal. capacity) .....                         | 9.90    |
| Dobbins No. 24-G Budget (3½-gal. capacity) .....                  | 7.90    |




Leader No. 293

## SPRAYER HOSE

|                             |     |
|-----------------------------|-----|
| ¾" Hose. Price per ft. .... | .20 |
|-----------------------------|-----|


## CHAMPION SPRAYER MODEL-1



by the operating lever. This lever also operates the brass agitator which keeps 4½ gallons of any liquid well mixed.

Price \$31.75, Postpaid

This all-purpose, portable sprayer is top in its field. The Champion combines in one unit all the desirable qualities of unusual power, simple operation, effective control, continuous agitation, and long life. Uniform pressure of 200 lbs. in the sturdy pressure cylinder is pumped as needed



Dobbins No. 702  
Hand Sprayer

## Household sprayers

f.o.b. Plant City or Kilgore stores

|   |        |
|---|--------|
| Brown No. 13, 1-quart capacity .....        | \$ .95 |
| Hudson No. 403, Misty, quart capacity ..... | .70    |
| Dobbins No. 702, quart capacity .....       | .60    |
| Dobbins No. 710, pint capacity .....        | .50    |

## Household continuous sprayers

f.o.b. Plant City or Kilgore stores

|   |        |
|---|--------|
| Brown No. 26 B.G., quart capacity.....            | \$1.90 |
| Hudson Cardinal No. 436, 3-quart capacity.....    | 2.40   |
| Hudson Mercury No. 433, 1-quart capacity.....     | 1.60   |
| Hudson Comet No. 431, 1-quart capacity.....       | 1.20   |
| Dobbins No. 43 G., 3-quart capacity.....          | 3.75   |
| Dobbins No. 703 Continuous, 1-quart capacity..... | 2.00   |
| Ken Sprayall (Trombone Type) No. 202.....         | 8.95   |
| No. 214 Redisprayer w/Extension .....             | 6.95   |
| Ken-Sprayall Little Giant No. 224 .....           | 4.95   |

**GAT SPRAYER.** Attaches to garden hose for dissolving, mixing and spraying Hy-Gro and other soluble plant foods.

|   |        |
|---|--------|
| Empty 2-lb. Hy-Gro jar with Gat Sprayer ..... | \$1.60 |
|---|--------|



## BRADSON SPRAY GUNS

For use with standard garden hose in obtaining pressure. Insect-O-Guns, Gard-N-Gun and Bug-Gun for spraying insecticides and fungicides.



Insect-O-Gun




Bug-Gun

Gro-Gun for application of soluble fertilizers and plant foods.


|                            |        |
|----------------------------|--------|
| Insect-O-Gun—Model-3 ..... | \$5.95 |
| Gard-N-Gun—Model-2 .....   | 3.95   |
| Gro-Gun—Model-20 .....     | 2.95   |
| Bug-Gun—Model-4 .....      | 2.65   |

## LAWN SPRINKLERS RAIN KING

|   |        |
|---|--------|
| Model-K Adjustable from 5 to 50 feet.....           | \$9.25 |
| Model-K-GSS., Rust resistant for coastal areas..... | 9.75   |
| Model-D Double nozzle adjustable sprinkler.....     | 6.50   |
| Model N1 Garden hose nozzle .....                   | 1.50   |
| Park Sprinkler Model 911.....                       | 2.95   |



Model K



Model D



# Miscellaneous Farm and Garden Tools


All prices quoted are f.o.b. Plant City or Kilgore stores



Planet Jr.  
No. 4 Hill  
and Drill  
Seeder

No. 4-D Seeder—Shipping weight 41 lbs. Price...\$27.95  
No. 4 Seeder—Shipping weight 50 lbs. Price... 34.95  
No. 300-A Seeder—Shipping weight 55 lbs. Price 36.95

Planet Jr. No. 219  
Fertilizer Distributor




No. 219 Fertilizer Distributor. Shipping weight 33 lbs. Price...\$21.50  
No. 215 Fertilizer Attachment for No. 219 Distributor. Shipping weight 17 lbs. Price ..... 10.95  
No. 8556X, Stand for No. 215. Shipping weight 2 lbs. Price.. 1.00  
No. 8858X, Covers for No. 215. Shipping weight 2 lbs. Price.. 1.40



Planet Jr.  
No. 220  
Fertilizer  
Distributor


Shipping  
Weight  
115 lbs.

No. 220 Fertilizer distributor.....\$57.50



Planet Jr.  
No. 13 Double  
and Single  
Wheel Hoe

No. 13 Double and Single Wheel Hoe—Shipping weight 27 lbs. Price.....\$17.50  
No. 92-D Harrow. Shipping weight 45 lbs. Price .....\$19.95  
90-D Plain with Lever. Shipping weight 54 lbs. Price ..... 23.95  
No. 4-D Five tooth plain Cultivator. Shipping weight 49 lbs. Price .... 18.95



Planet Jr.  
No. 92-D  
Harrow

Planet Jr. Grass Edger No. 2. Shipping wt. 3 1/4 lbs. Price .....\$2.95




Planet Jr.  
No. 17 1/2 Single  
Wheel Hoe

No. 17 Single Wheel Hoe. Shipping wt. 22 lbs. ....\$14.95

No. 19 Wheel Hoe. Shipping weight 18 lbs. Price ..... 9.75


No. 17 1/2 Single Wheel Hoe. Shipping wt. 18 lbs. ...\$13.50  
No. 119 High Wheel Plow. Shipping wt. 18 lbs. .... 9.50



BPI (2 HP) Planet  
Jr. Tractor with  
rubber tires  
Price \$190.00

B-12 cultivator attachment as illustrated. Shipping wt. 18 lbs. Price .....\$34.95

B-8 (2 1/2 h.p.) Tractor with 5.00 x 12 Rubber Tires. Shipping weight 197 lbs.....\$224.00  
Tuffy (1 h.p.) Shipping weight 90 lbs. .... 114.00  
H. W. (5 h.p.) Tractor with 5.50 x 16 Rubber Tires. Shipping weight 567 lbs. Price..... 400.00  
Write for Planet Jr. Tractor catalog and prices on tools and attachments for tractor.



Dobbins Corn Planter  
No. 925

## HAND CORN PLANTER


Dobbins all steel one handle automatic No. 925.....\$4.20

## CYCLONE SEEDER

The Cyclone will sow many acres per hour, adjustable to any amount per acre.

In ordering seeder to be used with power take-off specify make and model of tractor to be used on. Electric motor model operates from 6 volt battery from jeep or truck.

Hand Seeder .....\$ 4.30  
Power Model M-1 Electric Motor ..... 45.00  
Power Model S-1 for Power Take-Off ..... 57.00  
Extension Hoppers for M-1 and S-1—1 1/2 bu. capacity.....\$7.00



## INSCO Soil Fumigant Applicator


A 4-gal. tank gravity flow applicator for single row application. Can be attached to horse drawn plow or small tractors.

Price f.o.b. Plant City or Kilgore stores .....\$15.00



# MISCELLANEOUS FARM and GARDEN TOOLS

Prepaid unless specified. Prices subject to change without notice



ECLIPSE builds precision power mowers with extra features for unmatched performance in low-cost maintenance of beautiful lawns. Balanced design, durable construction, dependable power, self propelled, precise cutting action and effortless handling make lawn mowing easier, faster and better.

|  |          |
|--|----------|
| Eclipse Lark 18-in. ....                         | \$128.80 |
| Eclipse Rocket 20-in. (Lever starter) .....      | 146.50   |
| Eclipse Parkhound 21-in. ....                    | 164.90   |
| Eclipse Tornado No. 800, 36-in. Sickle Bar ..... | 235.00   |


Prices f.o.b. Plant City or Kilgore stores  
Sharpeners for reel type mowers available

|                                  |         |
|----------------------------------|---------|
| Eclipse—16-inch Hand Mower ..... | \$34.50 |
| Zephyr 16-inch .....             | 30.00   |
| Vogue 16-inch .....              | 27.00   |
| Arlington 16-inch .....          | 22.50   |

Prices f.o.b. Plant City or Kilgore stores

## NO. 811—3 PC. GARDEN SET

Sturdy, attractive hand made tools finished in baked green enamel. Set includes trowel, fork and cultivator.



|                            |              |
|----------------------------|--------------|
| No. 800 Trowel .....       | 25c each     |
| No. 801 Fork .....         | 25c each     |
| No. 802 Cultivator .....   | 25c each     |
| No. 804 Transplanter ..... | 25c each     |
| No. 805 Weed Cutter .....  | 25c each     |
| No. 811 3-piece Set .....  | .75c per set |

## WISS GARDEN TOOLS

Postpaid

### FLOWER SHEARS

|   |        |
|---|--------|
| FH-4 Cutter Holder, Chrome Plated 6½" ..... | \$2.25 |
|---|--------|




No. 808 Draw Cut Pruning Shears

|                                |        |
|--------------------------------|--------|
| No. 808 Draw Cut .....         | \$3.95 |
| No. 607-R Pruning Shears ..... | 2.75   |
| No. C-1 Orange Clippers .....  | 2.40   |



No. 8½-E Hedge Shears

|                             |      |
|-----------------------------|------|
| No. 9-A Hedge Shears .....  | 4.50 |
| No. 8½-E Hedge Shears ..... | 4.95 |
| No. 9½-E Hedge Shears ..... | 5.50 |



No. 224-T Lopping Shears

|                         |        |
|-------------------------|--------|
| No. 20-V 20 inch .....  | \$4.95 |
| No. 321-T 21 inch ..... | 5.95   |
| No. 224-T 24 inch ..... | 6.25   |




No. 5600 Forged Grass Shears

|  |        |
|--|--------|
| No. 5600 Forged Grass Shear .....      | \$2.75 |
| No. 500 Grass Shears .....             | 1.35   |
| No. 900 Grass Shears—Long Handle ..... | 4.25   |
| No. 801 "Quick Trim" .....             | 1.75   |
| No. 700 Grass Shear .....              | 1.95   |

## LAWN FERTILIZER DISTRIBUTORS

f.o.b. Plant City or Kilgore stores




### MASTERS MODEL 30

50-lb. capacity. All steel construction, tubular steel handle with two "bicycle type" rubber grips. Rubber tires. Finest home lawn spreader ever developed. Price \$12.95.

### MASTERS MODEL 25-D

20-lb. capacity all steel construction with tubular steel handle, rubber handle grip. Steel wheels with rubber tires. For the customers who want "something better" in a small spreader. Price \$10.25.



### MASTERS HANDI-CART MODEL 54

The Masters Handi-Cart has an outstanding reputation among users as a sturdy, dependable tool for hundreds of moving jobs around the home and garden. Capacity 3 cu. feet. Price \$9.50.



# Miscellaneous Farm and Garden Supplies

Prices subject to change without notice



## BIRD NEPONSET PAPER POTS Nutrient Treated

Made of tough waterproof paper, adapted for growing plants from seed to transplanting age. Can be sold in pots. Plants get no setback when transplanted from pots. Performs the major function of clay pots at a fraction of the cost.

## NEPONSET RED PAPER POTS

Neponset Red Paper pots are designed for shipping plants which have been grown in clay pots or in other containers. Shipping costs are lower by using Neponset Red

Pots because there is considerably less weight and less breakage, and the clay pots can be saved for further use.

|               | By mail postpaid<br>100 pots |        | f.o.b. Plant City<br>1000 pots |         |
|---------------|------------------------------|--------|--------------------------------|---------|
|               | Vita                         | Red    | Vita                           | Red     |
| 2 inch .....  | \$2.20                       | \$1.35 | \$14.75                        | \$ 8.65 |
| 2½ inch ..... | 2.40                         | 1.45   | 15.75                          | 8.80    |
| 3 inch .....  | 2.70                         | 1.55   | 17.70                          | 9.30    |
| 3½ inch ..... | 3.50                         | 1.95   | 23.15                          | 12.40   |
| 4 inch .....  | 3.95                         | 2.40   | 26.25                          | 15.00   |
| 5 inch .....  | 5.20                         | 2.95   | 34.25                          | 18.60   |
| 6 inch .....  | 6.75                         | 3.60   | 45.35                          | 22.90   |

## PLANT BANDS



PICK UP LIKE THIS



SQUEEZE AND PRESS  
CORNERS



SET IN PLACE



Postpaid per 100      f.o.b. Plant City per 1000

(Contains sufficient fertilizer for 10 weeks)

## Vita-Bands 10

|                          |        |        |
|--------------------------|--------|--------|
| 1¾ x 1¾ x 2½ inches..... | \$ .85 | \$4.10 |
| 2 x 2 x 2½ inches.....   | .95    | 4.70   |
| 2 x 2 x 3 inches.....    | 1.05   | 5.35   |
| 2½ x 2½ x 3 inches.....  | 1.20   | 6.20   |
| 3 x 3 x 3 inches.....    | 1.35   | 7.15   |
| 4 x 4 x 3 inches.....    | 1.70   | 9.35   |

## Vita Band-D

A nutrient treated band which disintegrates in the soil.

|                   | Postpaid<br>per 100 | f.o.b.<br>Plant City<br>per 1000 |
|-------------------|---------------------|----------------------------------|
| 1½ x 1½ x 2½..... | \$ .75              | \$3.90                           |
| 1¾ x 1¾ x 2½..... | .85                 | 4.20                             |
| 2 x 2 x 2½.....   | .95                 | 4.75                             |
| 2½ x 2½ x 3.....  | 1.20                | 6.25                             |
| 4 x 4 x 3.....    | 1.70                | 9.50                             |

## TOM THUMB FLATS

|                                     | Per 100 | f.o.b.<br>Per 1000 |
|-------------------------------------|---------|--------------------|
| No. 1 (for bands 1¾ x 1¾ x 2½) .... | \$5.10  | \$46.15            |
| No. 2 (for bands 2 x 2 x 2½) .....  | 5.65    | 51.30              |

# Terra-Lite Vermiculite



**TERRA-LITE (Vermiculite).** An excellent light weight medium for starting cuttings, seed in flats, paper pots or plant bands. Can be mixed with soil in seedbeds. Increases germination by reducing damping-off. Holds moisture for several days without rewatering.

4-qt. bag 39c; ½-bu. bag 89c;

2-bu. bag \$1.75; 4-cu. ft. bag \$2.35,

f.o.b. Plant City

Write for complete descriptive pamphlet on Terra-Lite.

## PEAT MOSS

Good to add humus and hold moisture in seedbeds. Also an excellent poultry litter.

Georgia Peat Humus, 1½ cu. ft. bag \$1.20 each  
5 cu. ft. bags \$3.50 each

## European Peat. Compressed in burlap

Wrapped bales 20" x 20" x 36" each \$4.65;

Bulk 10c per lb.

f.o.b. Plant City

## SPHAGNUM MOSS

Burlap Wrapped Bales 14" x 16" x 45" each \$4.25

Bulk - 40c per lb.

f.o.b. Plant City



## RUBBER STRIPS

For Budding and Grafting

Postpaid

| Size                        | Length | Width    | Approx.<br>pcs. to<br>the lb. | Price<br>per lb. | Price<br>per 5<br>lb. pkg. |
|-----------------------------|--------|----------|-------------------------------|------------------|----------------------------|
| No. 520                     | 5 in.  | 3-16 in. | 1400                          | \$2.25           | \$10.00                    |
| No. 620                     | 6 in.  | 3-16 in. | 1080                          | 2.25             | 10.00                      |
| No. 838                     | 8 in.  | 3-8 in.  | 450                           | 2.25             | 10.00                      |
| Less than 1 lb. 20c per oz. |        |          |                               |                  |                            |

## AIRWRAP

Coated plastic adapted to grove and plant propagation. Coated with nutrient and rooting hormones.

9" width—8 yd. Rolls..... 1.95

9" x 8" Price per Sheet.....\$ .06

9" width—Less than 8 yds. per yd..... .25

Write for quotations on larger quantities.

## AIRLAYERING ROOTING KIT

For plant propagation. Kit contains sphagnum moss, plastic wraps and tying strips. Complete instructions in box.

Kit complete (Prepaid) .....\$1.95



## GREEN THUMB GLOVE

Ladies' sizes, small, medium and large. Prepaid...\$ .89




# Miscellaneous Farm and Garden Supplies

Prices subject to change without notice

## 'TWIST-EMS'

Postpaid

Help you support plants quickly, neatly, permanently.



|                                 |     |
|---------------------------------|-----|
| Package of 35—8-inch size ..... | 10c |
| Package of 50—8-inch size ..... | 15c |
| Box of 100—8-inch size .....    | 25c |
| Box of 200—4-inch size .....    | 25c |
| Box of 100—16-inch size .....   | 50c |

### UTILITY TWIST-EMS

|                       |                |
|-----------------------|----------------|
| 1000 1/4" x 8" .....  | \$1.15 prepaid |
| 1000 1/4" x 16" ..... | 3.90 prepaid   |


### TRAIN-ETTS

New Weatherized trellis netting, ideal support for sweet peas and any vining flower or vegetable plants. Comes in three sizes. No. 1, 60 x 72 in. 65c; No. 2, 60 x 96 in. 85c; No. 3, 60 x 180 in. \$1.25, postpaid.

### BAMBOO CANE STAKES

Dyed green pencil-thickness

|  |         |
|--|---------|
| 3 ft. Stakes, per 1000 f.o.b. Plant City ..... | \$12.50 |
| 3 ft. Stakes, per 100 postpaid .....           | 1.50    |
| Less than 100 2 1/2c each postpaid.            |         |
| 4 ft. Stakes, per 1000 f.o.b. Plant City ..... | 16.00   |
| 4 ft. Stakes, per 100 postpaid .....           | 2.00    |
| Less than 100 3c each postpaid.                |         |



### LABELS—Plant and garden (postpaid)

|  | Per 100 | Per M  |
|--|---------|--------|
| 4" Pot Labels—Painted—(1M per ctn.) ....     | \$ .40  | \$3.25 |
| 5" Pot Labels—Painted—(1M per ctn.) ....     | .45     | 3.60   |
| 6" Pot Labels—Painted—(1M per ctn.) ....     | .50     | 4.00   |
| 8" Garden Labels—Painted—(250 per ctn.) ..   | .95     | 8.65   |
| 10" Garden Labels—Painted—(250 per ctn.) ..  | 1.05    | 9.60   |
| 12" Garden Labels—Painted—(100 per ctn.) ..  | 1.75    | 16.00  |
| 3 1/2" Copper Wired—Painted—(1M per ctn.) .. | .55     | 4.20   |

### GRAFTING WAX—Postpaid

A prepared wax recommended for all hand and brush grafting operations. 1/4 lb. 30c; 1/2 lb. 45c; 1 lb. 75c.

### GRAFTING TAPE

|   |        |
|---|--------|
| Grafting Tape 1/4-in. wide, 60-yd. Roll ..... | \$1.00 |
| Grafting Tape 1/2-in. wide, 60-yd. Roll ..... | 1.30   |
| Grafting Tape 1-in. wide, 60-yd. Roll .....   | 2.00   |

### BUDDING KNIVES—Postpaid

|   |        |
|---|--------|
| No. 11 Budding Knife, brass lining, 4 1/2 inch, Price | \$3.50 |
| No. 21SH Florist Knife, 2 blades, Price .....         | 3.00   |
| No. 8 1/2C Grafting Blade, Price .....                | 1.60   |
| No. 9C Budding Blade, Price .....                     | 1.60   |

### HOTKAPS

Hotkaps are "individual hothouses" for every plant—protecting them against frost, rain, wind, insects and crusting. They force plant growth, making hardier, healthier plants and producing crops two or three weeks earlier. They may be used with practically any crop that needs protection, but are especially suited for use with vine crops such as tomatoes, cucumbers, cantaloupes and watermelons.



|  |        |
|--|--------|
| 25 pkg. Complete with Paper Setter, each ..... | \$ .95 |
| 100 with Fibreboard Garden Setter .....        | 3.15   |
| 250 with Fibreboard Garden Setter .....        | 5.90   |
| 1000 without Setter .....                      | 18.35  |
| 5000 without Setter, per 1000 .....            | 18.20  |
| 10000 without Setter, per 1000 .....           | 18.05  |
| 25000 without Setter, per 1000 .....           | 17.90  |
| Fibreboard Garden Setter, alone .....          | .20    |
| Steel Setter, each .....                       | 1.95   |

Prices quoted are f.o.b. Plant City and all Kilgore stores

### LABEL PASTE

Ours is a genuine wheat paste, especially made for pasting labels on vegetables and fruit boxes. Mixes easily in cold or warm water. Will not discolor labels. Unaffected by water or atmospheric changes after drying.

Prices quoted are f.o.b. Plant City and all Kilgore stores  
25c per lb.; 100 lbs. \$22.00

### TWINE (Jute)

Soft vegetable 2 ply (app. 1520 ft. to lb.) tying twine, excellent quality. Used for tying tomato plants and bunching vegetables.

Per lb. 47c; 50 to 60-lb. reels, per lb. 45c.  
1/2-lb. balls.—30c each.

### COTTON TWINE (Variegated)

Used for tying pole beans. 2 1/2-lb. cones. Price per cone \$3.00.

f.o.b. Plant City

### RAFFIA

The plain or natural color comes in uneven weight hanks. 1 to 3 lbs. 80c lb.; 10 lbs. \$7.50. Prepaid.

### TARPAULINS

Write for prices stating size desired.

### SUDBURY SOIL

#### TEST KIT

##### Home Garden Model

Complete equipment with instructions and solutions for 20 tests, for Nitrogen, Phosphorous, Potash and Acidity.  
\$2.98 Postpaid



##### JR. PROFESSIONAL MODEL

Contains complete equipment and instructions—solutions for 50 tests for nitrogen, phosphorous, potash and acidity (pH). Transparent plastic case. Price \$5.89 Postpaid.

Two Minute Acidity Kit .....\$1.49


Write for prices on larger models.



# CANNING DEPARTMENT

Prices subject to change without notice

Florida offers the greatest opportunity for home canning of any state in the Union, due to the fact that every month in the year produces some fruit or vegetable, and yet there is probably less canning done here than in almost any other state; but we are becoming more thrifty each year in this respect.



## **BURPEE PRESSURE COOKER**


Pressure cooker or canner made by forming heavy plate aluminum. Very strong — half the weight of "cast" aluminum. Guaranteed not to crack, break or split. It will not stain because there are no "pores" in this metal to fill up with food and create a bad odor.

### **MODEL AR25**

Holds 10 qt. or 17 pt. cans or 7 qt. or 14 pt. jars. Equipped with canning basket, recipe book, and cooking pans.

Postpaid

Burpee AR25 Heavy Type Aluminum .....\$24.95



*Simple, accurate  
and foolproof.*

*Insert can, turn  
crank, and can  
is automatically  
sealed.*

*So simple!*

*Complete canning  
recipe book with  
each machine.*

## **DIXIE SIMPLIFIED AUTOMATIC HOME CAN SEALER**

Will seal No. 1, No. 2, No. 2½ and No. 3 Sanitary cans.  
Prepaid

With No. 2 or No. 3 chuck and base.....\$35.00  
Extra chuck and base ..... 8.50


Home canning is fine training for the children, both girls and boys, as it teaches thrift. In some states the women and children add several hundred dollars to the family income during the year from home canning, thus securing money for many other things. It matters not whether products are used at home or sold, it's money saved just the same.

Hardly is there a day in Florida when there is not some luscious vegetable or fruit that can easily be conserved for later use. Meats, poultry and fish can also be conserved by modern safe and scientific methods—same methods as used by commercial canners. A pantry full of wholesome home-canned products means a better balanced diet for the entire family the year round. It is fascinating, too, avoids waste and builds a bank account.

## **HOT-LIFT CAN TONGS**



Hot-Lift Can Tongs, price Postpaid .....\$ .60



## **BURPEE SIMPLEX SEALER WITH CAN SAVER ATTACHMENT**

*The Leader in Can Sealers*

Seals cans quickly and easily. Set the can in position, spin the crank with the right hand and move a lever with the left hand. Made of hardened steel for long life. Rustproof. Equipped with attachment which simplifies changing from one size can to another. Clamps on a table. Seals, opens and reflanges two sizes of cans, the No. 2 and No. 3. Tested canning recipes and illustrated instructions with every sealer. Net weight 11 lbs. Price \$15.90, prepaid.

## **BURPEE PRESSURE CANNER**


Pressure canner made by forming heavy plate aluminum. Very strong —half the weight of "cast" aluminum. Guaranteed not to crack, break or split. It will not stain because there are no "pores" in this metal to fill up with food and create a bad odor.

### **MODEL AR1525**

Holds 15 qt. or 25 pt. cans, or 14 qt. or 21 pt. jars. Equipped with 2 canning baskets, recipe book, and pans.

Postpaid

Burpee AR1525 Aluminum.....\$29.95





# CANNING DEPARTMENT

## SANITARY CANS



Our sanitary open top cans are ideal for home canning. The finest fruits or vegetables and all your hard work can be wasted if the cans are not right.

Our cans for home canning are uniform, accurate, dependable, and strong. For 30 years they have been recognized as cans of highest quality, preferred by the majority of home canners.

## CANS

*f.o.b. Plant City*

|  | No. in<br>Case | Case   | 1,000   |
|--|----------------|--------|---------|
| No. 2 Plain Sanitary cans and caps..           | 100            | \$4.30 | \$42.15 |
| No. 2 Plain Sanitary cans and caps..           | 24             | 1.15   | 45.25   |
| No. 3 Plain Sanitary cans and caps..           | 100            | 6.30   | 61.90   |
| No. 3 Plain Sanitary cans and caps..           | 24             | 1.65   | 66.05   |
| No. 2 Sanitary lacquered<br>cans and caps..... | 100            | 4.50   | 43.85   |
| No. 2 Lacquered cans and caps.....             | 24             | 1.15   | 46.95   |
| No. 3 Sanitary lacquered<br>cans and caps..... | 100            | 6.60   | 64.70   |
| No. 3 Lacquered cans and caps.....             | 24             | 1.70   | 68.90   |

These cans are of the highest quality, uniform, accurate, dependable, and strong.

Friction top or syrup cans are used for packaging syrup, honey and for many other purposes.



10-LB. SYRUP CAN

|  | No. in<br>Case | Case   | 1,000    |
|--|----------------|--------|----------|
| 1½ Friction-top cans and caps..  | 36             | \$2.10 | \$ 57.15 |
| No. 2½ Friction-top (approx.<br>¼ gal.) cans and caps.....   | 24             | 2.10   | 84.75    |
| No. 5 Friction-top (approx. ½<br>gal.) cans and caps .....   | 50             | 6.15   | 121.60   |
| No. 10 Friction-top (approx.<br>1 gal.) cans and caps .....  | 50             | 8.80   | 175.15   |
| 5 gal. Square Honey Cans—each 75c; case (16)   |                |        | \$10.55  |
| 50-lb. Lard Can—each 75c; case of 12—  |                |        | \$8.55   |
| Corrugated Shipping Cartons for 5-gal. Honey cans,<br>each 25c; 10 for \$2.30; 25 \$5.50; f.o.b. Plant City. |                |        |          |

## CANNING SUPPLIES

*(Prepaid)*

**Per 100**

|   |        |
|---|--------|
| Extra Covers for No. 2 Plain Sanitary cans.....   | \$ .80 |
| Extra Covers for No. 3 Plain Sanitary cans.....   | 1.05   |
| Extra Covers for No. 2 Enameled Sanitary cans.... | .90    |
| Extra Covers for No. 3 Enameled Sanitary cans.... | 1.15   |

## CANNING LABELS

*(Prepaid)*

Beans, Corn, Blackeye Peas, Tomato and Syrup.

|   | 25     | 100    | 1,000  |
|---|--------|--------|--------|
| Price, No. 1½ and 2½, Syrup....                   | \$ .25 | \$ .50 | \$3.50 |
| Price, No. 2 and 3, Vegetables....                | .30    | 1.00   | 5.25   |
| Price, No. 5 and No. 10, Syrup..                  | .25    | .60    | 5.00   |
| Rubber stamp, two-line \$1.75; three-line \$2.00. |        |        |        |

## ATLAS FRUIT JARS

*f.o.b. Plant City*

Atlas Jars are made of clear crystal glass of highest quality giving a natural color to packed fruits and vegetables. Double tested for strength. Best preserving jar on the market and one that is guaranteed to give entire satisfaction when properly used.



## Atlas Square Mason Jars

This style jar has been on the market for over 50 years but has recently been improved by the strong shoulder feature.

## FRUIT JARS

*(f.o.b. Plant City)*

|   | Per doz. |
|---|----------|
| Atlas Square Mason Jars,<br>½-pint size ..... | \$1.00   |
| Atlas Square Mason Jars, pint size..          | 1.05     |
| Atlas Square Mason Jars, quart size           | 1.15     |
| Atlas Square Mason, ½-gal. size.....          | 1.50     |
| Atlas Wide Mouth, pint size.....              | 1.15     |
| Atlas Wide Mouth, quart size.....             | 1.30     |

## Caps for all type jars

**Per doz.**

|  |        |
|--|--------|
| Atlas Seal All Standard Caps, 2-piece.....         | \$ .30 |
| Atlas Seal All Standard Lids only.....             | .15    |
| 63 MM 2-piece Caps for Coffee and Relish Jars..... | .30    |
| 63 MM Lids only for Coffee and Relish Jars.....    | .15    |
| Rubber Rings for Old Style Zinc Caps.....          | .10    |
| Wide Mouth Caps (Rings and Lids).....              | .45    |
| Wide Mouth (Lids Only).....                        | .25    |

## JELLY GLASS

|  |        |
|--|--------|
| No. 100 1/3 pt. size, Tumbler Style..... | \$ .70 |
| No. 25 1/3 pt. size, Ribbed Style.....   | .75    |
| No. 92 ½ pt. size, Tumbler Style.....    | .75    |
| No. 3 ½ pt. size, Ribbed Style.....      | .80    |



## HONEY JARS

*f.o.b. Plant City*

|                              | Utility<br>Round<br>per doz. | Queenline<br>per doz. |
|------------------------------|------------------------------|-----------------------|
| ½-lb., 2 doz. to case.....   | \$ .55                       | \$ .60                |
| 1-lb., 2 doz. to case.....   | .65                          | .80                   |
| 2-lb., 1 doz. to case.....   | .85                          | 1.00                  |
| No. 5 Glass Honey Pail.....  | 1.75                         | .....                 |
| No. 10 Glass Honey Pail..... | 2.00                         | .....                 |



QUEENLINE





# Bee-Keepers' Supplies, Bee-Hives and Equipment

Prices subject to change without notice and are f.o.b. Plant City or any Kilgore store. If you wish your shipment to be made by mail, please add postage when ordering. Most all Bee Supplies are exempt from State Sales Tax except Hive Bodies and Covers.

The following is only a partial list of our large and complete line of bee supplies carried in stock in all Kilgore's Florida stores and in Kilgore's mail order department at Plant City, Florida. We can always quickly get whatever a customer wants from the factory if we happen to be out of stock, in which case the price will be f.o.b. factory, Medina, Ohio. 5% discount will be allowed on orders of \$100.00 or more. Shipped direct from factory (average time of delivery two to three weeks).

We buy Bees Wax. Write for price.

Write for a complete catalog of Bee-keepers' supplies.

## Brood Foundation—(3-Ply)

8x16 $\frac{3}{4}$  in.—7 sheets to pound

|     |                 | Weight | Price  |
|-----|-----------------|--------|--------|
| H-1 | 10-Sheet Carton | 3 lbs. | \$1.98 |
| H-3 | 50-sheet Carton | 9 lbs. | 9.10   |

## Medium Brood

For standard frames

8x16 $\frac{3}{4}$  in.—8 sheets to pound

|      |                 | Weight | Price  |
|------|-----------------|--------|--------|
| H-20 | 10-sheet Carton | 2 lbs. | \$1.70 |
| H-22 | 50-sheet Carton | 8 lbs. | 7.98   |

## Special Brood Foundation

8x16 $\frac{3}{4}$  in.—10 sheets to pound

|      |                  | Weight  | Price   |
|------|------------------|---------|---------|
| H-32 | 10-sheet Carton  | 2 lbs.  | \$ 1.32 |
| H-34 | 50-sheet Carton  | 7 lbs.  | 6.15    |
| H-36 | 200-sheet Carton | 33 lbs. | 24.30   |

## Medium Brood

For shallow frames

4 $\frac{1}{2}$ x16 $\frac{3}{4}$  in.—16 sheets to pound

|      |                 | Weight | Price  |
|------|-----------------|--------|--------|
| H-26 | 10-sheet Carton | 2 lbs. | \$1.05 |
| H-28 | 50-sheet Carton | 5 lbs. | 4.60   |

## Medium Brood, Crimp-Wired

8 $\frac{1}{2}$ x16 $\frac{3}{4}$  in.—7 sheets to pound

|       |                 | Weight  | Price  |
|-------|-----------------|---------|--------|
| M-350 | 10-sheet Carton | 3 lbs.  | \$2.05 |
| M-350 | 50-sheet Carton | 10 lbs. | 9.20   |

## Medium Brood, Crimp-Wired

4 $\frac{3}{4}$ x16 $\frac{3}{4}$ —12 sheets to pound

|       |                 | Weight | Price  |
|-------|-----------------|--------|--------|
| M-352 | 10-sheet Carton | 2 lbs. | \$1.25 |
| M-352 | 50-sheet Carton | 6 lbs. | 5.60   |

## Thin Super

|       |  |        |        |
|-------|--|--------|--------|
| J-1.  | 1-lb. Carton, 3 $\frac{7}{8}$ x15 $\frac{3}{4}$ in.,<br>28 sheets to lb. | 2 lbs. | \$1.48 |
| J-3.  | 5-lb. Carton, 3 $\frac{7}{8}$ x15 $\frac{3}{4}$ in.,<br>28 sheets to lb. | 7 lbs. | 6.95   |
| J-22. | 1-lb. Carton, 4 $\frac{1}{2}$ x16 $\frac{1}{2}$ in.,<br>22 sheets to lb. | 2 lbs. | 1.48   |
| J-24. | 5-lb. Carton, 4 $\frac{1}{2}$ x16 $\frac{1}{2}$ in.,<br>22 sheets to lb. | 7 lbs. | 6.95   |
| J-27. | 1-lb. Carton, 5x16 $\frac{1}{2}$ in.,<br>22 sheets to lb.                | 2 lbs. | 1.48   |
| J-29. | 5-lb. Carton, 5x16 $\frac{1}{2}$ in.,<br>22 sheets to lb.                | 7 lbs. | 6.95   |

## Standard 10-Frame Hives

One-story standard hive with aluminum cover  
16 $\frac{1}{4}$ x20 in.

Hives in the flat are furnished with nails, 135-lb. tin rabbets punched for nails, reversible bottom boards, metal cover. Hoffman frames, and complete directions for nailing.

A-1. Knocked down without foundation. Wt. 31 lbs. 8.95

## One-story standard hive, with excelsior (wood) cover

A-6. Knocked down without foundation. Wt. 27 lbs. 7.35

## Shallow 10-Frame Extracting Supers.

|       |   |        |
|-------|---|--------|
| C-12. | 1 KD Empty only, no fittings. Wt. 5 lbs.  | \$1.55 |
| C-13. | 5 KD Empty only, no fittings. Wt. 28 lbs. | 6.95   |

## Beeway 10-Frame Section Supers

|      |   |        |
|------|---|--------|
| D-5. | 1 KD Without sections or foundation.<br>Wt. 7 lbs.  | \$2.60 |
| D-6. | 5 KD Without sections or foundation.<br>Wt. 36 lbs. | 11.95  |

## List of Inside Fixtures for Supers

|       |  |                          |        |
|-------|--|--------------------------|--------|
| E-1   | 70 Beeway Section Holders (4 $\frac{1}{4}$ x4 $\frac{1}{4}$ x1 $\frac{7}{8}$<br>in.) KD  | Wt. 3 lbs.               | \$ .90 |
| E-2.  | 50 Beeway Section Holders (4 $\frac{1}{4}$ x4 $\frac{1}{4}$ x1 $\frac{7}{8}$<br>in.), KD | Wt. 15 lbs.              | 5.00   |
| E-3.  | 100 Beeway Section Holders (4 $\frac{1}{4}$ x4 $\frac{1}{4}$ x1 $\frac{7}{8}$<br>in.) KD | Wt. 23 lbs.              | 9.95   |
| E-15. | 10 Beeway Wood Separators, 1/10 in.<br>thick   | Wt. 2 lbs.               | .70    |
| E-35. | 100 Super Springs (3 used to a<br>super)   | Wt. 1 $\frac{1}{2}$ lbs. | 1.55   |

## Covers, Metal, 10-Frame

|      |                                      |             |        |
|------|--------------------------------------|-------------|--------|
| B-1. | 1 KD Metal Cover with inner<br>cover | Wt. 11 lbs. | \$3.60 |
| B-2. | 5 KD Metal Cover with inner<br>cover | Wt. 51 lbs. | 15.75  |

## The Excelsior Wood Cover, 10-Frame

|      |                                  |             |        |
|------|----------------------------------|-------------|--------|
| B-4. | 5 KD Excelsior All-Wood<br>Cover | Wt. 31 lbs. | \$8.35 |
|------|----------------------------------|-------------|--------|

## Inner Covers, 10-Frame


|      |   |             |        |
|------|---|-------------|--------|
| B-8. | 1 KD Inner Cover, without bee<br>escape | Wt. 3 lbs.  | \$ .98 |
| B-9. | 5 KD Inner Cover, without bee<br>escape | Wt. 12 lbs. | 4.25   |

## Bottom Boards, 10-Frame

|       |                                 |             |        |
|-------|---------------------------------|-------------|--------|
| B-12. | 1 KD Reversible Bottom<br>Board | Wt. 7 lbs.  | \$1.95 |
| B-13. | 5 KD Reversible Bottom<br>Board | Wt. 37 lbs. | 8.35   |

## Comb Honey Sections

|     |          |             |         |
|-----|----------|-------------|---------|
|     | Mill Run |             |         |
| F-1 | 100      | Wt. 7 lbs.  | \$ 3.45 |
| F-3 | 500      | Wt. 32 lbs. | 15.45   |



Full Depth 10-Frame Extracting Supers or Hive Bodies



## BEE-KEEPERS' SUPPLIES (Continued)

For information on Beekeeping in Florida, write the Florida Agricultural Experiment Station, Mailing Department, Gainesville, Florida, for free copy of Bulletin No. 151, entitled Beginning Beekeeping

This publication is illustrated and contains much valuable information for beginners.

### Frames


Furnished in the flat with nails. All frames are furnished with solid bottoms unless otherwise specified.

|  | 10     | 50     | 100     |
|--|--------|--------|---------|
| Standard frames with heavy top bars and solid bottom bar, 9½x17½ in. | G-1    | G-2    | G-6     |
| Wt. 46 lbs. per 100  | \$1.85 | \$7.80 | \$14.95 |

|  |        |        |         |
|--|--------|--------|---------|
| Standard frames. Heavy top bar. 2 place bottom bar. Wt. 45 lbs., per 100 | G-14   | G-15   | G-16    |
|  | \$1.85 | \$7.80 | \$14.95 |

|  |        |         |
|--|--------|---------|
| Shallow Extracting Frames, with regular Hoffman top bar, with solid bottom bar. Wt. 38 lbs., per 100 | G-25   | G-26    |
|  | \$6.95 | \$13.70 |

|   |        |        |
|---|--------|--------|
| Shallow frames, L. T. thick top 2 piece bottom bars | G-24   | G-27   |
|   | \$1.80 | \$6.95 |




L 15

### Honey Extractors

|      |  | Wt. |          |
|------|--|-----|----------|
| L-15 | 1 2-frame reversing (9½x16 in. pocket) | 100 | \$ 65.90 |
| L-20 | 1 20-frame Simplicity                  | 300 | 199.50   |
| L-25 | 1 4-frame reversing (9½x16 in. pocket) | 215 | 142.00   |
| L-35 | 1 3-frame Novice, hand power           | 52  | 34.50    |
| L-45 | 1 45-frame Simplicity                  | 470 | 318.00   |

Large Electric Extractor can be supplied upon order



### Bee Escape

|     |                                   |        |
|-----|-----------------------------------|--------|
| O-5 | Root Double Escape without Board. |        |
|     | Wt. 1 oz.                         | \$ .35 |


### Queen Excluder, 10-Frame

Bees pass through rapidly, without hesitation.

|  | Each   | 10      |
|--|--------|---------|
| 10-frame Standard 3 wire and wood, 16¼x20 in. Wt. 4 lbs. | B-30   | B-31    |
|  | \$1.60 | \$15.25 |
| 10-frame Special 7-wire and wood. Wt. 4 lbs.             | B-35   | B-36    |
|  | \$1.65 | \$16.25 |
| Unbound Zinc Excluders                                   | B-40   | B-41    |
|  | \$1.30 | \$12.25 |

### Wire for Frames

|       |                                   | Wt.   |        |
|-------|-----------------------------------|-------|--------|
| G-61. | ¼-lb. Spool Tinned Wire, 350 ft.  | 8 oz. | \$ .30 |
| G-62. | ½-lb. Spool Tinned Wire, 700 ft.  | 1 lb. | .50    |
| G-63. | 1-lb. Spool Tinned Wire, 1400 ft. | 2 lb. | .80    |
| G-64. | 5-lb. Spool Tinned Wire, 7000 ft. | 7 lb. | 3.75   |
| N-65. | Spur Wire Imbedder                | 8 oz. | .42    |



### Bristle Bee Brush

|      |                              |        |
|------|------------------------------|--------|
| N-85 | Bristle Bee Brush. Wt. 4 oz. | \$ .65 |
|------|------------------------------|--------|

### Steel Hive Tool

|       |                               |        |
|-------|-------------------------------|--------|
| N-81. | Hive Tool. 10-inch. Wt. 6 oz. | \$1.15 |
|-------|-------------------------------|--------|

### Queen and Drone Trap (10-frame)

|      |                       |        |
|------|-----------------------|--------|
| O-8. | Wire Trap. Wt. 2 lbs. | \$1.35 |
|------|-----------------------|--------|

### Feeders

|      |                                       |        |
|------|---------------------------------------|--------|
| O-2. | Boardman Feeder, with cap. Wt. 12 oz. | \$ .35 |
|------|---------------------------------------|--------|

### Standard Smoker

|      |  |        |
|------|--|--------|
| N-3. | Standard Tin Smoker, 3¼x7 in. Wt. 2 lbs. | \$2.35 |
|------|--|--------|

|      |                                  |      |
|------|----------------------------------|------|
| N-7. | Big Jumbo Tin Smoker. Wt. 3 lbs. | 3.80 |
|------|----------------------------------|------|



### Leather and Drill Bee Gloves

|                                |            |        |
|--------------------------------|------------|--------|
| Leather (medium or large)      | Wt. 2 lbs. | \$3.85 |
| Drill (small, medium or large) | Wt. 1 lb.  | 1.55   |



### Veils

|       |   |        |
|-------|---|--------|
| N-10. | Root Indestructible Bee Veil. Wt. 1 lb. | \$1.85 |
| N-12. | Delphos Steel Bee Veil. Wt. 1 lb.       | 2.35   |
| N-15. | Cotton Tulle Veil. Wt. 3 lbs.           | 1.40   |



# INDEX

|   | Page |
|---|------|
| <b>SEED AND PLANTING INFORMATION</b>                                      |      |
| Home Gardens for Better Health  | 2    |
| Important Directions to Customers   | 1    |
| Kilgore Stores, Locations   | 49   |
| Location and Plan of the Vegetable Garden                                 | 3    |
| Making a Seedbed  | 3    |
| Ordering and Mailing Information  | 1    |
| Preparation of the Soil for the Vegetable Garden                          | 3    |
| Reference Table for Florida Vegetable Garden                              | 4    |
| Sales Tax Information   | 2    |
| Storing and Handling Seed   | 34   |
| What Corn, Grass, Clover and Field Crops and When to Sow Seeds in Florida | 5    |
| What Vegetables and When to Sow Seeds in Florida                          | 5    |
| What Flowers and When to Sow Seeds in Florida                             | 5    |
| What Will Your Harvest Be?  | 34   |

## SEED

|                                |       |
|--------------------------------|-------|
| Alfalfa                        | 41    |
| Beans, Bush, Pole, Lima, Green | 6-8   |
| Beans, Velvet                  | 46    |
| Beets                          | 9     |
| Beets, Spinach                 | 9     |
| Beets, Sugar and Stock         | 9     |
| Bene or Sesame                 | 41    |
| Broccoli                       | 9     |
| Brussels Sprouts               | 9     |
| Cabbage                        | 10-11 |
| Cabbage, Chinese               | 11    |
| Cantaloupe                     | 12    |
| Carrots                        | 13    |
| Cauliflower                    | 12    |
| Celery                         | 14-15 |
| Chicken Corn                   | 43    |
| Chufas                         | 43    |
| Clover, Alyce                  | 41    |
| Clover, White Dutch            | 42    |
| Clover, Hubam                  | 41    |
| Clover, Floranna               | 41    |
| Clover, Kenland Red            | 42    |
| Clover, Ladino                 | 42    |
| Collard                        | 15    |
| Corn, Field                    | 38-39 |
| Corn, Roasting Ear             | 39    |
| Corn, Sweet                    | 18    |
| Cowpeas, Field and Table       | 42    |
| Crotalaria                     | 43    |
| Cucumber                       | 16-17 |
| Dill                           | 20    |
| Eggplant                       | 19    |
| Egyptian Wheat                 | 43    |
| Endive                         | 20    |
| Flower Seed                    | 48    |
| Field Seed                     | 41-46 |
| Garlic                         | 20    |
| Grass, Lawn                    | 39-40 |
| Grass, Pasture                 | 41    |
| Gladiolus Bulbs                | 47    |
| Hegari (Higear)                | 46    |
| Hairy Indigo                   | 43    |
| Kale                           | 20    |
| Kohl Rabi                      | 20    |
| Leek                           | 20    |
| Lettuce                        | 21    |
| Lupine                         | 43-44 |
| Millet, Pearl or Cattail       | 44    |
| Mustard                        | 22    |
| Oats                           | 44    |
| Okra                           | 23    |
| Onion                          | 24    |
| Onion Sets                     | 24    |
| Parsley                        | 25    |
| Parsnip                        | 25    |
| Peanuts                        | 44-45 |
| Peas, English or Garden        | 25    |
| Peas, Southern (Cowpeas)       | 42    |
| Pepper                         | 26-27 |
| Potatoes                       | 28    |
| Pumpkin                        | 28    |
| Radish                         | 29    |
| Rape, Dwarf Essex              | 45    |
| Rhubarb                        | 28    |
| Rice                           | 45    |
| Rose Bushes                    | 47    |
| Roselle                        | 28    |
| Rutabaga                       | 35    |
| Rye                            | 45    |
| Sage                           | 29    |

|                   |       |
|-------------------|-------|
| Sesbania          | 46    |
| Shallu            | 43    |
| Sorghum           | 46    |
| Spinach           | 29    |
| Squash            | 30-31 |
| Strawberry Plants | 47    |
| Sunflower         | 46    |
| Swiss Chard       | 9     |
| Tomato            | 32-33 |
| Turnip            | 35    |
| Watermelon        | 36-37 |
| Wheat, Egyptian   | 43    |

## INSECTICIDE AND FUNGICIDE INFORMATION

|   |       |
|---|-------|
| INTRODUCTION                                    | 50    |
| SANITATION                                      | 50    |
| SAFE COMBINATION OF PESTICIDES                  | 51    |
| DON'TS (PESTICIDES NOT TO USE ON CERTAIN CROPS) | 51    |
| SEED TREATMENT                                  | 52    |
| PESTICIDE DILUTION TABLE                        | 52    |
| STANDARD HOUSEHOLD MEASUREMENTS                 | 52    |
| AMT. OF SPRAY AND DUST PER ACRE                 | 52    |
| INSECT AND DISEASE RECOMMENDATIONS              | 53-58 |
| Insects   | 53-58 |
| Livestock Pests                                 | 58    |
| Termite Control                                 | 58    |

## INSECTICIDES 59-60

|                                   |    |
|-----------------------------------|----|
| Alderin 2 Spray                   | 59 |
| Aramite 50W                       | 59 |
| Arsenate of Lead                  | 59 |
| BENZENE HEXACHLORIDE PREPARATIONS | 59 |
| LINDANE PREPARATIONS (ISOTOX)     | 59 |
| Black Leaf 40                     | 59 |
| Calcium Arsenate Powder           | 59 |

|                        |    |
|------------------------|----|
| CHLORDANE PREPARATIONS | 59 |
| Corn Earworm Oil       | 59 |

|                   |    |
|-------------------|----|
| DDT PREPARATIONS  | 59 |
| Dieldrin Emulsion | 59 |
| Karathane         | 59 |

|                       |    |
|-----------------------|----|
| KRYOCIDE PREPARATIONS | 60 |
| Malathion             | 60 |
| Lime Sulphur Solution | 60 |
| Oil Emulsion          | 60 |
| Ortho Garden Spray    | 60 |
| Para-Scalecide        | 60 |

|                        |    |
|------------------------|----|
| PARATHION PREPARATIONS | 60 |
| Paris Green            | 60 |
| POISON BAITS           | 60 |
| Red Arrow Insect Spray | 60 |

|                       |    |
|-----------------------|----|
| RHOTHANE PREPARATIONS | 60 |
| Rotenone Dust         | 60 |
| Scalecide             | 60 |

|                        |    |
|------------------------|----|
| TOXAPHENE PREPARATIONS | 60 |
| Vapotone XX Spray      | 60 |
| Volck Oil Spray        | 60 |

## FUNGICIDES 61-62

|         |    |
|---------|----|
| Arasan  | 61 |
| Ceresan | 61 |

## COPPER COMPOUNDS 61

|                                |    |
|--------------------------------|----|
| COPPER A COMPOUND PREPARATIONS | 61 |
| KILCOP "53" PREPARATIONS       | 61 |

## DITHANE PREPARATIONS 61

|                      |    |
|----------------------|----|
| FERMATE PREPARATIONS | 61 |
| Manzate              | 61 |
| Orthocide            | 61 |
| Parzate              | 61 |
| Phygon               | 61 |
| Semesan              | 61 |
| Semesan Bel          | 62 |
| Spergon              | 62 |
| Sulfocide            | 62 |
| Sulphur              | 62 |
| Tersan 75            | 62 |
| Zerlate              | 62 |

|   |    |
|---|----|
| INSECTICIDE and FUNGICIDE DUST COMBINATIONS | 62 |
|---|----|

## HOUSEHOLD and GARDEN PESTICIDES 63

|                    |     |
|--------------------|-----|
| FUMIGANTS          | 64* |
| Carbon Di Sulphide | 64  |
| Grain Protectant   | 64  |
| Cyanogas           | 64  |
| D-D Soil Fumigant  | 64  |
| Soilfume 80-20     | 64  |
| Soilfume Caps      | 64  |

## SPREADERS-STICKERS and EMULSIFIERS 64

|                             |    |
|-----------------------------|----|
| RODENTICIDES and VERMICIDES | 65 |
| Carbolineum                 | 65 |
| No-Nib'l Rabbit Repellent   | 65 |
| Ant Killers                 | 65 |
| Roach Killers               | 65 |
| Rat and Mouse Baits         | 65 |

## WEED KILLERS 65

## LIVE STOCK PREPARATIONS 66

## POULTRY PREPARATIONS 66

## FERTILIZERS 67

## SOLUBLE PLANT FOODS 67

## NUTRIENTS 68

|                        |    |
|------------------------|----|
| Activo                 | 68 |
| Adco                   | 68 |
| Es-Min-el              | 68 |
| Manganese Sulphate     | 68 |
| Kilgore's CMZ          | 68 |
| Sulphur-Manganese Dust | 68 |
| Kilgore Iron-Aid       | 68 |
| Aluminum Sulphate      | 68 |
| Hydrated Lime          | 68 |
| Zinc Sulphate          | 68 |

## HORMONES 68

|                 |    |
|-----------------|----|
| Hormodin Powder | 68 |
| Fruitone        | 68 |
| Rootone         | 68 |
| Transplantone   | 68 |

## SEED INOCULANTS 69

## POULTRY SUPPLIES 69

## KNAPSACK and HAND DUSTERS 70

## RESPIRATORS 70

## FLAME SPRAYERS 70

## POWER and HAND SPRAYERS 71

## LAWN SPRINKLERS 71

## FARM and GARDEN TOOLS 72

|                               |    |
|-------------------------------|----|
| Hand Corn Planter             | 72 |
| Cyclone Seeder                | 72 |
| Inso Soil Fumigant Applicator | 72 |

## LAWN and GARDEN TOOLS 73

## MICELLANEOUS FARM and GARDEN SUPPLIES 74-75

|                |    |
|----------------|----|
| Hot Kaps       | 75 |
| Label Paste    | 75 |
| Jute Twine     | 75 |
| Cotton Twine   | 75 |
| Raffia         | 75 |
| Soil Test Kits | 75 |

## CANNING DEPARTMENT 76-77

|                  |    |
|------------------|----|
| Pressure Cookers | 76 |
| Can Sealers      | 76 |
| Cans             | 77 |
| Fruit Jars       | 77 |
| Jelly Glasses    | 77 |
| Honey Jars       | 77 |

## BEE-KEEPERS SUPPLIES 78-79





## PLANT CITY, FLORIDA

Exp. or Frt. Office.....Date.....19.....

[SEE OTHER SIDE]



[illegible]

**DOES THIS ORDER** include everything you need in vegetable, flower and field crop seeds, fertilizers, sprays, dusts, spraying and dusting machines, garden tools, cans and canning equipment, poultry and bee supplies? Extra order blanks and envelopes sent upon application.

PLEASE WRITE BELOW

The names of neighbors or friends who might be interested in receiving a copy of our annual catalog.

| NAMES | POST OFFICE | P.O. Box, Street, or R.F.D. | STATE |
|-------|-------------|-----------------------------|-------|
|       |             |                             |       |
|       |             |                             |       |
|       |             |                             |       |
|       |             |                             |       |
|       |             |                             |       |

**Tear Off Along This Line**



Plant

**KILGORE'S**  
*Bred-Rite*  
**SEEDS**

TRADE MARK


*For Better Crops  
in  
Florida*



Early Prolific Straightneck  
Squash  
(See Page 31)




Florida Market  
(Phomopsis Resistant)  
Eggplant  
(See Page 19)



Cocozelle  
(Italian Vegetable Marrow)  
Squash  
(See Page 30)




Improved Rutgers Tomato  
(See Page 32)



Red Core Chantenay Carrot  
(See Page 13)



# KILGORE'S *Bred-Rite* SEEDS




Ocoee  
White Boston  
Lettuce  
(See Page 21)




Copenhagen  
Market  
(Early Strain No. 86)  
Cabbage  
(See Page 10)




Florida Giant Pepper  
(See Page 26)



Florida Giant  
(Black Diamond)  
Watermelon  
(See Page 37)



Shogoin  
(Foliage)  
Turnip  
(See Page 35)



Early  
Green Market  
(Marketer)  
Cucumber  
(See Page 16)



Fordhook Fancy Mustard  
(See Page 22)

## THE KILGORE SEED COMPANY

General Offices and Mail Order Department • • • PLANT CITY, FLORIDA